



**EUROPEAN COMMISSION  
Directorate General  
Joint Research Centre**

**IRMM**

# **CERTIFIED REFERENCE MATERIALS 2014**

Institute for Reference Materials and Measurements (IRMM)  
Standards for Innovation and sustainable Development Unit  
Retieseweg 111  
B - 2440 Geel, Belgium  
Fax: +32-(0)14-590 406  
Tel.: +32-(0)14-571 705  
e-mail: [jrc-irmm-rm-distribution@ec.europa.eu](mailto:jrc-irmm-rm-distribution@ec.europa.eu)  
Online catalogue:  
[http://irmm.jrc.ec.europa.eu/reference\\_materials\\_catalogue/catalogue/Pages/index.aspx](http://irmm.jrc.ec.europa.eu/reference_materials_catalogue/catalogue/Pages/index.aspx)  
Information on CRMs and on IRMM in general: <http://irmm.jrc.ec.europa.eu>



## TABLE OF CONTENTS

INTRODUCTION .....	v
<b>1 MATERIALS RELATED TO ENVIRONMENTAL ANALYSIS .....</b>	<b>1</b>
1.1 PURE MATERIALS AND SYNTHETIC MIXTURES .....	1
1.2 MATRIX MATERIALS .....	10
1.2.1 CERTIFIED FOR THE TOTAL ELEMENT CONTENT .....	10
1.2.2 CERTIFIED FOR THE EXTRACTABLE ELEMENT CONTENT AND SPECIES .....	19
1.2.3 CERTIFIED FOR ORGANIC POLLUTANTS.....	21
1.2.4 OTHERS.....	25
<b>2 MATERIALS RELATED TO THE ANALYSIS OF FOOD AND FEEDING STUFF...</b>	<b>25</b>
2.1 PURE MATERIALS AND SYNTHETIC MIXTURES .....	25
2.2 MATRIX MATERIALS .....	28
2.2.1 CERTIFIED FOR GMO CONTENT.....	28
2.2.2 CERTIFIED FOR NATURAL TOXINS AND XENOBIOTICS.....	36
2.2.3 CERTIFIED FOR THE TOTAL ELEMENT CONTENT .....	40
2.2.4 CERTIFIED FOR PROXIMATES AND CONVENTIONAL PROPERTIES.....	42
2.2.5 CERTIFIED FOR MICROBIOLOGICAL PROPERTIES .....	47
2.2.6 CERTIFIED FOR VETERINARY DRUGS.....	49
2.2.7 CERTIFIED FOR IDENTITY.....	51
2.2.8 OTHERS.....	52
<b>3 MATERIALS RELATED TO CLINICAL CHEMISTRY.....</b>	<b>52</b>
3.1 PURE STANDARDS AND SYNTHETIC MATERIALS .....	52
3.2 MATRIX MATERIALS .....	53
3.2.1 CERTIFIED FOR THE HORMONE CONTENT.....	533
3.2.2 CERTIFIED FOR THE TOTAL ELEMENT CONTENT AND OTHER PROPERTIES.....	544
3.2.3 CERTIFIED FOR PROTEIN CONTENT.....	55
3.2.4 CERTIFIED FOR CATALYTIC ACTIVITY .....	57
3.2.5 CERTIFIED FOR DNA SEQUENCE .....	58
3.2.6 OTHERS.....	58
<b>4 MATERIALS CERTIFIED FOR PHYSICAL PROPERTIES.....</b>	<b>59</b>
4.1 CERTIFIED FOR THERMAL PROPERTIES.....	59
4.2 CERTIFIED FOR MECHANICAL PROPERTIES .....	60
4.3 CERTIFIED FOR MORPHOLOGICAL PROPERTIES .....	62
4.4 CERTIFIED FOR OPTICAL PROPERTIES .....	64

<b>5 MATERIALS RELATED TO INDUSTRIAL APPLICATIONS .....</b>	<b>65</b>
5.1 CERTIFIED FOR COMPOSITION.....	65
5.2 CERTIFIED FOR TRACE ELEMENT CONTENT .....	67
5.3 OTHERS.....	72
<b>6 MATERIALS RELATED TO ISOTOPIC MEASUREMENTS.....</b>	<b>73</b>
6.1 CERTIFIED FOR ISOTOPE ABUNDANCE RATIO (AMOUNT RATIO).....	73
6.2 CERTIFIED FOR ISOTOPE AMOUNT CONTENT .....	75
<b>INDEX.....</b>	<b>77</b>
<b>NUMERICAL LIST .....</b>	<b>77</b>
<b>ALPHABETICAL LIST .....</b>	<b>88</b>

## INTRODUCTION

Public confidence in measurement results is important in many aspects of modern society, including consumer protection in food consumption, health-care, environmental protection, and fair trade. Certified Reference Materials (CRMs) are cornerstones of modern analytical quality assurance because they allow calibration of instruments, validation of methods, and quality control of methods and laboratories based on traceability and comparability of measurement results.

The Institute for Reference Materials and Measurements (IRMM) provides

**IRMM certified reference materials**, produced by the EC-JRC-IRMM

**BCR® certified reference materials** (BCR® is a registered trademark of the EC), for which production was supported by research funding of the European Commission, DG Research

and

**ERM® certified reference materials** (ERM® is a registered trademark of the EC), a new brand launched through the ERM® Initiative ([www.erm-crm.org](http://www.erm-crm.org)).

These CRMs are produced according to specific Guidelines of the European Commission which take into account the relevant ISO Guides 34 and 35.

The ERM® reference materials have undergone an additional uncompromising peer evaluation by the ERM® partners:

Bundesanstalt für Materialforschung und –prüfung (BAM), Germany  
LGC Standards, United Kingdom  
IRMM, JRC, European Commission

to guarantee highest quality and reliability.

## Certificates

Certificates carry a certified value with its uncertainty which is traceable either to a SI unit or an internationally accepted reference. The intended use for each CRM is stated on the certificate. CRMs are stored under controlled conditions which ensure their stability. Monitoring programmes have been set up to control CRM stability during the whole shelf-life.

At present IRMM's Reference Materials Unit offers about 550 different CRMs. A complete list of these CRMs, including description and price as well as the IRMM CRM catalogue and all certificates and certification reports can be accessed directly via the IRMM homepage:

<http://irmm.jrc.ec.europa.eu>

To order a free hardcopy of the catalogue simply write to:

[jrc-irmm-rm-distribution@ec.europa.eu](mailto:jrc-irmm-rm-distribution@ec.europa.eu)

## **Availability**

As CRMs should become regularly used items in any measurement laboratory, IRMM's Reference Materials Unit continues its effort to supply CRMs in sufficient amount to cover market needs. However, in exceptional cases where only limited amounts are available, it reserves the right to restrict and/or refuse orders. The reference materials contained in this catalogue are made available world-wide through the IRMM's Reference Materials Unit and its authorised distributors.

They can be purchased from the

European Commission, JRC,  
Institute for Reference Materials and Measurements (IRMM)  
Reference Materials Unit  
Retieseweg 111, B-2440 Geel, Belgium  
Tel. +32-(0)14-571 705, Fax +32-(0)14-590 406  
[jrc-irmm-rm-distribution@ec.europa.eu](mailto:jrc-irmm-rm-distribution@ec.europa.eu)

or from

### **AUTHORISED DISTRIBUTORS OF CRMs**

LGC STANDARDS GmbH  
Postfach 10 09 55  
D-46485 Wesel  
Germany  
Tel. +49 281 9887 0  
Fax +49 281 9887 299  
e-mail: [de@lgcstandards.com](mailto:de@lgcstandards.com)  
<http://www.lgcstandards.com>

SIGMA-ALDRICH CHEMIE GmbH  
Industriestrasse 25  
CH-9471 Buchs  
Switzerland  
Tel. +41 81 755 2828  
Fax +41 81 755 2815  
e-mail: [flukatec@sial.com](mailto:flukatec@sial.com)  
<http://www.sigmaaldrich.com/irmm>

Sigma-Aldrich RTC Inc.  
P.O. Box 1346, 2931 Soldier Springs Road  
Laramie, WY 82070  
USA  
Tel. +1 307 742 5452  
Fax +1 307 745 7936  
e-mail: [RTCSalesgroup@sial.com](mailto:RTCSalesgroup@sial.com)  
<http://www.RT-corp.com>

ARMI  
1 Perimeter Road, Suite 200  
Manchester, NH 03103  
USA  
Tel. +1 603 935 4100  
Fax +1 603 935 4101  
e-mail: [sales@armi.com](mailto:sales@armi.com)  
<http://www.armi.com>

Industrial Analytical  
Industrial Analytical House  
4 Indianapolis Road, Kyalami Business Park  
Kyalami 1684  
Republic of South Africa  
Tel. +27 11 466 4321  
Fax +27 11 466 4611  
e-mail: [info@industrialanalytical.co.za](mailto:info@industrialanalytical.co.za)  
<http://www.industrialanalytical.co.za>

Prices and sales conditions are freely determined by each distributor and can be obtained from them on request. They can provide advice and help in ordering and importing reference materials.

If samples are purchased through **UNAUTHORISED COMPANIES**, the IRMM cannot be held responsible for the integrity of the materials (especially in case of potentially unstable materials) nor, for the accuracy and/or completeness of the accompanying information (certificates, reports, etc.).

# 1 MATERIALS RELATED TO ENVIRONMENTAL ANALYSIS

## 1.1 PURE MATERIALS AND SYNTHETIC MIXTURES

	Substance	Purity (g/g)		
	<b>Polycyclic aromatic compounds</b>			
BCR-046	BENZO[b]CHRYSENE	0.994	+	0.006
		-		0.008
BCR-047	BENZO[b]FLUORANTHENE	0.997	4	0.002 6
BCR-048R	BENZO[k]FLUORANTHENE (unit size 10 mg)	0.997	+	0.003
		-		0.004
BCR-049	BENZO[j]FLUORANTHENE	0.997	±	0.003
		±		0.006
BCR-050	BENZO[e]PYRENE	0.991	+	0.009
		-		0.010
BCR-052	BENZO[ghi]PERYLENE	0.992	3	0.002 1
BCR-077R	1-METHYLCHRYSENE (unit size 10 mg)	0.991	+	0.007
BCR-078R	2-METHYLCHRYSENE (unit size 10 mg)	0.993	±	0.005
BCR-079R	3-METHYLCHRYSENE (unit size 10 mg)	0.993	±	0.005
BCR-080R	4-METHYLCHRYSENE (unit size 10 mg)	0.994	±	0.004
BCR-081R	5-METHYLCHRYSENE (unit size 10 mg)	0.997	3	0.001 3
BCR-082R	6-METHYLCHRYSENE (unit size 10 mg)	0.998	2	0.001 3
BCR-091	ANTHANTHRENE	0.996	±	0.004
BCR-092	10-AZABENZO[a]PYRENE	0.996	±	0.006
BCR-093R	1-METHYLBENZ[a]ANTHRACENE (unit size 10 mg)	0.996	±	0.005
BCR-094	DIBENZ[a,c]ANTHRACENE	0.996	±	0.004
BCR-095	DIBENZ[a,j]ANTHRACENE	0.997	8	0.002 5
BCR-096	DIBENZO[a,j]PYRENE	0.997	2	0.002 5
BCR-097	BENZO[a]FLUORANTHENE	0.996	±	0.004
BCR-133	DIBENZO[a,e]PYRENE	0.996	+	0.004
		-		0.005
BCR-134	BENZO[c]PHENANTHRENE	0.996	8	0.001 4
BCR-136R	BENZO[b]NAPHTHO[2,3-d]THIOPHENE (unit size 10 mg)	0.994	±	0.006
BCR-137R	BENZO[b]NAPHTHO[1,2-d]THIOPHENE (unit size 10 mg)	0.996	6	0.002 9
BCR-138	DIBENZ[a,h]ANTHRACENE	0.990	±	0.007
BCR-139	BENZO[ghi]FLUORANTHENE	0.995	±	0.004
BCR-140	BENZO[c]CHRYSENE	0.996	+	0.004
		-		0.005
BCR-153R	DIBENZ[a,h]ACRIDINE (unit size 10 mg)	0.999	2	0.000 6
BCR-154	DIBENZ[a,j]ACRIDINE	0.999	0	0.000 7
		-		0.001 0
BCR-155	DIBENZ[a,c]ACRIDINE	0.999	1	0.000 7
		-		0.000 8
BCR-156R	DIBENZ[c,h]ACRIDINE (unit size 10 mg)	0.993	6	0.002 1
BCR-157	BENZ[a]ACRIDINE	0.998	2	0.001 8
BCR-158	BENZ[c]ACRIDINE	0.998	7	0.001 3
		-		0.001 8
BCR-159	DIBENZO[a,h]PYRENE	0.993	±	0.007
BCR-160R	FLUORANTHENE (unit size 10 mg)	0.996	+	0.004
		-		0.005
BCR-168	PICENE (unit size 10 mg)	0.998	+	0.001 3
		-		0.004
BCR-177R	PYRENE (unit size 10 mg)	0.998	0	0.000 4

Availability: Amber vials containing about 100 mg of powdered material.

	Substance	Purity (g/g)		
	<b>Polycyclic aromatic compounds</b>			
BCR-152	DIBENZ[a,i]ACRIDINE	0.998	5	0.001 0
		-		0.000 8
BCR-265	DIBENZO[a,e]FLUORANTHENE	0.998	5	0.001 6
		-		0.001 0
BCR-266	7H-DIBENZO[c,g]CARBAZOLE	0.997	1	0.001 6
BCR-267	INDENO[1,2,3-cd]FLUORANTHENE	0.998	6	0.000 9
		-		0.000 8
BCR-269	CHRYSENE	0.992	8	0.002 8
BCR-270	TRIPHENYLENE	0.998	4	0.001 0
		-		0.000 6
BCR-271	BENZ[a]ANTHRACENE	0.998	4	0.000 9
BCR-272	CORONENE	0.998	9	0.000 6
		-		0.000 4

Availability: Amber vials containing about 20 mg of powdered material.

<b>Nitro-polycyclic aromatic hydrocarbons</b>		
BCR-305	1-NITROPYRENE	0.997 6 + 0.000 7
BCR-306	1-NITRONAPHTALENE	0.996 9 ± 0.001 0
BCR-307	2-NITRONAPHTALENE	0.997 7 + 0.000 9
BCR-308		- 0.001 1
BCR-309	9-NITROANTHRACENE	0.997 5 ± 0.001 0
BCR-310	6-NITROCHRYSENE	0.989 ± 0.004
	3-NITROFLUORANTHENE	0.996 8 + 0.001 2
BCR-311	6-NITROBENZO[a]PYRENE	- 0.002 1
BCR-312	2-NITRO-7-METHOXYNAPHTHO[2,1-b]FURAN	0.997 8 + 0.000 8 - 0.001 0
		0.998 4 ± 0.000 7

Availability: Amber vials containing about 10 mg of powdered material.

<b>Oxygenated polycyclic aromatic hydrocarbons</b>		
BCR-337	DIBENZO[b,d]FURAN	0.987 ± 0.007
BCR-338	4H-CYCLOPENTA[def]PHENANTHREN-4-ONE	0.995 1 ± 0.003 0
BCR-339	6H-BENZO[c,d]PYREN-6-ONE	0.988 ± 0.009
BCR-340	BENZO[b]NAPHTHO[1,2-d]FURAN	0.997 + 0.003
BCR-341		- 0.005
BCR-342	BENZO[b]NAPHTHO[2,1-d]FURAN	0.996 + 0.004 - 0.005
	BENZO[a]FLUORENONE	0.997 9 + 0.002 1 - 0.002 2

Availability: Amber vials containing about 10 mg of powdered material.

<b>Polychlorinated biphenyls</b>		
IUPAC No.		
BCR-289	8	2,4' - DICHLOROBIPHENYL
		0.996 3 + 0.005 - 0.001 8
BCR-290	20	2,3,3' - TRICHLOROBIPHENYL
BCR-291	28	2,4,4' - TRICHLOROBIPHENYL
BCR-293	52	2,2',5,5' - TETRACHLOROBIPHENYL
BCR-296	138	2,2',3,4,4',5' - HEXACHLOROBIPHENYL
BCR-297	153	2,2',4,4',5,5' - HEXACHLOROBIPHENYL
BCR-298	180	2,2',3,4,4',5,5' - HEPTACHLOROBIPHENYL
		0.995 7 ± 0.001 4

Availability: Amber vials containing about 25 mg of powdered material.

### **Polychlorinated biphenyls in iso-octane (BCR-365)**

IUPAC No.	Content in mg/kg	Concentration in g/m <sup>3</sup> at 25 °C <sup>1)</sup>
8	11.4 ± 0.4	(7.8 ± 0.2)
20	15.2 ± 0.9	(10.5 ± 0.7)
28	24.8 ± 1.1	(17.1 ± 0.8)
35	14.3 ± 0.8	(9.8 ± 0.5)
52	14.8 ± 0.6	(10.2 ± 0.4)
101	14.4 ± 0.6	(9.9 ± 0.4)
118	14.9 ± 0.8	(10.3 ± 0.6)
138	8.6 ± 0.6	(5.9 ± 0.5)
153	14.2 ± 0.6	(9.8 ± 0.4)
180	15.2 ± 0.6	(10.4 ± 0.3)

1) Not certified concentrations (g/m<sup>3</sup>) were calculated from the certified content assuming a density of iso-octane of 687.77 kg/m<sup>3</sup> at 25 °C.

Availability: Unit consisting of a pair of dark glass ampoules, each containing 2 cm<sup>3</sup> of 2,2,4-Trimethylpentane (iso-octane) sealed under nitrogen.

The pair of ampoules is supplied in a metal can which is packed with absorbent material.

## Polychlorodibenzo-p-dioxins (PCDDs) and polychlorodibenzofurans (PCDFs)

### BCR-614 SOLUTION – S0

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
2,3,7,8-T <sub>4</sub> CDD	0.137	0.004	0.098 3	0.002 9
1,2,3,7,8-P <sub>5</sub> CDD	0.698	0.014	0.501	0.010
1,2,3,4,7,8-HCDD	0.688	0.021	0.494	0.015
1,2,3,6,7,8-HCDD	0.696	0.006	0.500	0.004
1,2,3,7,8,9-HCDD	0.705	0.008	0.506	0.006
1,2,3,4,6,7,8-HCDD	1.400	0.020	1.005	0.014
1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	1.396	0.007	1.001	0.005
2,3,7,8-T <sub>4</sub> CDF	0.139 7	0.001 1	0.100 2	0.000 8
1,2,3,7,8-P <sub>5</sub> CDF	0.707	0.013	0.507	0.009
2,3,4,7,8-P <sub>5</sub> CDF	0.698	0.005	0.501	0.004
1,2,3,4,7,8-HCDF	0.700	0.006	0.502	0.005
1,2,3,6,7,8-HCDF	0.698	0.005	0.501	0.004
1,2,3,7,8,9-HCDF	0.699	0.009	0.502	0.007
2,3,4,6,7,8-HCDF	0.694	0.007	0.498	0.005
1,2,3,4,6,7,8-HCDF	1.396	0.008	1.001	0.006
1,2,3,4,7,8,9-HCDF	1.394	0.030	1.001	0.022
1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	1.397	0.024	1.002	0.017

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

### BCR-614 SOLUTION – S0

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDD	13.95	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDD	13.9	0.4	10.00	0.23
<sup>13</sup> C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDD	13.94	0.24	10.00	0.17
<sup>13</sup> C-1,2,3,7,8,9-HCDD	13.95	0.10	10.01	0.07
<sup>13</sup> C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	27.87	0.16	20.00	0.12
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDF	13.96	0.09	10.02	0.07
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDF	13.94	0.24	10.00	0.17
<sup>13</sup> C-2,3,4,7,8-P <sub>5</sub> CDF	13.95	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,4,7,8-HCDF	13.90	0.07	9.97	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDF	13.93	0.10	10.00	0.08
<sup>13</sup> C-1,2,3,7,8,9-HCDF	13.93	0.10	10.00	0.07
<sup>13</sup> C-2,3,4,6,7,8-HCDF	13.93	0.09	10.00	0.06
<sup>13</sup> C-1,2,3,4,6,7,8-HCDF	27.92	0.20	20.03	0.15
<sup>13</sup> C-1,2,3,4,7,8,9-HCDF	27.87	0.24	20.00	0.17
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	27.88	0.25	20.01	0.18
<sup>13</sup> C-1,2,3,4-T <sub>4</sub> CDD	13.94	0.08	10.00	0.06

**BCR-614 SOLUTION – S1**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
2,3,7,8-T <sub>4</sub> CDD	0.273	0.008	0.196	0.006
1,2,3,7,8-P <sub>5</sub> CDD	1.394	0.027	1.000	0.020
1,2,3,4,7,8-HCDD	1.37	0.05	0.986	0.030
1,2,3,6,7,8-HCDD	1.391	0.010	0.998	0.007
1,2,3,7,8,9-HCDD	1.408	0.015	1.011	0.011
1,2,3,4,6,7,8-HCDD	2.80	0.04	2.006	0.028
1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	2.787	0.010	2.000	0.007
2,3,7,8-T <sub>4</sub> CDF	0.279 0	0.002 1	0.200 2	0.001 5
1,2,3,7,8-P <sub>5</sub> CDF	1.412	0.025	1.013	0.018
2,3,4,7,8-P <sub>5</sub> CDF	1.395	0.008	1.001	0.006
1,2,3,4,7,8-HCDF	1.398	0.011	1.003	0.008
1,2,3,6,7,8-HCDF	1.393	0.009	1.000	0.006
1,2,3,7,8,9-HCDF	1.397	0.017	1.002	0.012
2,3,4,6,7,8-HCDF	1.387	0.012	0.995	0.009
1,2,3,4,6,7,8-HCDF	2.787	0.012	2.000	0.009
1,2,3,4,7,8,9-HCDF	2.78	0.06	2.00	0.05
1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	2.79	0.05	2.00	0.04
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDD	13.95	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDD	13.9	0.4	10.00	0.23
<sup>13</sup> C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
<sup>13</sup> C-1,2,3,7,8,9-HCDD	13.94	0.10	10.01	0.07
<sup>13</sup> C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	27.86	0.16	19.99	0.11
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDF	13.96	0.09	10.01	0.07
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDF	13.93	0.24	10.00	0.17
<sup>13</sup> C-2,3,4,7,8-P <sub>5</sub> CDF	13.94	0.06	10.00	0.05
<sup>13</sup> C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
<sup>13</sup> C-1,2,3,7,8,9-HCDF	13.92	0.10	9.99	0.07
<sup>13</sup> C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
<sup>13</sup> C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.14
<sup>13</sup> C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	27.87	0.25	20.00	0.18
<sup>13</sup> C-1,2,3,4-T <sub>4</sub> CDD	13.93	0.07	10.00	0.05

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

**BCR-614 SOLUTION – S2**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
2,3,7,8-T <sub>4</sub> CDD	1.09	0.04	0.785	0.023
1,2,3,7,8-P <sub>5</sub> CDD	5.57	0.11	4.00	0.08
1,2,3,4,7,8-HCDD	5.49	0.17	3.94	0.12
1,2,3,6,7,8-HCDD	5.56	0.04	3.992	0.027
1,2,3,7,8,9-HCDD	5.63	0.06	4.04	0.05
1,2,3,4,6,7,8-HCDD	11.18	0.16	8.02	0.11
1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	11.15	0.04	8.000	0.027
2,3,7,8-T <sub>4</sub> CDF	1.116	0.008	0.801	0.006
1,2,3,7,8-P <sub>5</sub> CDF	5.65	0.10	4.05	0.07
2,3,4,7,8-P <sub>5</sub> CDF	5.58	0.03	4.004	0.022
1,2,3,4,7,8-HCDF	5.59	0.05	4.01	0.04
1,2,3,6,7,8-HCDF	5.57	0.04	3.999	0.024
1,2,3,7,8,9-HCDF	5.59	0.07	4.01	0.05
2,3,4,6,7,8-HCDF	5.55	0.05	3.98	0.04
1,2,3,4,6,7,8-HCDF	11.15	0.05	8.00	0.04
1,2,3,4,7,8,9-HCDF	11.14	0.24	7.99	0.17
1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	11.16	0.19	8.01	0.14
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDD	13.95	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDD	13.9	0.4	10.00	0.23
<sup>13</sup> C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
<sup>13</sup> C-1,2,3,7,8,9-HCDD	13.94	0.10	10.01	0.07
<sup>13</sup> C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	27.86	0.16	19.99	0.11
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDF	13.96	0.09	10.01	0.07
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDF	13.93	0.24	10.00	0.17
<sup>13</sup> C-2,3,4,7,8-P <sub>5</sub> CDF	13.94	0.06	10.00	0.05
<sup>13</sup> C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
<sup>13</sup> C-1,2,3,7,8,9-HCDF	13.93	0.10	9.99	0.07
<sup>13</sup> C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
<sup>13</sup> C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.15
<sup>13</sup> C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	27.87	0.25	20.00	0.18
<sup>13</sup> C-1,2,3,4-T <sub>4</sub> CDD	13.93	0.08	10.00	0.06

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

**BCR-614 SOLUTION – S3**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
2,3,7,8-T <sub>4</sub> CDD	5.47	0.16	3.92	0.12
1,2,3,7,8-P <sub>5</sub> CDD	27.9	0.6	20.0	0.4
1,2,3,4,7,8-HCDD	27.5	0.9	19.7	0.6
1,2,3,6,7,8-HCDD	27.81	0.19	19.96	0.14
1,2,3,7,8,9-HCDD	28.17	0.30	20.21	0.21
1,2,3,4,6,7,8-HCDD	55.9	0.8	40.1	0.6
1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	55.74	0.19	40.00	0.14
2,3,7,8-T <sub>4</sub> CDF	5.58	0.04	4.003	0.029
1,2,3,7,8-P <sub>5</sub> CDF	28.2	0.5	20.3	0.4
2,3,4,7,8-P <sub>5</sub> CDF	27.90	0.16	20.02	0.11
1,2,3,4,7,8-HCDF	27.96	0.22	20.06	0.16
1,2,3,6,7,8-HCDF	27.87	0.17	20.00	0.12
1,2,3,7,8,9-HCDF	27.9	0.4	20.04	0.24
2,3,4,6,7,8-HCDF	27.73	0.23	19.90	0.17
1,2,3,4,6,7,8-HCDF	55.74	0.24	40.00	0.17
1,2,3,4,7,8,9-HCDF	55.7	1.2	40.0	0.9
1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	55.8	1.0	40.0	0.7
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDD	13.95	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDD	13.9	0.4	10.00	0.23
<sup>13</sup> C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
<sup>13</sup> C-1,2,3,7,8,9-HCDD	13.95	0.10	10.01	0.07
<sup>13</sup> C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	27.87	0.16	20.00	0.11
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDF	13.96	0.09	10.02	0.07
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDF	13.93	0.24	10.00	0.17
<sup>13</sup> C-2,3,4,7,8-P <sub>5</sub> CDF	13.94	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,4,7,8-HCDF	13.90	0.07	9.97	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDF	13.93	0.11	10.00	0.08
<sup>13</sup> C-1,2,3,7,8,9-HCDF	13.93	0.10	10.00	0.07
<sup>13</sup> C-2,3,4,6,7,8-HCDF	13.93	0.09	10.00	0.06
<sup>13</sup> C-1,2,3,4,6,7,8-HCDF	27.91	0.20	20.03	0.14
<sup>13</sup> C-1,2,3,4,7,8,9-HCDF	27.87	0.24	20.00	0.17
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	27.88	0.25	20.00	0.18
<sup>13</sup> C-1,2,3,4-T <sub>4</sub> CDD	13.93	0.07	10.00	0.05

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

**BCR-614 SOLUTION – S4**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
2,3,7,8-T <sub>4</sub> CDD	27.3	0.8	9.6	0.6
1,2,3,7,8-P <sub>5</sub> CDD	139.3	2.7	100.0	2.0
1,2,3,4,7,8-HCDD	137	5	98.6	3.0
1,2,3,6,7,8-HCDD	139.1	1.0	99.8	0.7
1,2,3,7,8,9-HCDD	140.8	1.5	101.1	1.1
1,2,3,4,6,7,8-HCDD	280	4	200.6	2.8
1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	278.7	1.0	200.0	0.7
2,3,7,8-T <sub>4</sub> CDF	27.89	0.21	20.02	0.15
1,2,3,7,8-P <sub>5</sub> CDF	141.2	2.5	101.3	1.8
2,3,4,7,8-P <sub>5</sub> CDF	139.5	0.8	100.1	0.6
1,2,3,4,7,8-HCDF	139.8	1.1	100.3	0.8
1,2,3,6,7,8-HCDF	139.3	0.9	100.0	0.6
1,2,3,7,8,9-HCDF	139.6	1.7	100.2	1.2
2,3,4,6,7,8-HCDF	138.7	1.2	99.5	0.9
1,2,3,4,6,7,8-HCDF	278.7	1.2	200.0	0.9
1,2,3,4,7,8,9-HCDF	278	6	200	5
1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	279	5	200	4
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDD	13.95	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDD	13.99	0.4	10.00	0.23
<sup>13</sup> C-1,2,3,4,7,8-HCDD	13.98	0.07	10.03	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
<sup>13</sup> C-1,2,3,7,8,9-HCDD	13.94	0.10	10.01	0.07
<sup>13</sup> C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	27.86	0.16	19.99	0.11
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDF	13.96	0.09	10.01	0.07
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDF	13.93	0.24	10.00	0.17
<sup>13</sup> C-2,3,4,7,8-P <sub>5</sub> CDF	13.94	0.06	10.00	0.05
<sup>13</sup> C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
<sup>13</sup> C-1,2,3,7,8,9-HCDF	13.92	0.10	9.99	0.07
<sup>13</sup> C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
<sup>13</sup> C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.15
<sup>13</sup> C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	27.87	0.25	20.00	0.18
<sup>13</sup> C-1,2,3,4-T <sub>4</sub> CDD	13.93	0.08	10.00	0.06

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

**BCR-614 SOLUTION – S5**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
2,3,7,8-T <sub>4</sub> CDD	109	4	78.5	2.3
1,2,3,7,8-P <sub>5</sub> CDD	557	11	400	8
1,2,3,4,7,8-HCDD	549	17	394	12
1,2,3,6,7,8-HCDD	556	4	399.1	2.7
1,2,3,7,8,9-HCDD	563	6	404	5
1,2,3,4,6,7,8-HCDD	1118	16	802	11
1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	1115	4	799.9	2.7
2,3,7,8-T <sub>4</sub> CDF	1116	0.8	80.1	0.6
1,2,3,7,8-P <sub>5</sub> CDF	565	0	405	7
2,3,4,7,8-P <sub>5</sub> CDF	558	3	400.4	2.2
1,2,3,4,7,8-HCDF	559	5	401	4
1,2,3,6,7,8-HCDF	557	4	399.9	2.4
1,2,3,7,8,9-HCDF	559	7	401	5
2,3,4,6,7,8-HCDF	555	5	398	4
1,2,3,4,6,7,8-HCDF	1115	5	800	4
1,2,3,4,7,8,9-HCDF	1114	24	799	17
1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	1116	19	801	14
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDD	13.95	0.06	10.01	0.05
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDD	13.9	0.4	10.00	0.23
<sup>13</sup> C-1,2,3,4,7,8-HCDD	13.98	0.7	10.03	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDD	13.93	0.24	10.00	0.17
<sup>13</sup> C-1,2,3,7,8,9-HCDD	13.95	0.10	10.01	0.07
<sup>13</sup> C-1,2,3,4,6,7,8-HCDD	27.9	0.6	20.0	0.4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	27.86	0.16	19.99	0.11
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDF	13.96	0.09	10.02	0.07
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDF	13.93	0.24	10.00	0.17
<sup>13</sup> C-2,3,4,7,8-P <sub>5</sub> CDF	13.94	0.06	10.00	0.05
<sup>13</sup> C-1,2,3,4,7,8-HCDF	13.89	0.07	9.97	0.05
<sup>13</sup> C-1,2,3,6,7,8-HCDF	13.93	0.11	9.99	0.08
<sup>13</sup> C-1,2,3,7,8,9-HCDF	13.93	0.10	9.99	0.07
<sup>13</sup> C-2,3,4,6,7,8-HCDF	13.93	0.09	9.99	0.06
<sup>13</sup> C-1,2,3,4,6,7,8-HCDF	27.90	0.20	20.02	0.15
<sup>13</sup> C-1,2,3,4,7,8,9-HCDF	27.86	0.24	19.99	0.17
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	27.87	0.25	20.00	0.18
<sup>13</sup> C-1,2,3,4-T <sub>4</sub> CDD	13.93	0.08	10.00	0.06

**BCR-614 SOLUTION – S6**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDF	139.3	2.3	100.0	1.7
<sup>13</sup> C-1,2,3,7,8,9-HCDF	139.4	0.9	100.0	0.7
<sup>13</sup> C-1,2,3,4,7,8,9-HCDF	278.7	2.4	200.0	1.7

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

**BCR-614 SOLUTION – S7**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDD	139.5	0.6	100.1	0.4
<sup>13</sup> C-1,2,3,7,8-P <sub>5</sub> CDD	139	4	99.9	2.4
<sup>13</sup> C-1,2,3,4,7,8-HCDD	139.8	0.7	100.3	0.5
<sup>13</sup> C-1,2,3,6,7,8-HCDD	139.3	2.4	100.0	1.7
<sup>13</sup> C-1,2,3,4,6,7,8-HCDD	279	6	200	4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDD	278.7	1.6	200.0	1.1
<sup>13</sup> C-2,3,7,8-T <sub>4</sub> CDF	139.5	0.9	100.1	0.6
<sup>13</sup> C-2,3,4,7,8-P <sub>5</sub> CDF	139.2	0.6	99.9	0.4
<sup>13</sup> C-1,2,3,4,7,8-HCDF	138.9	0.6	99.7	0.5
<sup>13</sup> C-1,2,3,6,7,8-HCDF	139.4	1.1	100.0	0.8
<sup>13</sup> C-2,3,4,6,7,8-HCDF	139.4	0.8	100.0	0.6
<sup>13</sup> C-1,2,3,4,6,7,8-HCDF	278.7	2.0	200.0	1.4
<sup>13</sup> C-1,2,3,4,6,7,8,9-O <sub>8</sub> CDF	278.7	2.5	200.0	1.8

**BCR-614 SOLUTION – S8**

Congener	Certified mass fraction <sup>1)</sup> (µg/kg)	Uncertainty <sup>2)</sup> (µg/kg)	Mass fraction expressed in concentration units <sup>3)</sup> (µg/L)	Uncertainty expressed in concentration units <sup>3)</sup> (µg/L)
<sup>13</sup> C-1,2,3,7,8,9-HCDD	558	4	400.5	2.7
<sup>13</sup> C-1,2,3,4-T <sub>4</sub> CDD	557.4	2.7	400.0	2.0

- 1) The certified mass fraction has been calculated from the purity of the individual PCDD/F compounds as assessed in a comprehensive study and the gravimetric preparation of the solution.
- 2) Uncertainties have been calculated by combining contributions from the purity study and the gravimetric preparation; details are given in the certification report.
- 3) Non-certified values.

Availability: BCR-614 Solutions S1-S7 contain about 1 mL solution and BCR-614 Solution S8 about 0.5 mL. The solutions are available individually or as set. A set consists of 11 ampoules, one of S2, S4-S8 and two of S1 and S3. In addition, the set contains an additional solution S9, which is not certified. This solution is intended as complementary tool for QA/QC purposes.

Substance	ERM-AC213		
	PAHs in Acetonitrile / Toluene (µg/g)		
Benz[a]anthracene	3.09	±	0.04
Chrysene	3.06	±	0.05
5-methylchrysene	3.08	±	0.07
Benzo[b]fluoranthene	3.05	±	0.05
Benzo[k]fluoranthene	3.06	±	0.08
Benzo[j]fluoranthene	3.05	±	0.10
Dibenz[a,h]anthracene	2.76	±	0.05
Benzo[ghi]perylene	3.07	±	0.05
Dibenzo[a,l]pyrene	2.85	±	0.10
Dibenzo[a,e]pyrene	2.97	±	0.10
Benzo[c]fluorene	(2.13	±	0.11)
Cyclopental[cd]pyrene	(2.96	±	0.12)
Dibenzo[a,i]pyrene	(2.37	±	0.15)

Values in brackets are not certified.

Availability: ERM-AC213 consists of 2 mL toluene containing 15 PAHs in an ampoule.

## 1.2 MATRIX MATERIALS

### 1.2.1 CERTIFIED FOR THE TOTAL ELEMENT CONTENT

Substance	BCR-142R Light sandy soil (mg/kg)			ERM-CC141 Loam soil (mg/kg)			BCR-143R Sewage sludge amended soil (mg/kg)		
As	0.34	±	0.04	9.9	±	1.5			
Cd	12.1	±	0.7	0.35	±	0.05	71.8	±	1.2
Co				8.5	±	0.5	12.3	±	0.3
Cr				86	±	8			
Cu	69.7	±	1.3	14.4	±	1.4	130.6	±	1.5
Hg	0.067	±	0.011	0.083	±	0.017	1.10	±	0.07
Mn	970	±	16	464	±	18	904	±	13
Ni	64.5	±	2.5	26.4	±	2.4	299	±	5
Pb	40.2	±	1.9	41	±	4	179.7	±	2.1
Zn	(101	±	6)	57	±	4	1055	±	14
Aqua regia soluble <sup>1)</sup>									
As				7.5	±	1.4			
Cd	0.249	±	0.010	0.25	±	0.04	72.0	±	1.8
Co	(10.2	±	0.6)	7.9	±	0.9	(11.8	±	1.0)
Cr				31	±	4	426	±	12
Cu	(69.8	±	1.0)	12.4	±	0.9	(128	±	7)
Hg				0.080	±	0.008	(1.10	±	0.06)
Mn	(800	±	50)	387	±	17	858	±	11
Ni	61.1	±	1.5	21.9	±	1.6	296	±	4
Pb	25.7	±	1.6	32.2	±	1.4	174	±	5
Zn	93.3	±	2.7	50	±	4	1063	±	16

Values in brackets are not certified.

Availability: Glass bottles containing about 50 g of powdered material; ERM-CC141 contains minimum 24 g.

<sup>1)</sup> Details of the analytical procedure to obtain the aqua regia soluble content of the elements are given in the certification report.

Substance	BCR-667 Estuarine sediment (mg/kg)			ERM-CC690 Calcareous soil (mg/kg)		
Br	(99.7	±	2.5)			
Cd	(0.67	±	0.11)			
Ce	56.7	±	2.5	49.1	±	2.5
Co	(23.0	±	1.3)			
Cr	(178	±	16)			
Cs	(7.8	±	0.7)			
Cu	(60	±	9)			
Dy	4.01	±	0.14	2.90	±	0.28
Er	2.35	±	0.15			
Eu	1.00	±	0.05			
Fe	(44800	±	1)			
Gd	4.41	±	0.12	3.2	±	0.4
Ho	0.80	±	0.06			
La	27.8	±	1.0	24.4	±	1.7
Lu	0.325	±	0.020			
Mn	(920	±	40)			
Nd	25.0	±	1.4	19.1	±	2.2
Ni	(128	±	9)			
Pb	(31.9	±	1.1)			
Pr	6.1	±	0.5			
Sb	(0.96	±	0.05)			
Sc	13.7	±	0.7	7.9	±	0.9
Se	(1.59	±	0.08)			
Sm	4.66	±	0.20	3.5	±	0.4
Ta	(0.876	±	0.017)			
Tb	0.682	±	0.017	0.50	±	0.07
Th	10.0	±	0.5	7.6	±	0.8
Tm	0.326	±	0.025	0.232	±	0.026
U	2.26	±	0.15	1.90	±	0.23
Yb	2.20	±	0.09	1.57	±	0.19
Zn	(175	±	13)			

Values in brackets are not certified.

Availability: BCR-667: Glass bottles containing about 40 g of powdered material.

ERM-CC690: Glass bottles containing about 70 g of powdered material. The report gives additional indicative values for As, Au, Co, Cr, Cs, Cu, Er, Eu, Fe, Hf, Ho, Lu, Ni, Pb, Pr, Sb, Ta, W, Y and Zn.

Substance	BCR-277R			BCR-280R			BCR-320R		
	Estuarine sediment (mg/kg)			Lake sediment (mg/kg)			Channel sediment (mg/kg)		
As	18.3	±	1.8	33.4	±	2.9	21.7	±	2.0
Cd	0.61	±	0.07	0.85	±	0.10	2.64	±	0.18
Co	22.5	±	1.4	16.8	±	0.9	9.7	±	0.6
Cr	188	±	14	126	±	7	59	±	4
Cu	63	±	7	53	±	6	46.3	±	2.9
Fe							25700	±	1300
Hg	0.128	±	0.017	1.46	±	0.20	0.85	±	0.09
Mn							910	±	50
Ni	130	±	8	69	±	5	27.1	±	2.2
Pb							85	±	5
Sc							5.2	±	0.4
Se	(0.58	±	0.11)	(0.46	±	0.09)	(0.96	±	0.18)
Sn	(6.5	±	1.8)	(9.5	±	1.7)	(9.4	±	1.7)
Th							5.3	±	0.4
Tl							0.65	±	0.08
U							1.56	±	0.20
V							46.5	±	2.8
Zn	178	±	20	224	±	25	318	±	20

Values in brackets are not certified.

Availability: Amber glass bottles containing 40 g of powder for BCR-277R and BCR-320R and 30 g for BCR-280R.

Substance	BCR-145R			BCR-146R			BCR-597		
	Sewage sludge (mixed origin) (mg/kg)			Sewage sludge (industrial origin) (mg/kg)			Sewage sludge (mg/kg)		
Cd	3.50	±	0.15	18.8	±	0.5			
Co	5.6	±	0.4	7.39	±	0.27			
Cr				196	±	7			
Cu	696	±	12	838	±	16			
Hg	2.01	±	0.22	8.6	±	0.4			
Mn	156	±	4	323	±	7			
Ni	247	±	7	70	±	5			
Pb	286	±	5	609	±	14			
Zn	2122	±	23	3060	±	60			
Aqua regia soluble <sup>1)</sup>									
Cd	(3.43	±	0.17)	18.4	±	0.4			
Co	(5.3	±	0.7)	6.5	±	0.4			
Cr	307	±	13	174	±	7			
Cu	707	±	9	831	±	16			
Hg	(1.99	±	0.08)	8.39	±	0.25			
Mn	(145	±	7)	298	±	9			
Ni	251	±	6	65.0	±	3.0			
Pb	282	±	9	583	±	17			
Zn	2140	±	50	3040	±	60			

Values in brackets are not certified.

Availability: Glass bottles containing about 50 g of powdered material for BCR-146R, 40 g for BCR-145R and BCR-597.

<sup>1)</sup> Details of the analytical procedure to obtain the aqua regia soluble content of the elements are given in the certification report.

Substance	ERM-CC580		
	Estuarine sediment (mg/kg)		
Total Hg	132	±	3
CH <sub>3</sub> Hg <sup>+</sup>	0.075	±	0.004

Availability: Glass bottles containing about 40 g powder.

Substance	BCR-038		
	Fly ash from pulverised coal (mg/kg)		
As	48.0	±	2.3
Cd	4.6	±	0.3
Co	53.8	±	1.9
Cr	192	±	10
Cu	176	±	9
Fe	$33.8 \times 10^3$	±	$0.7 \times 10^3$
Hg	2.10	±	0.15
Mn	479	±	16
Na	$3.74 \times 10^3$	±	$0.15 \times 10^3$
Pb	262	±	11
Zn	581	±	29

Availability: BCR-038 in ampoules containing about 5 g.

Substance	BCR-176R		
	Fly ash (mg/kg)		
As	54	±	5
Cd	226	±	19
Co	26.7	±	1.6
Cr	810	±	70
Cu	1050	±	70
Fe	13100	±	500
Hg	(1.60)	±	(0.23)
Mn	(730)	±	(50)
Ni	117	±	6
Pb	5000	±	500
Sb	850	±	50
Se	18.3	±	1.9
Tl	1.32	±	0.21
V	(35)	±	(6)
Zn	16800	±	400

Values in brackets are not certified.

The report gives additional indicative values for Ag, Au, Ba, Br, Ce, Cs, Eu, Hf, La, Na, Rb, Sc, Ta, Th and W.

Availability: Amber glass bottles containing about 40 g of powdered material.

Substance	BCR-723		
	Trace elements in road dust (µg/kg)		
Pd	6.1	±	1.9
Pt	81.3	±	2.5
Rh	12.8	±	1.3

Availability: Brown glass bottles with screw cap containing approximately 25 g of powder.

Substance	ERM-CZ120 Elements in fine dust (PM <sub>10</sub> -like) (mg/kg)		
As	7.1	±	0.7
Cd	0.90	±	0.22
Pb	113	±	17
Ni	58	±	7

Availability: Vial containing approximately 0.5 g of fine dust.

Substance	BCR-060 Lagarosiphon major (Aquatic plant) (mg/kg)		BCR-596 Trapa natans (Aquatic plant) (mg/kg)	
	Al	Cd	Cr	Cu
Al	4180	±	120	
Cd	2.20	±	0.10	
Cr				36.3
Cu	51.2	±	1.9	±
Hg	0.34	±	0.04	1.7
Mn	1760	±	60	
Pb	64	±	4	
Zn	313	±	8	

Availability: CRMs are provided in units of 25 g.

Substance	BCR-100 Beech leaves (g/kg)		
Al	0.435	±	0.004
Ca	5.30	±	0.05
Cl	1.49	±	0.06
Cr	0.008 0	±	0.000 6
K	9.94	±	0.20
Mg	0.878	±	0.017
N	26.29	±	0.25
P	1.55	±	0.04
S	2.69	±	0.04

Availability: BCR-100 is provided in units of 30 g.

Substance	BCR-129 Hay powder (g/kg)			BCR-402 White clover (mg/kg)		
As				0.093	±	0.010
Ca	6.40	±	0.10			
Co				0.178	±	0.008
I	$0.167 \times 10^{-3}$	±	$0.024 \times 10^{-3}$			
K	33.8	±	0.8			
Mg	1.45	±	0.04			
Mo				6.93	±	0.19
N	37.2	±	0.5			
P	2.36	±	0.07			
S	3.16	±	0.04			
Se				6.70	±	0.25
Zn	$32.1 \times 10^{-3}$	±	$1.7 \times 10^{-3}$			
Kjeldahl-N	34.2	±	0.4			

Availability: CRMs are provided in powder form in bottles containing approximately for BCR-129 30 g, BCR-402 25 g.

Note: BCR-402 was produced from white clover grown on a ground specially rich in selenium. This explains the high content of this element.

Substance	ERM-CD281 Rye grass (mg/kg)		
As	0.042	±	0.010
B	5.5	±	0.5
Ca		(6.3 g/kg)	
Cd	0.120	±	0.007
Cr	24.8	±	1.3
Cu	10.2	±	0.5
Fe		(0.18 g/kg)	
Hg	0.0164	±	0.0022
K		(34 g/kg)	
Mg		(1.6 g/kg)	
Mn	82	±	4
Mo	2.22	±	0.12
Na		(4.0 g/kg)	
Ni	15.2	±	0.6
P		(2.8 g/kg)	
Pb	1.67	±	0.11
S		(3.4 g/kg)	
Sb	0.042	±	0.007
Se	0.023	±	0.004
Si		(1.3 g/kg)	
Sn	0.062	±	0.011
Zn	30.5	±	1.1

Values in brackets are not certified.

Availability: Amber glass vial containing approximately 10 g.

Substance	BCR-414 Plankton (mg/kg)		
As	6.82	±	0.28
Cd	0.383	±	0.014
Co	(1.43	±	0.06)
Cr	23.8	±	1.2
Cu	29.5	±	1.3
Fe	(1.85	±	0.19
Hg	0.276	±	0.018
K	(7.55	±	0.17)
Mn	299	±	13
Mo	(1.35	±	0.20)
Ni	18.8	±	0.8
Pb	3.97	±	0.19
Sc	(0.54	±	0.02)
Se	1.75	±	0.10
Sr	(261	±	25)
V	8.10	±	0.18
Zn	111.6	±	2.5

Values in brackets are not certified.

Availability: CRM is provided in powder form in bottles containing approximately 5 g.

Substance	BCR-482 Lichen (mg/kg)			ERM-CD200 Bladderwrack ( <i>Fucus vesiculosus</i> ) (mg/kg)		
Al	1103	±	24			
As	0.85	±	0.07	55	±	4
Cd	0.56	±	0.02	0.95	±	0.06
Cr	4.12	±	0.15			
Cu	7.03	±	0.19	1.71	±	0.18
Hg	0.48	±	0.02	0.0186	±	0.0016
Ni	2.47	±	0.07			
Pb	40.9	±	1.4	0.51	±	0.06
Se				0.088	±	0.010
Zn	100.6	±	2.2	25.3	±	1.7

Availability: BCR-482 is provided in powder form in bottles containing approximately 15 g.

ERM-CD200 is provided in powder form in bottles containing approximately 5 g.

Substance	BCR-670 Lemna minor (Aquatic plant) (duck weed) (µg/kg)		
As	(1980	±	190)
Cd	(75.5	±	2.5)
Ce	990	±	40
Cr	(2050	±	100
Cs	(77	±	10)
Cu	(1820	±	300)
Dy	79	±	7
Er	44.0	±	2.8
Eu	23.2	±	1.5
Gd	98	±	8
Ho	15.8	±	1.8
La	487	±	20
Lu	6.3	±	0.5
Mo	(560	±	70)
Nd	473	±	15
Pb	(2060	±	120)
Pr	121	±	6
Sc	191	±	11
Sm	94	±	7
Tb	14.0	±	1.1
Th	159	±	18
Tm	5.7	±	0.7
U	82	±	8
Y	460	±	60
Yb	40	±	4
Zn	(24000	±	2100)

Values in brackets are not certified.

Availability: Glass bottles containing about 10 g of powdered material.

Substance	BCR-063R Skim milk powder (natural)		
Ca	g/kg	13.49	±
Cl	g/kg	9.94	±
Cu	mg/kg	0.602	±
Fe	mg/kg	2.32	±
I	mg/kg	0.81	±
K	g/kg	17.68	±
Mg	g/kg	1.263	±
N (total)	g/kg	62.3	±
Na	g/kg	4.37	±
P	g/kg	11.10	±
Pb	µg/kg	18.5	±
Zn	mg/kg	49.0	±

Availability: Glass bottles containing 50 g of powdered material.

Substance	ERM-CE278k Mussel tissue (mg/kg)		
As	6.7	±	0.4
Ag	(0.044	±	0.016)
Cd	0.336	±	0.025
Cr	0.73	±	0.22
Cu	5.98	±	0.27
Fe	161	±	8
Hg	0.071	±	0.007
I	1.4	±	0.4
Mn	4.88	±	0.24
Ni	0.69	±	0.15
Pb	2.18	±	0.18
Rb	2.46	±	0.16
Se	1.62	±	0.12
Sr	19.0	±	1.2
Zn	71	±	4

Values in brackets are not certified.

Availability: CRMs are provided in powder form in bottles containing approximately 8 g.

Substance	BCR-668 Mussel tissue (µg/kg)		
As	(7100	±	500)
Cd	(275	±	11)
Ce	89	±	7
Cr	(370	±	60)
Cs	(13.8	±	1.5)
Dy	8.9	±	0.6
Er	4.5	±	0.5
Eu	2.79	±	0.16
Gd	13.0	±	0.6
Ho	(1.8	±	0.6)
La	80	±	6
Lu	0.389	±	0.024
Mo	(1990	±	150)
Nd	54	±	4
Pr	12.3	±	1.1
Sc	(8.5	±	1.8)
Sm	11.2	±	0.8
Tb	1.62	±	0.12
Th	10.7	±	1.2
Tm	0.48	±	0.08
U	56	±	5
Y	59	±	5
Yb	(2.8	±	0.5)
Zn	(70700	±	400)

Values in brackets are not certified.

Availability: Glass bottles containing about 10 g of powdered material.

Substance	BCR-463 Tuna fish (mg/kg)			ERM-CE464 Tuna fish (mg/kg)		
Total Hg	2.85	±	0.16	5.24	±	0.10
CH <sub>3</sub> Hg <sup>+</sup>	3.04	±	0.16	5.50	±	0.17

Availability: Glass bottles containing about 15 g.

Substance	BCR-505 Trace elements in estuarine water (nmol/kg)			BCR-579 Coastal sea-water (ng/kg)		
Cd	0.80	±	0.04			
Co	(0.99	±	0.26)			
Cu	29.4	±	1.5			
Fe	(19	±	4)			
Hg				1.9	±	0.5
Ni	24.1	±	2.0			
Pb	(0.24	±	0.14)			
Zn	172	±	11			

Values in brackets are not certified.

Availability: BCR-505 is provided in 1 L polyethylene bottles and BCR-579 in 1 L glass bottles.

Substance	ERM-CA408 Simulated rainwater (low contents) (mg/L)		
<b>Mass concentration:</b>			
Ammonium	0.910	±	0.028
Cl	1.96	±	0.07
Fluoride	0.194	±	0.008
Mg	0.145	±	0.022
NO <sub>3</sub>	2.01	±	0.09
Ortho-phosphate	1.00	±	0.05
SO <sub>4</sub>	1.46	±	0.04
<b>Electrochemical property:</b>			
Conductivity (20 °C)	18.7	±	1.8 µS/cm
pH (20 °C)	6.3	±	0.6

Availability: ERM-CA408 is provided in units of about 95 mL in flame-sealed ampoules.

Substance	BCR-479 Freshwater (low contents)			BCR-480 Freshwater (high contents)		
Nitrate						
As amount of substance content	214	±	4	µmol/kg	885	±
As mass fraction	13.3	±	0.3	mg/kg	54.9	±
					13	µmol/kg
					0.8	mg/kg

Availability: Units of about 100 mL in white glass ampoules.

Substance	BCR-611 Bromide in ground water based on IC-measurements (low contents) (µg/kg)			BCR-612 Bromide in ground water based on IC-measurements (high contents) (µg/kg)			
Br	93	±	4		252	±	10

Availability: Set of four brown glass ampoules of 25 mL.

Substance	BCR-609 Ground water (low contents) (µg/kg)	BCR-610 Ground water (high contents) (µg/kg)	ERM-CA616 Ground water (high carbonate content) (mg/L)	BCR-617 Ground water (low carbonate content) (mg/kg)
Al	47.7	± 1.6	159	± 4
As	1.20	± 0.12	10.8	± 0.4
Cd	0.164	± 0.012	2.94	± 0.08
Cu	2.48	± 0.09	45.7	± 1.5
Pb	1.63	± 0.04	7.78	± 0.13
Ca			42.6	± 1.4
Cl			44.6	± 10.9
K			5.79	± 0.15
Mg			10.1	± 0.3
Mn			27.9	± 0.8
Na			2.24	± 0.10
NO <sub>3</sub>			426	± 5 µS/cm
Ortho-phosphate			7.12	± 0.18
PO <sub>4</sub>				25.8 ± 0.5
SO <sub>4</sub>				26.3 ± 0.5
Conductivity (20 °C)				
pH (20 °C)				

Availability: BCR-609 and BCR-610 are provided in 500 mL PE bottles; ERM-CA616 consists of about 95 mL natural groundwater in a flame-sealed ampoule; BCR-617 is provided in 75 mL glass ampoules.

Substance	ERM-CA615 Groundwater			
As	9.9	±	0.7	µg/L
Cd	0.106	±	0.011	µg/L
Fe	5.11	±	0.26	mg/L
Hg	0.037	±	0.004	µg/L
Mn	107	±	5	µg/L
Ni	25.3	±	1.1	µg/L
Pb	7.1	±	0.6	µg/L

Availability: One unit consists of about 95 mL natural groundwater in a flame-sealed ampoule.

Substance	ERM-CA713 Wastewater µg/L			
As	10.8	±	0.3	
Cd	5.09	±	0.20	
Cr	20.9	±	1.3	
Cu	101	±	7	
Fe	445	±	27	
Hg	1.84	±	0.11	
Mn	95	±	4	
Ni	50.3	±	1.4	
Pb	49.7	±	1.7	
Se	4.9	±	1.1	

Availability: ampoule containing approximately 100 mL of wastewater effluent acidified with HNO<sub>3</sub> to about pH 2.

## 1.2.2 CERTIFIED FOR THE EXTRACTABLE ELEMENT CONTENT AND SPECIES

Substance		BCR-483 Sewage sludge amended soil (mg/kg)			BCR-484 Sewage sludge amended (terra rossa) soil (mg/kg)			BCR-700 Organic-rich soil (mg/kg)		
<u>EDTA:</u>	Cd	24.3	±	1.3	0.509	±	0.030	65.2	±	3.5
	Cr	28.6	±	2.6				10.1	±	0.9
	Cu	215	±	11	88	±	4	89.4	±	2.8
	Ni	28.7	±	1.7	1.39	±	0.11	53.2	±	2.8
	Pb	229	±	8	47.9	±	2.6	103	±	5
	Zn	612	±	20	152	±	7	510	±	17
<u>Acetic acid:</u>	Cd	18.3	±	0.6	0.48	±	0.04	67.5	±	2.8
	Cr	18.7	±	1.0				19.0	±	1.1
	Cu	33.5	±	1.6	33.9	±	1.4	36.3	±	1.6
	Ni	25.8	±	1.0	1.69	±	0.16	99.0	±	5.1
	Pb	2.10	±	0.25	1.17	±	0.16	4.85	±	0.38
	Zn	620	±	24	193	±	7	719	±	24
<u>Calcium chloride extractable content</u>	Cd	(0.45	±	0.05)			(< 0.08)			
	Cr	(0.35	±	0.09)			(< 0.09)			
	Cu	(1.2	±	0.4)			(0.67	±	0.29)	
	Ni	(1.4	±	0.2)			(< 0.05)			
	Pb	(< 0.06)					(< 0.06)			
	Zn	(8.3	±	0.7)			(0.31	±	0.17)	
<u>Sodium nitrate extractable content</u>	Cd	(0.08	±	0.03)			(< 0.05)			
	Cr	(0.30	±	0.07)			(< 0.03)			
	Cu	(0.89	±	0.22)			(0.48	±	0.15)	
	Ni	(0.65	±	0.07)			(0.023	±	0.005)	
	Pb	(< 0.03)					(< 0.06)			
	Zn	(2.7	±	0.8)			(0.09	±	0.04)	
<u>Ammonium nitrate extractable content</u>	Cd	(0.26	±	0.05)			(0.003	±	0.002)	
	Cr	(0.27	±	0.10)			(< 0.06)			
	Cu	(1.2	±	0.3)			(1.1	±	0.4)	
	Ni	(1.1	±	0.3)			(0.033	±	0.017)	
	Pb	(0.020	±	0.013)			(< 0.06)			
	Zn	(6.5	±	0.9)			(0.17	±	0.05)	

Values in brackets are not certified.

Availability: BCR-483 and -484 are provided in glass bottles containing about 70 g of powder. BCR-700 is provided in glass bottles containing about 40 g of powder.

Substance		BCR-684 River sediment (mg/kg)		
NaOH-extractable P		550	±	21
HCl-extractable P		536	±	28
Inorganic P		1113	±	24
Organic P		209	±	9
Conc. HCl-extract. P		1373	±	35

Availability: Glass bottles containing about 35 g of powdered material.

Substance	BCR-701 Lake sediment (mg/kg)		
<u>Extractable mass fraction based on dry mass</u>			
<u>Step 1:</u> Cd	7.3	±	0.4
Cr	2.26	±	0.16
Cu	49.3	±	1.7
Ni	15.4	±	0.9
Pb	3.18	±	0.21
Zn	205	±	6
<u>Step 2:</u> Cd	3.77	±	0.28
Cr	45.7	±	2.0
Cu	124	±	3
Ni	26.6	±	1.3
Pb	126	±	3
Zn	114	±	5
<u>Step 3:</u> Cd	0.27	±	0.06
Cr	143	±	7
Cu	55	±	4
Ni	15.3	±	0.9
Pb	9.3	±	2.0
Zn	46	±	4
<u>Mass fraction based on dry mass</u>			
Cd	(0.13	±	0.08)
Cr	(62.5	±	7.4)
Cu	(38.5	±	11.2)
Ni	(41.4	±	4.0)
Pb	(11.0	±	5.2)
Zn	(95	±	13)

Values in brackets are not certified.

Availability: Glass bottles containing about 20 g of powdered material.

Substance	BCR-462 Coastal sediment (µg/kg)			BCR-646 Freshwater sediment (µg/kg)		
	54	±	15	480	±	80
Dibutyltin (DBT)	68	±	12	770	±	90
Monobutyltin (MBT)				610	±	120
Triphenyltin (TPhT)				29	±	11
Diphenyltin (DPhT)				36	±	8
Monophenyltin (MPhT)				69	±	18

Availability: Glass bottle containing about 25 g of powder for BCR-462 and 40 g of powder for BCR-646.

Substance	ERM-CC580 Estuarine sediment (mg/kg)		
Total Hg	132	±	3
CH <sub>3</sub> Hg <sup>+</sup>	0.075	±	0.004

Availability: Glass bottles containing about 40 g powder.

Substance	BCR-605 Urban dust (µg/kg)		
Trimethyllead (TriML)	7.9	±	1.2

Availability: Glass bottles containing about 15 g of powder.

	<b>BCR-545</b> Welding dust loaded on a filter (g/kg)		
Cr (VI) total leachable Cr	40.2	±	0.6
	39.5	±	1.3

Availability: Glass fibre filter loaded with welding dust containing about 100 µg Cr (VI).

Substance	<b>ERM-CE477</b> Mussel tissue (mg/kg)		
Tributyltin (TBT)	2.20	±	0.19
Dibutyltin (DBT)	1.54	±	0.12
Monobutyltin (MBT)	1.50	±	0.28

Availability: Glass bottle containing about 14 g of powder.

Substance	<b>BCR-463</b> Tuna fish (mg/kg)	<b>ERM-CE464</b> Tuna fish (mg/kg)				
Total Hg	2.85	±	0.16	5.24	±	0.10
CH <sub>3</sub> Hg <sup>+</sup>	3.04	±	0.16	5.50	±	0.17

Availability: Glass bottles containing about 15 g powder.

Substance	<b>BCR-627</b> Tuna fish tissue	<b>ERM-BC211</b> Rice				
Total As	4.8	±	0.3 mg/kg	260	±	13 µg/kg
Dimethylarsinic acid	2.0	±	0.3 µmol/kg	119	±	13 µg/kg
Sum of arsenite and arsenate				124	±	11 µg/kg
Arsenobetaine	52	±	3 µmol/kg			

Availability: BCR-627: Glass bottles containing about 10 g powder

ERM-BC211: Vial containing about 10 g of powder

### 1.2.3 CERTIFIED FOR ORGANIC POLLUTANTS

Substance	<b>ERM-CZ100</b> Fine dust (PM <sub>10</sub> -like) (mg/kg)		
Benz[a]anthracene	0.91	±	0.07
Benzo[a]pyrene	0.72	±	0.05
Benzo[b]fluoranthene	1.42	±	0.14
Benzo[j]fluoranthene	0.75	±	0.14
Benzo[k]fluoranthene	0.67	±	0.06
Dibenzo[a,h]anthracene	0.18	±	0.04
Indeno[1,2,3-c,d]pyrene	1.07	±	0.10
Sum of Benzo[b]fluoranthene, benzo[k]fluoranthene and benzo[j]fluoranthene	2.84	±	0.21

Availability: Vial containing about 0.5 g of fine dust.

Substance	BCR-524 Contaminated industrial soil (mg/kg)		
Pyrene	173	±	11
Benz[a]anthracene	22.5	±	1.8
Benzo[a]pyrene	8.6	±	0.5
Benzo[e]pyrene	10.6	±	1.4
Benzo[b]fluoranthene	13.5	±	1.6
Benzo[k]fluoranthene	6.2	±	0.6
Benzo[b]naphtho[2,1-d]thiophene	3.8	±	0.6
Indeno[1,2,3-cd]pyrene	5.1	±	0.4
Pentachlorophenol	0.034	±	0.005

Availability: Glass bottle containing about 40 g of powder.

Substance	BCR-535 Freshwater harbour sediment (mg/kg)		
Pyrene	2.52	±	0.18
Benz[a]anthracene	1.54	±	0.10
Benzo[a]pyrene	1.16	±	0.10
Benzo[e]pyrene	1.86	±	0.13
Benzo[b]fluoranthene	2.29	±	0.15
Benzo[k]fluoranthene	1.09	±	0.15
Indeno[1,2,3-cd]pyrene	1.56	±	0.14

Availability: Glass bottle containing about 40 g of powder.

Polychlorinated biphenyls (IUPAC No.)	BCR-481 Industrial soil (mg/kg)			BCR-536 Freshwater harbour sediment (µg/kg)		
28				44	±	5
52				38	±	4
101	37	±	3	44	±	4
105				3.5	±	0.6
118	9.4	±	0.7	28	±	3
128	9.1	±	0.8	5.4	±	1.2
138				27	±	4
149	97	±	7	49	±	4
153	137	±	7	50	±	4
156	7.0	±	0.5	3.0	±	0.4
163				17	±	3
170	52	±	4	13.4	±	1.4
180	124	±	6	22	±	2

Availability: BCR-481 is provided in brown glass bottles with a polyethylene insert containing approximately 25 g of soil. BCR-536 is provided in a glass bottle containing about 40 g of powder.

Substance	BCR-529 Industrial (sandy) soil			BCR-530 Industrial (clay) soil		
3,4-dichlorophenol	0.23	±	0.04 mg/kg	6.0	±	0.5 mg/kg
2,4,5-trichlorophenol	1.51	±	0.10 mg/kg	40	±	7 mg/kg
Pentachlorophenol	0.23	±	0.04 mg/kg	0.47	±	0.09 mg/kg
2,3,7,8 - TCDD (D48)	4.5	±	0.6 µg/kg			
1,2,3,7,8 - PeCDD (D54)	0.44	±	0.05 µg/kg			
1,2,3,4,7,8 - HxCDD (D66)	1.22	±	0.21 µg/kg			
1,2,3,6,7,8 - HxCDD (D67)	5.4	±	0.9 µg/kg	0.061	±	0.011 µg/kg
1,2,3,7,8,9 - HxCDD (D70)	3.0	±	0.4 µg/kg	0.0218	±	0.0029 µg/kg
2,3,7,8 - TCDF (F83)	0.078	±	0.013 µg/kg			
1,2,3,7,8 - PeCDF (F94)	0.145	±	0.028 µg/kg	0.24	±	0.04 µg/kg
2,3,4,7,8 - PeCDF (F114)	0.36	±	0.07 µg/kg	0.62	±	0.07 µg/kg
1,2,3,4,7,8 - HxCDF (F118)	3.4	±	0.5 µg/kg	0.321	±	0.016 µg/kg
1,2,3,6,7,8 - HxCDF (F121)	1.09	±	0.15 µg/kg	0.186	±	0.023 µg/kg
1,2,3,7,8,9 - HxCDF (F124)	0.022	±	0.010 µg/kg			
2,3,4,6,7,8 - HxCDF (F130)	0.37	±	0.05 µg/kg	0.126	±	0.012 µg/kg

Availability: Amber glass bottles containing about 50 g of dried soil.

Substance	BCR-677 Sewage sludge (ng/kg)		
2,3,7,8 - T <sub>4</sub> CDD (D48)	1.51	±	0.16
1,2,3,7,8 - P <sub>5</sub> CDD (D54)	4.1	±	0.9
1,2,3,6,7,8 - H <sub>6</sub> CDD (D67)	235	±	16
1,2,3,7,8,9 - H <sub>6</sub> CDD (D70)	79	±	7
1,2,3,4,6,7,8 - H <sub>7</sub> CDD (D73) O <sub>8</sub> CDD (D75)	3.5 x 10 <sup>3</sup>	±	0.4 x 10 <sup>3</sup>
2,3,7,8 - T <sub>4</sub> CDF (F83)	12.7 x 10 <sup>3</sup>	±	0.8 x 10 <sup>3</sup>
1,2,3,7,8 - P <sub>5</sub> CDF (F94)	45	±	4
2,3,4,7,8 - P <sub>5</sub> CDF (F114)	24.8	±	1.6
1,2,3,4,7,8 - H <sub>6</sub> CDF (F118)	16.9	±	1.5
1,2,3,6,7,8 - H <sub>6</sub> CDF (F121)	14.5	±	1.6
1,2,3,7,8,9 - H <sub>6</sub> CDF (F124)	6.1	±	0.8
2,3,4,6,7,8 - H <sub>6</sub> CDF (F130)	0.84	±	0.29
1,2,3,4,6,7,8 - H <sub>7</sub> CDF (F131)	5.6	±	0.6
1,2,3,4,7,8,9 - H <sub>7</sub> CDF (F134) O <sub>8</sub> CDF (F135)	62	±	3
	6.3	±	0.8
	177	±	7

Availability: BCR-677 consists of approximately 40 g of dried sewage sludge in amber glass bottles.

Substance	BCR-490 Fly ash (μg/kg)		
2,3,7,8 - T <sub>4</sub> CDD (D48)	0.169	±	0.012
1,2,3,7,8 - P <sub>5</sub> CDD (D54)	0.67	±	0.04
1,2,3,4,7,8 - H <sub>6</sub> CDD (D66)	0.95	±	0.11
1,2,3,6,7,8 - H <sub>6</sub> CDD (D67)	4.8	±	0.4
1,2,3,7,8,9 - H <sub>6</sub> CDD (D70)	2.84	±	0.17
2,3,7,8 - T <sub>4</sub> CDF (F83)	0.90	±	0.05
1,2,3,7,8 - P <sub>5</sub> CDF (F94)	1.71	±	0.12
2,3,4,7,8 - P <sub>5</sub> CDF (F114)	1.85	±	0.11
1,2,3,4,7,8 - H <sub>6</sub> CDF (F118)	2.37	±	0.12
1,2,3,6,7,8 - H <sub>6</sub> CDF (F121)	2.64	±	0.14
1,2,3,7,8,9 - H <sub>6</sub> CDF (F124)	0.34	±	0.05
2,3,4,6,7,8 - H <sub>6</sub> CDF (F130)	2.47	±	0.17

Availability: BCR-490 consists of approximately 30 g of fly ash in amber glass bottles.

Substance	BCR-615 Fly ash (low level) (ng/kg)		
2,3,7,8 - T <sub>4</sub> CDD (D48)	27	±	5
1,2,3,7,8 - P <sub>5</sub> CDD (D54)	92	±	12
1,2,3,4,7,8 - H <sub>6</sub> CDD (D66)	74	±	12
1,2,3,6,7,8 - H <sub>6</sub> CDD (D67)	103	±	13
1,2,3,7,8,9 - H <sub>6</sub> CDD (D70)	108	±	16
1,2,3,4,6,7,8 - H <sub>7</sub> CDD (D73) O <sub>8</sub> CDD (D75)	0.87 x 10 <sup>3</sup>	±	0.13 x 10 <sup>3</sup>
2,3,7,8 - T <sub>4</sub> CDF (F83)	1.75 x 10 <sup>3</sup>	±	0.20 x 10 <sup>3</sup>
1,2,3,7,8 - P <sub>5</sub> CDF (F94)	86	±	28
2,3,4,7,8 - P <sub>5</sub> CDF (F114)	176	±	26
1,2,3,4,7,8 - H <sub>6</sub> CDF (F118)	125	±	20
1,2,3,6,7,8 - H <sub>6</sub> CDF (F121)	203	±	21
1,2,3,7,8,9 - H <sub>6</sub> CDF (F124)	204	±	23
2,3,4,6,7,8 - H <sub>6</sub> CDF (F130)	13.3	±	2.0
1,2,3,4,6,7,8 - H <sub>7</sub> CDF (F131)	130	±	15
1,2,3,4,7,8,9 - H <sub>7</sub> CDF (F134) O <sub>8</sub> CDF (F135)	0.75 x 10 <sup>3</sup>	±	0.09 x 10 <sup>3</sup>
	61	±	6
	0.29 x 10 <sup>3</sup>	±	0.04 x 10 <sup>3</sup>

Availability: BCR-615 consists of approximately 50 g of dried fly ash in amber glass bottles.

Substance	BCR-683 Beech wood (mg/kg)		
Benz[a]anthracene	6.5	±	0.7
Benzo[a]pyrene	3.4	±	0.4
Benzo[e]pyrene	9.3	±	1.0
Benzo[b]fluoranthene	5.8	±	0.6
Benzo[k]fluoranthene	2.58	±	0.29
Pentachlorophenol	3.6	±	0.5

Availability: Glass bottle containing about 60 g of powder.

Polychlorinated biphenyls (IUPAC No.)	BCR-682 Mussel tissue (µg/kg)			BCR-718 Canned fresh herring (µg/kg)			BCR-719 Canned fresh chub (ng/kg)			
	28	0.30	±	0.07	0.41	±	0.04	196	±	6
52	0.78	±		1.00	±		13.6	±	0.4	
77										
81					2.12	±	0.06			
101					0.63	±	0.06			
105					1.78	±	0.07			
118	2.6	±	0.3		0.62	±	0.101	20.0	±	0.8
126					2.97	±	0.11			
128										
138	4.6	±	0.8							
138 + 163					2.58	±	0.11			
149	5.7	±	0.9		4.62	±	0.10			
153	9.2	±	0.8		0.19	±	0.09			
156										
169					0.350	±	0.026	1.80	±	0.15
170	0.17	±	0.05		0.795	±	0.027			
180	0.77	±	0.07							

Availability: BCR-682, BCR-718 and BCR-719 are provided in sealed tin cans containing approximately 70 g fresh mussel tissue.

Polychlorinated biphenyls (IUPAC No.)	BCR-420 Waste mineral oil (low level) (mg/kg)			BCR-449 Waste mineral oil (high level) (mg/kg)			
	28	0.61	±	0.06	0.80	±	0.07
52					31.4	±	1.8
101	1.45	±	0.18		57.2	±	1.9
105					17.4	±	1.0
118	1.69	±	0.14		46.6	±	2.4
128					12.5	±	0.7
153	0.92	±	0.06		39.0	±	1.7
156					6.9	±	0.5
170					6.6	±	0.6
180	0.195	±	0.017		10.4	±	0.4

Availability: BCR-420 is provided in units of about 7.5 g in glass ampoules and BCR-449 in units of about 50 g in glass ampoules.

## 1.2.4 OTHERS

Parameter	IRMM443-1 (EUROSOIL 1)	IRMM443-2 (EUROSOIL 2)	IRMM443-3 (EUROSOIL 3)
K <sub>f</sub> of atrazine <sup>(1)</sup>	7.0 ± 1.5	2.7 ± 0.7	2.4 ± 0.7
1/n of atrazine <sup>(1)</sup>	0.91 ± 0.11	0.93 ± 0.12	0.91 ± 0.13
K <sub>f</sub> of 2,4-D <sup>(1)</sup>	2.5 ± 1.0	0.99 ± 0.30	1.31 ± 0.28
1/n of 2,4-D <sup>(1)</sup>	0.9 ± 0.4	0.96 ± 0.15	0.93 ± 0.15
K <sub>f</sub> of lindane <sup>(1)</sup>		48 ± 11	
1/n of lindane <sup>(1)</sup>		0.98 ± 0.15	
pH in water <sup>(2)</sup>	6.21 ± 0.30	8.1 ± 0.9	6.2 ± 0.4
pH in 0.01M CaCl <sub>2</sub> <sup>(2)</sup>	5.65 ± 0.24	7.5 ± 0.8	5.5 ± 0.4

Parameter	IRMM443-4 (EUROSOIL 4)	IRMM443-5 (EUROSOIL 5)	IRMM443-7 (EUROSOIL 7)
K <sub>f</sub> of atrazine <sup>(1)</sup>	0.7 ± 0.4	13 ± 6	4.8 ± 1.1
1/n of atrazine <sup>(1)</sup>	0.87 ± 0.22	0.9 ± 0.4	0.92 ± 0.15
K <sub>f</sub> of 2,4-D <sup>(1)</sup>	0.39 ± 0.21	18 ± 7	8.2 ± 1.8
1/n of 2,4-D <sup>(1)</sup>	0.9 ± 0.4	0.9 ± 0.4	0.88 ± 0.15
K <sub>f</sub> of lindane <sup>(1)</sup>	8.3 ± 2.2		
1/n of lindane <sup>(1)</sup>	0.96 ± 0.12		
pH in water <sup>(2)</sup>	7.5 ± 0.7	4.1 ± 1.5	5.1 ± 0.8
pH in 0.01M CaCl <sub>2</sub> <sup>(2)</sup>	6.8 ± 0.6	3.1 ± 1.1	4.3 ± 0.7

(1) Determination according OECD Testguideline 106.

(2) Measurement based on ISO Standard 10390.

Uncertainty express as estimated expanded uncertainty as defined in the Guide to the Expression of Uncertainty in Measurement (GUM).

Availability: Brown glass bottles with 200 g of air-dried fine soil (< 2 mm).

## 2 MATERIALS RELATED TO THE ANALYSIS OF FOOD AND FEEDING STUFF

### 2.1 PURE MATERIALS AND SYNTHETIC MIXTURES

BCR-123 Ethanol			
Parameter	Ethanol H	Ethanol M	Ethanol L
(D/H) <sub>I</sub>	$109.65 \times 10^{-6}$ ± $0.20 \times 10^{-6}$	$101.69 \times 10^{-6}$ ± $0.17 \times 10^{-6}$	$90.30 \times 10^{-6}$ ± $0.18 \times 10^{-6}$
(D/H) <sub>II</sub>	$119.76 \times 10^{-6}$ ± $0.25 \times 10^{-6}$	$130.94 \times 10^{-6}$ ± $0.21 \times 10^{-6}$	$122.20 \times 10^{-6}$ ± $0.4 \times 10^{-6}$
R	2.184 ± 0.005	2.575 ± 0.006	2.708 ± 0.009

Availability: Units of 3 sealed NMR tubes containing respectively H-, M-, and L-ethanols, to which the tetramethylurea internal standard and the C<sub>6</sub>F<sub>6</sub> lock substance are added. 10 mm (BCR-123A) or 15 mm (BCR-123B) O.D. NRM tubes can be supplied.

Parameter	Unit	BCR-656 (96% ethanol)	
(D/H) <sub>I</sub> by <sup>2</sup> H-NMR	ppm	102.84	± 0.20
(D/H) <sub>II</sub> by <sup>2</sup> H-NMR	ppm	132.07	± 0.30
R by <sup>2</sup> H-NMR		2.570	± 0.005
δ <sup>13</sup> C <sub>VPDB</sub> by IRMS	‰	-26.91	± 0.07
Alcoholic grade t <sub>D</sub>	w/w %	94.61	± 0.05

Availability: BCR-656: Units of 25 mL of 96 % vol. neutral ethanol from wine in glass bottle.

Parameter	Unit	BCR-657 (Sugar)	BCR-658 (Synthetic wine)	BCR-659 (Synthetic wine)	BCR-660 (Ethanol in water)
(D/H) <sub>I</sub> by <sup>2</sup> H-NMR	ppm				102.90 ± 0.16
(D/H) <sub>II</sub> by <sup>2</sup> H-NMR	ppm				131.95 ± 0.23
R by <sup>2</sup> H-NMR					2.567 ± 0.005
$\delta^{13}\text{C}_{\text{VPDB}}$ by IRMS	‰	-10.76 ± 0.04			-26.72 ± 0.09
$\delta^{18}\text{O}_{\text{VSMOW}}$ of water from wine by IRMS	‰		-7.19 ± 0.04	-7.18 ± 0.02	
(D/H) <sub>w</sub> of water (IRMS)	ppm				148.68 ± 0.14
Alcoholic grade t <sub>D</sub>	w/w %				11.96 ± 0.06 <sup>1)</sup>

1) in v/v %

Availability: BCR-656: Units of 25 mL of 96 % vol. neutral ethanol from wine in glass bottle;

BCR-657: Units of approx. 1 g of dry glucose in a sealed amber vial;

BCR-658: Units of 25 mL of synthetic wine solution in glass bottle;

BCR-659: Units of 25 mL of synthetic wine solution in glass bottle;

BCR-660: Units of 450 mL of aqueous ethanol solution in glass bottle.

### Tetramethylurea (STA-003m)

Tetramethylurea (TMU) which is used as Internal Standard in routine SNIF-NMR analysis is available in 500 mL quantities. The D/H nominal value of tetramethylurea batches is determined by multiple calibration at 61.45/400 MHz, 61.4/400 MHz and 45.05/400 MHz for deuterium and given in an accompanying analytical report.

Substance	BCR-423 (RM) Aflatoxin M <sub>1</sub> in chloroform (µg/mL)
Aflatoxin M <sub>1</sub>	(9.93)

Value in brackets is not certified.

Availability: Sealed ampoules containing about 2.5 mL.

Compound	BCR-663 Saxitoxin in acetic acid
Saxitoxin·2HCl	<u>Mass fraction</u> (mg/kg)

Availability: BCR-663 is available in ampoules containing 1 mL.

Substance	ERM-AC699 Zearalenone in acetonitrile
ZON	<u>Mass concentration</u> (µg/mL)

Availability: ERM-AC699 is supplied in ampoules filled and sealed under nitrogen in amounts of 4 mL.

Substance	<b>ERM-AC057</b> Aflatoxin B1 in acetonitrile		
Aflatoxin B1	<u>Mass fraction</u> ( $\mu\text{g/g}$ ) 3.79 $\pm$ 0.11	<u>Mass concentration at 20 °C</u> ( $\mu\text{g/mL}$ ) (2.97 $\pm$ 0.09)	

Values in brackets are not certified.

Availability: ERM-AC057 is supplied in amber glass ampoules filled with 4 mL.

Substance	<b>ERM-AC058</b> Aflatoxin B2 in acetonitrile		
Aflatoxin B2	<u>Mass fraction</u> ( $\mu\text{g/g}$ ) 3.80 $\pm$ 0.08	<u>Mass concentration at 20 °C</u> ( $\mu\text{g/mL}$ ) (2.98 $\pm$ 0.06)	

Values in brackets are not certified.

Availability: ERM-AC058 is supplied in amber glass ampoules filled with 4 mL.

Substance	<b>ERM-AC059</b> Aflatoxin G1 in acetonitrile		
Aflatoxin G1	<u>Mass fraction</u> ( $\mu\text{g/g}$ ) 3.78 $\pm$ 0.13	<u>Mass concentration at 20 °C</u> ( $\mu\text{g/mL}$ ) (2.96 $\pm$ 0.10)	

Values in brackets are not certified.

Availability: ERM-AC059 is supplied in amber glass ampoules filled with 4 mL.

Substance	<b>ERM-AC060</b> Aflatoxin G2 in acetonitrile		
Aflatoxin G2	<u>Mass fraction</u> ( $\mu\text{g/g}$ ) 3.80 $\pm$ 0.07	<u>Mass concentration at 20 °C</u> ( $\mu\text{g/mL}$ ) (2.98 $\pm$ 0.06)	

Values in brackets are not certified.

Availability: ERM-AC060 is supplied in amber glass ampoules filled with 4 mL.

Substance	<b>IRMM-315</b> 4-Deoxynivalenol in acetonitrile		
4-Deoxynivalenol	<u>Mass fraction</u> ( $\mu\text{g/g}$ ) 25.1 $\pm$ 1.2	<u>Mass concentration</u> ( $\mu\text{g/mL}$ ) (19.7 $\pm$ 0.9)	

Values in brackets are not certified.

Availability: IRMM-315 is supplied in amber glass ampoules filled with 4 mL.

Substance	<b>IRMM-316</b> Nivalenol in acetonitrile		
Nivalenol	<u>Mass fraction</u> ( $\mu\text{g/g}$ ) 24.0 $\pm$ 1.1	<u>Mass concentration</u> ( $\mu\text{g/mL}$ ) (18.8 $\pm$ 0.9)	

Values in brackets are not certified.

Availability: IRMM-316 is supplied in amber glass ampoules filled with 4 mL.

## 2.2 MATRIX MATERIALS

### 2.2.1 CERTIFIED FOR GMO CONTENT

The materials were prepared by quantitative mixing of non-genetically modified powder and genetically modified powder, produced from ground seed with the help of a dry-mixing technique, and are intended for the calibration of methods for the detection of genetically modified food.

#### CRMs for genetically modified Roundup Ready™ soya beans (ERM-BF410k)

Six CRMs of dried soya bean powder with different mass fractions of genetically modified (Roundup Ready™) soya beans were produced by IRMM.

	Certified value Roundup Ready mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF410ak	< 0.7	-
ERM-BF410bk	1.0	0.5
ERM-BF410dk	10.0	1.0
ERM-BF410gk	100	7

Availability: Vials containing about 1 g of soya bean powder.

#### CRMs for genetically modified Bt-176 maize (ERM-BF411)

Six CRMs of dried maize powder with different mass fractions of genetically modified (Bt-176) maize were produced by IRMM.

	Certified value Bt-176 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF411a	< 0.14	-
ERM-BF411b	1.00	0.29
ERM-BF411c	5.0	0.6
ERM-BF411d	10.0	0.8
ERM-BF411e	20.0	1.1
ERM-BF411f	50.0	1.8

Availability: Vials containing about 1 g of maize powder.

#### CRMs for genetically modified Bt-11 maize (ERM-BF412)

Six CRMs of dried maize powder with different mass fractions of genetically modified (Bt-11) maize were produced by IRMM.

	Certified value Bt-11 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF412a	< 0.12	-
ERM-BF412b	0.98	0.29
ERM-BF412c	4.9	0.6
ERM-BF412d	9.8	0.9
ERM-BF412e	19.6	1.3
ERM-BF412f	48.9	2.1

Availability: Vials containing about 1 g of maize powder.

## CRMs for genetically modified MON 810 maize (ERM-BF413k)

Four CRMs of dried maize powder with different mass fractions of genetically modified (MON 810) maize were produced by IRMM.

	Certified value MON 810 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF413ak	< 0.9	-
ERM-BF413ck	4.9	1.0
ERM-BF413ek *	19.8	1.5
ERM-BF413gk	99	5

Availability: Vials containing about 1 g of maize powder.

\* ERM-BF413ek is also certified for the DNA copy number ratio.

	Certified value MON 810 DNA copy number ratio (%)	Uncertainty (%)
ERM-BF413ek	0.77	0.08

	ERM-AD413 DNA fragments per plasmid	
Fragment of 5' plant-P35S junction DNA/plasmid	1	(negligible uncertainty)
Fragment of hmg DNA/plasmid	1	(negligible uncertainty)
Ratio between the number of 5' plant-P35S junction and hmg fragments in the plasmid by duplex rt-PCR <sup>(1)</sup> and simplex rt-PCR <sup>(2)</sup>	(1.00 <sup>1</sup> (1.04 <sup>2</sup>	± 0.06) ± 0.06)

Values in brackets are not certified.

Availability: ERM-AD413 is available in vials.

## CRMs for genetically modified GA21 maize (ERM-BF414)

Six CRMs of dried maize powder with different mass fractions of genetically modified (GA21) maize were produced by IRMM.

	Certified value GA21 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF414a	< 0.8	-
ERM-BF414b	1.0	0.8
ERM-BF414c	4.9	1.0
ERM-BF414d	9.9	1.1
ERM-BF414e	17.2	1.2
ERM-BF414f	42.9	1.7

Availability: Vials containing about 1 g of maize powder.

## CRMs for genetically modified NK603 maize (ERM-BF415)

Six CRMs of dried maize powder with different mass fractions of genetically modified (NK603) maize were produced by IRMM.

	Certified value NK603 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF415a	< 0.4	-
ERM-BF415b	1.0	0.4
ERM-BF415c	4.9	0.5
ERM-BF415d	9.8	0.7
ERM-BF415e *	19.6	0.9
ERM-BF415f	49.1	1.3

Availability: Vials containing about 1 g of maize powder.

\* ERM-BF415e is also certified for the DNA copy number ratio.

	Certified value NK603 maize DNA copy number ratio (%)	Uncertainty (%)
ERM-BF415e	0.95	0.11

	ERM-AD415 DNA fragments per plasmid	
	Number	
Fragment of 3' insertion-specific DNA / pIRMM-0086	1	(negligible uncertainty)
Fragment of <i>hmg</i> DNA / pIRMM-0086	1	(negligible uncertainty)

Availability: ERM-AD415 is available in vials.

## CRMs for genetically modified MON 863 maize (ERM-BF416)

Four CRMs of dried maize powder with different mass fractions of genetically modified (MON 863) maize were produced by IRMM.

	Certified value MON 863 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF416a	< 1.0	-
ERM-BF416b	1.0	- 0.3 ; + 1.0
ERM-BF416c	9.8	- 0.7 ; + 1.2
ERM-BF416d	98.5	- 2.2 ; + 2.5

Availability: Vials containing about 1 g of maize powder.

## CRMs for genetically modified MON 863 x MON 810 maize (ERM-BF417)

Four CRMs of dried maize powder with different mass fractions of genetically modified (MON 863 x MON 810) maize were produced by IRMM.

	Certified value MON 863 x MON 810 mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF417a	< 1.0	-
ERM-BF417b	1.0	- 0.2 ; + 1.0
ERM-BF417c	9.8	- 0.7 ; + 1.2
ERM-BF417d	98.5	- 2.0 ; + 2.4

Availability: Vials containing about 1 g of maize powder.

### CRMs for genetically modified 1507 maize (ERM-BF418)

Four CRMs of dried maize powder with different mass fractions of genetically modified (1507) maize were produced by IRMM.

	Certified value 1507 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF418a	< 0.5	-
ERM-BF418b	1.0	- 0.2 ; + 0.6
ERM-BF418c	9.9	- 0.6 ; + 0.8
ERM-BF418d	98.6	- 1.7 ; + 2.0

Availability: Vials containing about 1 g of maize powder.

### CRMs for genetically modified H7-1 sugar beet (ERM-BF419)

Two CRMs of dried sugar beet powder with different mass fractions of genetically modified (H7-1) sugar beet were produced by IRMM.

	Certified value H7-1 sugar beet mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF419a	0	0
ERM-BF419b	1000	0

Availability: Vials containing about 1 g of sugar beet powder.

### CRMs for genetically modified 3272 maize (ERM-BF420)

Three CRMs of dried maize powder with different mass fractions of genetically modified (3272) maize were produced by IRMM.

	Certified value 3272 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF420a	< 1.3	-
ERM-BF420b	9.8	1.2
ERM-BF420c	98	8

Availability: Vials containing about 1 g of maize powder.

### CRMs for genetically modified EH92-527-1 potato (ERM-BF421)

Two CRMs of dried potato powder with different mass fractions of genetically modified (EH92-527-1) potato were produced by IRMM.

	Certified value Number fraction of EH92-527-1 potato / total number of potatoes [%]	Certified property Identity	Uncertainty
ERM-BF421a	0	potato without the EH92-527-1 event EH92-527-1 potato	Not applicable
ERM-BF421b	100		

Availability: ERM-BF421a: vials containing about 1 g of potato powder, ERM-BF421b: vials containing about 0.5 g of potato powder.

### **CRMs for genetically modified 281-24-236 x 3006-210-23 cotton seed (ERM-BF422)**

Four CRMs of dried cotton seed powder with different mass fractions of genetically modified (281-24-236 x 3006-210-23) cotton seed were produced by IRMM.

	Certified value 281-24-236 x 3006-210-23 cotton seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF422a	< 0.5	-
ERM-BF422b	> 979	-
ERM-BF422c	10.0	1.7
ERM-BF422d	100	16

Availability: Vials containing about 1 g of cotton seed powder.

### **CRMs for genetically modified MIR604 maize (ERM-BF423)**

Four CRMs of dried maize powder with different mass fractions of genetically modified (MIR604) maize were produced by IRMM.

	Certified value MIR604 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF423a	< 0.9	-
ERM-BF423b	1.0	- 0.3 ; + 1.0
ERM-BF423c	9.8	- 0.9 ; + 1.3
ERM-BF423d	98.5	- 2.6 ; + 2.9

Availability: Vials containing about 1 g of maize powder.

### **CRMs for genetically modified 59122 maize (ERM-BF424)**

Four CRMs of dried maize powder with different mass fractions of genetically modified (59122) maize were produced by IRMM.

	Certified value 59122 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF424a	< 1.2	-
ERM-BF424b	1.0	- 0.2 ; + 1.2
ERM-BF424c	9.9	- 0.8 ; + 1.4
ERM-BF424d	98.7	- 5.8 ; + 5.9

Availability: Vials containing about 1 g of maize powder.

### **CRMs for genetically modified 356043 soya seed (ERM-BF425)**

Four CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed were produced by IRMM.

	Certified value 356043 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF425a	< 0.5	-
ERM-BF425b	1.0	0.4
ERM-BF425c *	10.0	1.1
ERM-BF425d	100	9

Availability: Vials containing about 1 g of soya seed powder.

\* ERM-BF425c is also certified for the DNA copy number ratio.

	Certified value 356043 soya DNA copy number ratio (%)	Uncertainty (%)
ERM-BF425c	0.85	0.11

Substance	<b>ERM-AD425</b> DNA fragments per plasmid	
	<u>Number</u>	
Fragment of 5' insert-to-plant junction DNA / pIRMM-0073	1	(negligible uncertainty)
Fragment of <i>le1</i> DNA / pIRMM-0073	1	(negligible uncertainty)

Availability: ERM-AD425 is available in vials.

### CRMs for genetically modified 305423 soya seed (ERM-BF426)

Four CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed were produced by IRMM.

	Certified value 305423 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF426a	< 0.8	-
ERM-BF426b	5.0	0.8
ERM-BF426c	10.0	1.0
ERM-BF426d	100	7

Availability: Vials containing about 1 g of soya seed powder.

### CRMs for genetically modified 98140 maize (ERM-BF427)

Four CRMs of dried maize seed powder with different mass fractions of genetically modified (98140) maize were produced by IRMM.

	Certified value 98140 maize seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF427a	< 0.4	-
ERM-BF427b	5.0	0.6
ERM-BF427c *	20.0	0.8
ERM-BF427d	100	4

Availability: Vials containing about 1 g of maize seed powder.

\* ERM-BF427c is also certified for the DNA copy number ratio.

	Certified value 98140 maize DNA copy number ratio (%)	Uncertainty (%)
ERM-BF427c	1.75	0.13

Substance	<b>ERM-AD427</b> DNA fragments per plasmid	
	<u>Number</u>	
Fragment of 5' insert-to-plant junction DNA / pIRMM-0090	1	(negligible uncertainty)
Fragment of <i>hmg</i> DNA / pIRMM-0090	1	(negligible uncertainty)

Availability: ERM-AD427 is available in vials.

### **CRMs for genetically modified GHB119 cotton (ERM-BF428)**

Three CRMs of dried cotton seed powder with different mass fractions of genetically modified (GHB119) cotton were produced by IRMM.

	Certified value GHB119 cotton seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF428a	< 0.2	-
ERM-BF428b	10	4
ERM-BF428c	100	11

Availability: Vials containing about 1 g of cotton seed powder.

### **CRMs for genetically modified T304-40 cotton (ERM-BF429)**

Three CRMs of dried cotton seed powder with different mass fractions of genetically modified (T304-40) cotton were produced by IRMM.

	Certified value T304-40 cotton seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF429a	< 0.4	-
ERM-BF429b	10	1.3
ERM-BF429c	100	11

Availability: Vials containing about 1 g of cotton seed powder.

### **CRMs for genetically modified AM04-1020 potato (ERM-BF430)**

Five CRMs of dried potato powder with different mass fractions of genetically modified (AM04-1020) potato were produced by IRMM.

	Certified value AM04-1020 potato mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF430a	0	-
ERM-BF430b	1000	-
ERM-BF430c	10	1.4
ERM-BF430d	40	5
ERM-BF430e	100	12

Availability: Vials containing about 1 g of potato powder.

### **CRMs for genetically modified AV43-6-G7 potato (ERM-BF431)**

Five CRMs of dried potato powder with different mass fractions of genetically modified potato AV43-6-G7 were produced by IRMM.

	Certified value AV43-6-G7 potato mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF431a	0	-
ERM-BF431b	1000	-
ERM-BF431c	9.9	1.3
ERM-BF431d	40	5
ERM-BF431e	99	10

Availability: Vials containing about 1 g of potato powder

### **CRMs for genetically modified DAS-68416-4 soya seed (ERM-BF432)**

Four CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed DAS-68416-4 were produced by IRMM.

	Certified value DAS-68416-4 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF432a	< 0.3	-
ERM-BF432b	5.0	0.6
ERM-BF432c	10.0	1.7
ERM-BF432d	100	13

Availability: Vials containing about 1 g of soya seed powder.

### **CRMs for genetically modified DAS-40278-9 maize (ERM-BF433)**

Four CRMs of dried maize powder with different mass fractions of genetically modified DAS-40278-9 maize were produced by IRMM.

	Certified value DAS-40278-9 maize mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF433a	< 0.3	-
ERM-BF433b	5.0	0.6
ERM-BF433c	10.0	0.9
ERM-BF433d	100	8

Availability: Vials containing about 1 g of maize powder.

### **CRMs for genetically modified 73496 rapeseed (ERM-BF434)**

Five CRMs of dried rapeseed powder with different mass fractions of genetically modified rapeseed 73496 were produced by IRMM.

	Certified value 73496 rapeseed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF434a	< 0.04	-
ERM-BF434b	>988	-
ERM-BF434c	1.00	0.15
ERM-BF434d	10.0	1.4
ERM-BF434e	100	12

Availability: Vials containing about 1 g of rapeseed powder.

### **CRMs for genetically modified PH05-026-0048 potato (ERM-BF435)**

Two CRMs for the detection of genetically modified PH05-026-0048 potato were produced by IRMM.

	Certified value PH05-026-0048 potato mass fraction (g/kg)
ERM-BF435a	< 0.4

  

	Certified identity
ERM-BF435b	Positive for event PH05-026-0048

Availability: Vials containing about 1 g of potato powder

## CRMs for genetically modified DAS-44406-6 soya seed (ERM-BF436)

Five CRMs of dried soya seed powder with different mass fractions of genetically modified soya seed DAS-44406-6 were produced by IRMM.

	Certified value DAS-44406-6 soya seed mass fraction (g/kg)	Uncertainty (g/kg)
ERM-BF436a	< 0.06	-
ERM-BF436b	>986	-
ERM-BF436c	1.00	0.14
ERM-BF436d	10.0	1.0
ERM-BF436e	100	9

Availability: Vials containing about 1 g of soya seed powder.

## 2.2.2 CERTIFIED FOR NATURAL TOXINS AND XENOBIOTICS

Substance	BCR-458 Coconut oil (µg/kg)	BCR-459 Coconut oil (µg/kg)
Pyrene	9.4 ± 1.5	< 0.9
Chrysene	4.9 ± 0.4	< 0.6
Benzo[k]fluoranthene	1.87 ± 0.18	< 0.2
Benzo[a]pyrene	0.93 ± 0.09	< 0.3
Benzo[ghi]perylene	0.97 ± 0.07	< 0.2
Indeno[1,2,3-cd]pyrene	1.00 ± 0.07	< 0.2

Availability: BCR-458 and BCR-459 are provided in ampoules containing approximately 45 g.

Polychlorinated biphenyls (IUPAC No.)	BCR-450 Natural milk powder (µg/kg)	ERM-BB444 Natural pork fat (blank) (µg/kg)	ERM-BB445 Spiked pork fat (very low level) (µg/kg)	ERM-BB446 Spiked pork fat (low level) (µg/kg)
28		< 2	14.8 ± 1.3	29.6 ± 2.1
52	1.16 ± 0.17	< 2	12.9 ± 0.9	25.5 ± 1.8
101		< 2	12.5 ± 1.2	30 ± 4
118	3.3 ± 0.4	< 2	12.7 ± 1.3	30.2 ± 2.7
138		< 2	14.6 ± 1.6	32 ± 4
153	19.0 ± 0.7	< 2	13.1 ± 1.1	30.8 ± 2.4
156	1.62 ± 0.20			
170	4.8 ± 0.6			
180	11.0 ± 0.7	< 2	12.6 ± 0.9	29.8 ± 2.5
sum (28, 52, 101, 118, 138, 153, 180)		< 14	93 ± 7	207 ± 11
*PDBE 47 γ-HCH (lindane)		(3.7) (5.7)	(3.9) (5.6)	(6.1) (4.6)

\* 2,2'4,4'-Tetrabromo-diphenylether

Values in brackets are not certified.

Availability: BCR-450 in brown glass bottles of about 20 g. ERM-BB444 to 446 in glass ampoules of about 5 g.

Polychlorinated biphenyls (IUPAC No.)	BCR-349 Cod liver oil ( $\mu\text{g/kg}$ )			ERM-BB350 Fish oil (ng/g)		
28	68	±	8	21.3	±	1.1
52	149	±	21	37.4	±	2.2
74				23.0	±	1.9
95				(38	±	47)
99				62	±	6
101	372	±	18	111	±	5
105				25.8	±	2.1
110				54.1	±	2.8
118	460	±	40	84	±	4
138				137	±	10
149				88	±	9
153	940	±	40	220	±	11
156				20.1	±	1.3
163				(43	±	73)
167				(17	±	27)
177				25.8	±	2.0
180	282	±	23	67	±	4
183				22.5	±	1.8
187				67	±	5
194				23.4	±	1.5
196				41	±	7

Values in brackets are not certified.

Availability: BCR-349 and ERM-BB350 are provided in sealed glass ampoules containing approximately 2 g fish oil.

Substance	BCR-598 Cod liver oil ( $\mu\text{g/kg}$ )		
HCB	55.7	±	2.0
$\alpha$ -HCH	42	±	3
$\beta$ -HCH	16	±	3
$\gamma$ -HCH	23	±	4
$\gamma$ -Chlordane	6.9	±	1.6
$\alpha$ -Chlordane	24.4	±	1.8
Oxychlordane	11.0	±	1.8
Transnonachlor	39	±	4
Dieldrin	59	±	4
p,p'-DDE	$0.61 \times 10^3$	±	$0.04 \times 10^3$
o,p'-DDD	30	±	4
p,p'-DDD	$0.40 \times 10^3$	±	$0.03 \times 10^3$
p,p'-DDT	$0.179 \times 10^3$	±	$0.018 \times 10^3$

Availability: BCR-598 is provided in sealed glass ampoules containing approximately 5 g under dry argon.

Substance	ERM-BB430 Pork fat (mg/kg)		
HCB	0.193	±	0.017
$\alpha$ -HCH	0.25	±	0.04
$\beta$ -HCH	0.109	±	0.010
$\gamma$ -HCH	(1.87	±	0.31)
$\beta$ -HEPO	0.213	±	0.016
Dieldrin	(0.21	±	0.05)
Endrin	(0.055	±	0.016)
p,p'-DDT	0.48	±	0.07
p,p'-DDD	0.222	±	0.022
p,p'-DDE	0.38	±	0.09

Values in brackets are not certified.

Availability: ERM-BB430 is provided in sealed amber glass ampoules containing approximately 5 g of material bottled under argon.

Substance	BCR-115 Animal feed (mg/kg)		
HCB	0.019	4	0.001
$\beta$ -HCH	0.023	4	0.002
$\gamma$ -HCH	0.021	8	0.002
Heptachlor	0.019	0	0.001
$\gamma$ -Chlordane	0.048		0.006
$\alpha$ -Endosulfan	0.046		0.004
Dieldrin	0.018	1	0.002
Endrin	0.046		0.006
$\alpha,p'$ -DDT	0.046		0.005
$p,p'$ -DDE	0.047		0.004

Availability: BCR-115 is provided in sealed hard glass ampoules containing approximately 30 g under dry N<sub>2</sub>. The sample is a homogeneous animal feed obtained from commonly used ingredients and enriched with organochlorine pesticides.

Substance	BCR-187 Milk powder ( $\mu$ g/kg)	
HCB	1.45	37.4
$\alpha$ -HCH	$\pm$ 0.21	$\pm$ 2.7
$\beta$ -HCH	1.80	$\pm$ 0.14
$\gamma$ -HCH	5.7	$\pm$ 1.2
$\beta$ -HEPO	$\pm$ 0.8	45.4
$p,p'$ -DDE	6.6	$\pm$ 2.9
Dieldrin	$\pm$ 0.6	32.0
Endrin		$\pm$ 1.9
$p,p'$ -DDT		51
		$\pm$ 4
		36.1
		$\pm$ 2.5
		6.2
		$\pm$ 0.9
		69
		$\pm$ 5

Availability: The samples are provided in sealed hard glass ampoules containing about 20 g (under Argon).

Substance	BCR-607 Natural spray dried milk powder (ng/kg)
2,3,7,8 - T <sub>4</sub> CDD (D48)	0.25
1,2,3,7,8 - P <sub>5</sub> CDD (D54)	$\pm$ 0.03
0.79	0.04
1,2,3,4,7,8 - H <sub>6</sub> CDD (D66)	0.42
1,2,3,6,7,8 - H <sub>6</sub> CDD (D67)	$\pm$ 0.07
0.98	0.11
1,2,3,7,8,9 - H <sub>6</sub> CDD (D70)	0.34
2,3,7,8 - T <sub>4</sub> CDF (F83)	$\pm$ 0.05
0.05	0.03
1,2,3,7,8 - P <sub>5</sub> CDF (F94)	0.054
2,3,4,7,8 - P <sub>5</sub> CDF (F114)	$\pm$ 0.013
1.81	0.13
1,2,3,4,7,8 - H <sub>6</sub> CDF (F118)	0.94
1,2,3,6,7,8 - H <sub>6</sub> CDF (F121)	$\pm$ 0.04
1.01	0.09
2,3,4,6,7,8 - H <sub>6</sub> CDF (F130)	1.07
	$\pm$ 0.05

Availability: Amber glass bottles containing approximately 100 g.

Substance	ERM-BC190 Rapeseed (colza)	ERM-BC366 Rapeseed (colza)	ERM-BC367 Rapeseed (colza)
	<u>Certified values</u>	<u>Certified values</u>	<u>Certified values</u>
Total glucosinolate content	23 $\pm$ 4    mmol/kg	11.9 $\pm$ 1.3    mmol/kg	99 $\pm$ 9    mmol/kg
Sulphur content	4.72 $\pm$ 0.22    g/kg	3.31 $\pm$ 0.17    g/kg	10.3 $\pm$ 0.5    g/kg

Availability: 20 g whole seed in an aluminium plastic laminated sachet sealed under nitrogen.

Substance	<b>BCR-262R</b> Defatted peanut meal (blank)	<b>BCR-263R</b> Defatted peanut meal (medium level)	<b>BCR-264</b> Defatted peanut meal (high level)	<b>BCR-375</b> Compound feed (very low level blank)	<b>ERM-BE375</b> Compound feed (very low level)	<b>ERM-BE376</b> Compound feed (high level)
Aflatoxin B1 Aflatoxin B2 Aflatoxin G1 Aflatoxin G2 Sum of aflatoxin B1, B2, G1 and G2	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  < 3	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  17.1 $\pm$ 2.4 3.0 $\pm$ 0.4 3.0 $\pm$ 0.5 (0.62 $\pm$ 0.21) (23.7 $\pm$ 2.5)	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  206 $\pm$ 13	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  < 1	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  2.6 $\pm$ 0.4 0.20 $\pm$ 0.04 0.40 $\pm$ 0.10 < 0.2	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  12.9 $\pm$ 1.8 0.68 $\pm$ 0.10 5.2 $\pm$ 0.8

Values in brackets are not certified.

Availability: Sachets sealed under vacuum containing about 100 g (BCR-263R) and about 150 g (BCR-264) of finely ground defatted peanut meal.

BCR-262R is available in 280 mL amber glass bottles containing about 100 g of the peanut meal, additionally sealed in foil-laminate sachet.

BCR-375 is supplied in units of about 50 g of a finely ground compound feed, ERM-BE375 and ERM-BE376 consist of 2 bottles filled with about 75 g of compound feedingstuff each.

	Description	<b>Aflatoxin M<sub>1</sub></b> ( $\mu\text{g}/\text{kg}$ )
<b>ERM-BD282</b>	Whole milk powder (zero level)	< 0.02
<b>ERM-BD283</b>	Whole milk powder (low level)	0.111 $\pm$ 0.018
<b>ERM-BD284</b>	Whole milk powder (high level)	0.44 $\pm$ 0.06

Availability: The materials are provided in units of 30 g in amber glass bottles filled and sealed under nitrogen.

Substance	<b>BCR-401R</b> Peanut butter (very low level)	<b>BCR-385R</b> Peanut butter (low level)
Aflatoxin B1	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  < 0.2	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  1.77 $\pm$ 0.30
Aflatoxin B2	< 0.2	0.48 $\pm$ 0.08
Aflatoxin G1	< 0.2	0.9 $\pm$ 0.4
Aflatoxin G2	< 0.2	0.30 $\pm$ 0.12
Total		3.5 $\pm$ 0.5

Availability: The samples are supplied in units of about 100 g in aluminium cans.

Substance	<b>BCR-377</b> Maize Flour (very low level blank)	<b>BCR-396</b> Wheat Flour (very low level blank)
DON	<u>Mass fraction</u> (mg/kg)  < 0.05	<u>Mass fraction</u> (mg/kg)  < 0.05

Availability: Sachets sealed under vacuum containing about 150 g of sealed finely ground flour.

Substance	<b>BCR-471</b> Wheat (blank)
Ochratoxin A	<u>Mass fraction</u> ( $\mu\text{g}/\text{kg}$ )  < 0.6

Availability: Units of about 55 g in foil-laminate pouches sealed under vacuum.

Compound	BCR-543 Mussel		
	<u>Mass fraction</u> (mg/kg)		
Saxitoxin·2HCl Saxitoxin·2HCl (if enriched with BCR-663) dc-saxitoxin	< 0.07	0.48 ± < 0.04	0.06

Availability: BCR-543 is available in heat-sealed laminate sachets containing about 15 g of material each. BCR-663 is available in ampoules containing 1 mL.

Substance	ERM-BC716 Maize	ERM-BC717 Maize
	<u>Mass fraction</u> (µg/kg)	<u>Mass fraction</u> (µg/kg)
ZON	< 5	83 ± 9

Availability: ERM-BC716 and ERM-BC717 are supplied in sachets containing at least 60 g.

### 2.2.3 CERTIFIED FOR THE TOTAL ELEMENT CONTENT

Substance	ERM-BD150 Skimmed milk powder	ERM-BD151 Skimmed milk powder
	(g/kg)	(g/kg)
Ca	13.9 ± 0.8	13.9 ± 0.7
Cl	9.7 ± 2.0	9.8 ± 1.2
K	17.0 ± 0.7	17.0 ± 0.8
Mg	1.26 ± 0.10	1.26 ± 0.07
Na	4.18 ± 0.19	4.19 ± 0.23
	(mg/kg)	(mg/kg)
Cd	0.0114 ± 0.029	0.106 ± 0.013
Cu	1.08 ± 0.06	5.00 ± 0.23
Fe	4.6 ± 0.5	53 ± 4
Hg	0.060 ± 0.007	0.52 ± 0.04
I	1.73 ± 0.14	1.78 ± 0.17
Mn	0.289 ± 0.018	0.29 ± 0.03
Pb	0.019 ± 0.004	0.207 ± 0.014
Se	0.188 ± 0.014	0.19 ± 0.04
Zn	44.8 ± 2.0	44.9 ± 2.3

Availability: Glass bottles containing 20 g of skimmed milk powder.

Substance	ERM-BB184 Bovine muscle (mg/kg)	ERM-BB186 Pig kidney (mg/kg)	ERM-BB422 Fish muscle (mg/kg)	BCR-185R Bovine liver (mg/kg)
As	0.0234 ± 0.0026	(0.008 ± 0.006)	12.7 ± 0.7	0.0330 ± 0.0029
Cd	0.0022 ± 0.0004	1.09 ± 0.05	0.0075 ± 0.0018	0.544 ± 0.017
Cu	2.31 ± 0.09	36.5 ± 1.8	1.67 ± 0.16	277 ± 5
Fe	75 ± 4	255 ± 13	9.4 ± 1.4	-
Hg	(0.0018 ± 0.0010)	(0.023 ± 0.011)	0.601 ± 0.030	-
I	-	-	1.4 ± 0.4	-
Mn	0.276 ± 0.013	7.26 ± 0.25	0.368 ± 0.028	11.07 ± 0.29
Pb	-	0.040 ± 0.005	-	0.172 ± 0.009
Se	0.45 ± 0.04	10.3 ± 0.9	1.33 ± 0.13	1.68 ± 0.14
Zn	146 ± 7	134 ± 5	16.0 ± 1.1	138.6 ± 2.1

Values in brackets are not certified.

Availability: BCR-185R is provided in units of 15 g as lyophilised powders, in screw-capped glass bottles.

ERM-BB184 is provided in units of 7 g as lyophilised powder, in brown-glass vials.

ERM-BB186 and ERM-BB422 are provided in units of 10 g as lyophilised powders, in brown-glass vials

Substance	BCR-191 Brown bread		
Cd µg/kg	28.4	±	1.4
Cu mg/kg	2.63	±	0.07
Fe mg/kg	40.7	±	2.3
Mn mg/kg	20.3	±	0.7
Pb µg/kg	187	±	14
Zn mg/kg	19.5	±	0.5

Availability: BCR-191 is provided in units of 25 g.

Substance	IRMM-804 Rice flour (mg/kg)		
As	0.049	±	0.004
Cd	1.61	±	0.07
Cu	2.74	±	0.24
Mn	34.2	±	2.3
Pb	0.42	±	0.07
Zn	23.1	±	1.9

Availability: Amber glass bottles containing about 15 g of powdered material.

Substance	BCR-273			BCR-274		
	Single cell protein (mg/g)			Single cell protein (µg/g)		
As				0.132	±	0.014
Ca				0.030	±	0.002
Cd				0.039	±	0.003
Co				13.1	±	0.4
Cu						
Fe	11.97	±	0.14			
K	0.156	±	0.004			
Mn	2.22	±	0.05			
N	121.6	±	0.8	51.9	±	1.2
P	26.8	±	0.4			
Pb				0.044	±	0.010
Se				1.03	±	0.05
Zn				42.7	±	1.0

Availability: BCR-273 and BCR-274 are provided in units of 10 g as a dry powder in sealed glass ampoules.

Substance	BCR-679 White cabbage		
B mg/kg	(27.7	±	1.9)
Ba mg/kg	(10.3	±	0.6)
Ca mg/kg	(7768	±	655)
Cd mg/kg	1.66	±	0.07
Cr mg/kg	(0.6	±	0.1)
Cu mg/kg	2.89	±	0.12
Fe mg/kg	55.0	±	2.5
Hg µg/kg	6.3	±	1.4
Mg mg/kg	(1362	±	127)
Mn mg/kg	13.3	±	0.5
Mo mg/kg	14.8	±	0.5
Ni mg/kg	27.0	±	0.8
P mg/kg	(3307	±	241)
Sb µg/kg	20.6	±	2.6
Sr mg/kg	11.8	±	0.4
Tl µg/kg	3.0	±	0.3
Zn mg/kg	79.7	±	2.7

Values in brackets are not certified.

Availability: BCR-679 is provided in units of 15 g.

Substance	ERM-BC381 Rye Flour	ERM-BC382 Wheat Flour	BCR-383 Haricots Verts (Beans)	ERM-BB384 Lyophilised pork muscle
<u>Major components (g / 100 g)</u>				
Glucose			(12.4)	
Fructose			(4.6)	
Sucrose			(1.0)	
N (Kjeldahl)	1.562 ± 0.014	1.851 ± 0.017	1.05 ± 0.04	14.2 ± 0.4
Fat	1.36 ± 0.16	1.39 ± 0.17		8.99 ± 0.20
Starch <sup>1)</sup>	72.2 ± 1.9	81.2 ± 1.7		
Starch & Sugars <sup>2)</sup>			(78.9) (10.9)	
Dietary Fibre (Englyst)			11.9 ± 0.6	
Dietary Fibre (AOAC 1985/1988) <sup>3)</sup>				
Ash at 550 °C	1.08 ± 0.11	0.60 ± 0.10	2.39 ± 0.10	4.51 ± 0.19
<u>Essential elements (g/kg)</u>				
Ca	0.32 ± 0.04	0.210 ± 0.018	2.85 ± 0.23	0.164 ± 0.021
K	3.35 ± 0.11	1.88 ± 0.08	7.8 ± 0.4	
Mg	0.567 ± 0.013	0.247 ± 0.010	(0.9)	1.03 ± 0.04
Na			0.075 ± 0.007	1.86 ± 0.15
P	2.01 ± 0.07	1.19 ± 0.07	(1.8)	8.7 ± 0.5

Values in brackets are not certified.

Availability: ERM-BC381 and ERM-BC382: 100 ml amber vial containing 37 g flour; ERM-BB384: 2 vials of 9 g lyophilised material; BCR-383: 100 g of powdered material in food grade laminated plastic/aluminium sachets sealed under nitrogen.

<sup>1)</sup> Mass fraction of polysaccharide in dry matter

<sup>2)</sup> Mass fraction of monosaccharides in dry matter

<sup>3)</sup> Prosky L. et al., J Assoc Off Anal Chem (1985) 68: 677-679, (1988) 71: 1017-1023

## 2.2.4 CERTIFIED FOR PROXIMATES AND CONVENTIONAL PROPERTIES

Substance	BCR-162R Soya-maize oil blend		
<u>Methyl ester of</u>	<u>Mass fraction fatty acid methyl ester / Total fatty acid methyl ester (g/100 g)</u>		
16:0 n-hexadecanoic acid	10.74	±	0.16
18:0 n-octadecanoic acid	2.82	±	0.04
18:1 n-octadecenoic acids		(26.5)	
18:2 n-octadecadienoic acids		(54.68)	
18:3 n-octadecatrienoic acids		(3.80)	
9c-18:1 n-octadecenoic acid	25.4	±	0.4
9c,12c-18:2 n-octadecadienoic acid	54.13	±	0.25
9c,12c,15c-18:3 n-octadecatrienoic acid	3.35	±	0.05

Values in brackets are not certified.

Availability: Each unit contains approximately 5.5 g soya-maize oil blend in 10 mL amber ampoule.

Substance	BCR-163 Beef-pork fat oil blend		
<u>Methyl ester of</u>	<u>Mass fraction fatty acid methyl ester / Total fatty acid methyl ester <sup>1)</sup> (g/100 g)</u>		
14:0 n-tetradecanoic acids	2.29	±	0.04
16:0 n-hexadecanoic acids	25.96	±	0.30
16:1 n-hexadecenoic acids	2.58	±	0.16
18:0 n-octadecanoic acids	18.29	±	0.17
18:1 n-octadecenoic acids	38.3	±	0.4
18:2 n-octadecadienoic acids	7.05	±	0.17
18:3 n-octadecatrienoic acids	0.86	±	0.14
<u>Sterols</u>	<u>Mass fraction (mg/100 g) in fat</u>		
Cholesterol	134	±	5

Availability: In units of 2 × 5 mL in dark glass ampoules sealed under nitrogen.

1) Includes any geometric (i.e. cis/trans) and positional isomers, expressed as mass fraction of total fatty acid (methyl esters) derived from triglycerides.

2) These components are included in the Certified Value for this group of fatty acids.

The report gives additional indicative values: Fatty Acids and "Total" Sterol Mass Fraction.

Component	BCR-633 Tracers in anhydrous butter fat (mg/kg)		
$\beta$ -Apo-8'-carotenic acid ethyl ester	26.5	$\pm$	1.4
$\beta$ -Sitosterol	530	$\pm$	29
Stigmasterol	147	$\pm$	11
<i>n</i> -Heptanoic acid triglyceride	1.06 * 10 <sup>3</sup>	$\pm$	0.04 * 10 <sup>3</sup>

Availability: BCR-633 is supplied in units of about 5 g each in amber glass ampoules, which were filled under inert gas conditions (nitrogen).

Component	IRMM-801 Cocoa Butter (%)		
1,3-dipalmitoyl-2-oleyl-glycerol	18.14	$\pm$	0.26
1-palmitoyl-2-oleoyl-3-stearoyl-glycerol	44.68	$\pm$	0.30
1,2-dioleoyl-3-palmitoyl-glycerol	2.26	$\pm$	0.16
1,3-distearoyl-2-oleoyl-glycerol	31.63	$\pm$	0.29
1,2-dioleoyl-3-stearoyl-glycerol	3.29	$\pm$	0.17

Availability: IRMM-801 is supplied in units of 5 g in brown amber glass ampoules sealed under argon/helium.

Component	BCR-519 Anhydrous butter fat (%)		
Cholesterol	0.30	$\pm$	0.03
C24	0.05	$\pm$	0.02
C26	0.25	$\pm$	0.03
C28	0.59	$\pm$	0.04
C30	1.15	$\pm$	0.05
C32	2.43	$\pm$	0.12
C34	5.64	$\pm$	0.18
C36	10.47	$\pm$	0.19
C38	12.53	$\pm$	0.22
C40	10.03	$\pm$	0.16
C42	6.69	$\pm$	0.10
C44	6.11	$\pm$	0.08
C46	6.86	$\pm$	0.08
C48	8.69	$\pm$	0.15
C50	11.40	$\pm$	0.24
C52	10.96	$\pm$	0.25
C54	5.89	$\pm$	0.13

Availability: The material consists of anhydrous butter fat and is supplied in units of two glass ampoules sealed under nitrogen, each containing approximately 5 mL fat. At normal ambient temperature BCR-519 is solid.

Compound	BCR-632 Butter fat					
	BCR-632A Pure butter fat (g/100 g)			BCR-632B Adulterated butter fat (g/100 g)		
C24	0.07	$\pm$	0.04	0.08	$\pm$	0.04
Cholesterol	0.289	$\pm$	0.012	0.278	$\pm$	0.011
C26	0.33	$\pm$	0.06	0.34	$\pm$	0.06
C28	0.74	$\pm$	0.07	0.75	$\pm$	0.06
C30	1.37	$\pm$	0.08	1.46	$\pm$	0.07
C32	2.83	$\pm$	0.14	3.30	$\pm$	0.12
C34	6.09	$\pm$	0.29	6.57	$\pm$	0.25
C36	10.7	$\pm$	0.5	11.1	$\pm$	0.4
C38	12.5	$\pm$	0.4	12.7	$\pm$	0.4
C40	10.05	$\pm$	0.19	10.07	$\pm$	0.17
C42	7.07	$\pm$	0.13	7.10	$\pm$	0.10
C44	6.68	$\pm$	0.12	6.57	$\pm$	0.12
C46	7.36	$\pm$	0.17	7.12	$\pm$	0.17
C48	8.74	$\pm$	0.21	8.42	$\pm$	0.19
C50	10.74	$\pm$	0.24	10.28	$\pm$	0.19
C52	9.8	$\pm$	0.4	9.36	$\pm$	0.28
C54	4.7	$\pm$	0.5	4.5	$\pm$	0.4

Availability: BCR-632 is available as a set of BCR-632A and BCR-632B: 2 amber glass ampoules each with approximately 5 mL.

	<b>BCR-121</b> Wholemeal flour (mg/kg)	<b>BCR-122</b> Margarine (mg/kg)	<b>BCR-431</b> Brussels sprouts (mg/kg)	<b>BCR-485</b> Mixed vegetables (mg/kg)	<b>BCR-487</b> Pig's liver (mg/kg)	<b>ERM-BD600</b> Whole milk powder (mg/kg)
A (all-trans-retinol)					3.8	± 0.6
A (all-trans-retinol and 13-cis-retinol)					4.1	± 0.8
B <sub>1</sub> (thiamin)	4.63 ± 0.39			3.07 ± 0.34	8.6	± 1.1
B <sub>2</sub> (riboflavin)				106.8 ± 5.6	4.5	± 0.6
B <sub>6</sub> (total pyridoxine)	4.10 ± 1.02			4.8 ± 0.8	16.7	± 1.4
B <sub>12</sub> (cyanocobalamin)		0.125 ± 0.007	4830 ± 240	19.3 ± 2.9		
C (total ascorbate)		241 ± 12		1.12 ± 0.09	0.32	± 0.07
D <sub>3</sub> (cholecalciferol)					74	± 11
E (tocopherol)						
Folate (total)	0.50 ± 0.07			86	± 15	
Niacin						
Trans-α-carotene						
Trans-β-carotene						
Total-α-carotene						
Total-β-carotene						
Lutein						
Lutein + zeaxanthin						
5-methyltetrahydro-folic acid (5-MTHF)						
				(2.14 ± 0.42)		

Values in brackets are not certified;

Availability: BCR-121: about 50 g unit size.

BCR-122: can filled with about 200 g.

BCR-431: about 20 g lyophilised and powdered material in food grade plastic/aluminium laminated sachets under nitrogen.

BCR-485: about 25 g unit size.

BCR-487: about 15 g unit size.

ERM-BD600: sachet with about 100 g.

	Description	Amylose mass fraction (g/100 g)
<b>BCR-465</b>	Rice flour (low level)	15.40 ± 0.30
<b>BCR-466</b>	Rice flour (medium level)	23.1 ± 0.5
<b>BCR-467</b>	Rice flour (high level)	27.7 ± 0.8

Availability: 10 g of ground rice flour in vacuum sealed laminated polyester/aluminium/polyethylene sachets.

Substance	<b>BCR-644</b> Artificial foodstuff	<b>BCR-645</b> Artificial foodstuff
<u>Mass fraction on dry mass basis (g / 100 g)</u>		
Fructose	16.2 ± 1.1	
Sucrose	10.81 ± 0.25	26.2 ± 0.8
Lactose	15.85 ± 0.29	27.8 ± 0.6
Starch / glucose	35.1 ± 1.2	25.2 ± 0.9

Availability: BCR-644 and BCR-645 are supplied in units of approximately 50 g in 125 mL amber glass bottles.

	<b>BCR-651</b> Beer (% ethanol v/v)	<b>BCR-652</b> Beer (% ethanol v/v)	<b>BCR-653</b> Wine (% ethanol v/v)
Ethanol	0.505 ± 0.006	0.051 ± 0.002	0.539 ± 0.007

Availability: Amber glass ampoule, flushed with nitrogen, containing 10 mL of sample.

Substance	ERM-BC381 Rye Flour	ERM-BC382 Wheat Flour	BCR-383 Haricots Verts (Beans)	ERM-BB384 Lyophilised pork muscle
<u>Major components (g / 100 g)</u>				
Glucose			(12.4)	
Fructose			(4.6)	
Sucrose			(1.0)	
N (Kjeldahl)	1.562 ± 0.014	1.851 ± 0.017	1.05 ± 0.04	14.2 ± 0.4
Fat	1.36 ± 0.16	1.39 ± 0.17		8.99 ± 0.20
Starch <sup>1)</sup>	72.2 ± 1.9	81.2 ± 1.7		
Starch & Sugars <sup>2)</sup>			(78.9) (10.9)	
Dietary Fibre (Englyst)			11.9 ± 0.6	
Dietary Fibre (AOAC 1985/1988) <sup>3)</sup>				
Ash at 550 °C	1.08 ± 0.11	0.60 ± 0.10	2.39 ± 0.10	4.51 ± 0.19
<u>Essential elements (g/kg)</u>				
Ca	0.32 ± 0.04	0.210 ± 0.018	2.85 ± 0.23	0.164 ± 0.021
K	3.35 ± 0.11	1.88 ± 0.08	7.8 ± 0.4	
Mg	0.567 ± 0.013	0.247 ± 0.010	(0.9)	1.03 ± 0.04
Na			0.075 ± 0.007	1.86 ± 0.15
P	2.01 ± 0.07	1.19 ± 0.07	(1.8)	8.7 ± 0.5

Values in brackets are not certified.

Availability: ERM-BC381 and ERM-BC382: 100 ml amber vial containing 37 g flour; ERM-BB384: 2 vials of 9 g lyophilised material; BCR-383: 100 g of powdered material in food grade laminated plastic/aluminium sachets sealed under nitrogen.

<sup>1)</sup> Mass fraction of polysaccharide in dry matter

<sup>2)</sup> Mass fraction of monosaccharides in dry matter

<sup>3)</sup> Prosky L. et al., J Assoc Off Anal Chem (1985) 68: 677-679, (1988) 71: 1017-1023

Substance	BCR-380R Whole milk powder (g / 100 g)	BCR-685 Skim milk powder (g / 100 g)
<u>Mass fraction</u>		
Crude protein (Kjeldahl-N x 6.38)	28.66 ± 0.28	38.2 ± 0.4
Fat	26.95 ± 0.16	0.96 ± 0.12
Lactose (anhydrous)	37.1 ± 1.0	
Ash	6.00 ± 0.13	

Availability: BCR-380R is supplied in units of about 100 g, BCR-685 in units of about 50 g, in amber glass bottles.

Quantity	BCR-446 Low oil content rapeseed	BCR-447 Medium oil content rapeseed
	<u>Mass fraction (g/100 g)</u>	<u>Mass fraction (g/100 g)</u>
'As is' Oil	39.49 ± 0.15	41.99 ± 0.15
Moisture and volatiles	7.01 ± 0.07	7.42 ± 0.07
'Dry basis' Oil	42.48 ± 0.15	45.36 ± 0.15

Availability: About 150 g of rapeseed in a specially laminated bag containing an oxygen absorber and sealed under dry argon.

Substance	BCR-708 Dairy feed	BCR-709 Pig feed	
<u>Mass fraction</u>			
Crude protein	240 ± 12	199 ± 5	g/kg
Crude oils and fats	65 ± 8	51 ± 14	g/kg
Crude fibre	93 ± 14	56 ± 12	g/kg
Crude ash	50.0 ± 3.0	42 ± 4	g/kg
Ca	4.8 ± 0.5	1.05 ± 0.16	g/kg
Cu	37 ± 4	173 ± 25	mg/kg
Mg	1.47 ± 0.22	1.89 ± 0.30	g/kg
P	4.7 ± 0.4	5.4 ± 0.7	g/kg

Availability: 100 mL amber glass bottle containing about 40 g of material.

Substance	ERM-BC514 Haricot beans (g/kg)	ERM-BC515 Carrot (g/kg)	ERM-BC516 Apple (g/kg)	ERM-BC517 Full fat soya flour (g/kg)	ERM-BD518 Bran breakfast cereal (g/kg)
<u>Dietary Fibres</u>					
Methods used:					
AOAC 1990	256 ± 5	311 ± 6	164 ± 4	126 ± 5	302 ± 8
Englyst (by gas chromatography)	198 ± 10	271 ± 6	137 ± 5	119 ± 7	241 ± 8
Uppsala	237 ± 15	298 ± 11	162 ± 8	128 ± 9	276 ± 18
AOAC 1992 MES-TRIS	259 ± 15	295 ± 4	149 ± 10	124 ± 21	305 ± 6
Englyst (by colorimetry)	201 ± 6	252 ± 12	134 ± 5	123 ± 8	250 ± 11

Availability: ERM-BC514 to BC517 and ERM-BD518 are supplied in units of approximately 25 g in food grade glass bottles sealed under vacuum.

	BCR-537 Plastic film A (mg/dm <sup>2</sup> )	BCR-538 Plastic film B (mg/dm <sup>2</sup> )	BCR-539 Plastic film C (mg/dm <sup>2</sup> )
Overall migration by total immersion in olive oil 10 days at 40 °C	8.3 ± 1.0		
Overall migration by single sided cell in olive oil 10 days at 40 °C		5.7 ± 0.7	
Overall migration by pouch in olive oil 10 days at 40 °C			6.1 ± 1.0

Availability: PET/foil/PE heat sealed pouches containing double thickness sheets of additive free linear low density polyethylene of (33 x 22.5) cm for BCR-537, (125 x 22.5) cm for BCR-538 and (45 x 25) cm for BCR-539.

Parameters (determined according to ISO and ICC standards)	BCR-563 Common wheat flour	
Protein content g/100 g dry matter basis	11.71	± 0.13
Ash content % dry matter basis	0.562	± 0.008
Falling number / s	319	± 15
Zeleny sedimentation volume / mL	44	± 1
<u>CHOPIN ALVEOGRAPH</u>		
P/mmH <sub>2</sub> O*	80.8	± 2.1
L/mm**	109	± 7.2
P/L	0.8	± 0.1
W/(10 <sup>-4</sup> Joules)***	289.9	± 10.4
<u>BRABENDER FARINOGRAPH</u>		
Maximum consistency / BU (Brabender Units)	499	± 5
Development time / min.	1.7	± 0.3
Stability / min.	2.5	± 0.4
Degree of softening / BU (Brabender Units)	87	± 9
<u>BRABENDER EXTENSOGRAPH</u>		
Maximum resistance / EU (Brabender Extensograph Units)	446	± 39
Resistance at 50 mm / EU (Brabender Extensograph Units)	261	± 30
Energy / (cm <sup>2</sup> )	119	± 14
Extensibility / mm	202	± 9
MOISTURE MASS FRACTION [g/100 g]	13.95	± 0.04

Availability: BCR-563 is provided in vacuum sealed laminated foil envelopes in units of approximately 360 g.

\* P: Mean curve height of Alveograph.

\*\* L: Mean curve length.

\*\*\* W: Area under Alveograph curve.

## 2.2.5 CERTIFIED FOR MICROBIOLOGICAL PROPERTIES

### BCR-528: Capsules filled with milk powder artificially contaminated by *Bacillus cereus* (ATCL 9139)

Procedure	Number of colony forming particles in one analytical portion <sup>1)</sup>		
	Certified value <sup>2)</sup>	95 % confidence limits	
MEYP (ISO 7932) after 24 h incubation	53.4	51.7	- 55.2
MEYP (ISO 7932) after 48 h incubation	53.7	52.1	- 55.4
PEMBA (L 00.00 - 25) <sup>3)</sup> after 24 h incubation	55.0	52.8	- 57.4
PEMBA (L 00.00 - 25) <sup>3)</sup> after 48 h incubation	55.8	53.6	- 58.0

Availability: BCR-528 is provided in containers holding 10 gelatine capsules filled with artificially contaminated milk powder.

- 1) Number of colony forming particles of *Bacillus cereus* determined in one analytical portion. Analytical portion: A volume of (0.100 ± 0.002) mL from 10 mL peptone saline solution in which one capsule has been reconstituted.
- 2) This value is the geometric mean of 11 accepted sets of data, independently obtained by 11 laboratories.
- 3) German Federal Food Law method number.

### IRMM-311: Genomic DNA (gDNA) of *Bacillus licheniformis* DSM 5749 in agarose inserts for Pulsed Field Gel Electrophoresis (PFGE)

<i>Sfi</i> digested DNA fragments in the size interval 50 kb – 90 kb		Fragment length (kb)		
Band no	1	89.6	±	4.7
	2	80.9	±	2.5
	3	75.3	±	2.7
	4	72.2	±	3.5
	5	66.9	±	1.9
	6	64.6	±	2.9
	7	60.3	±	1.3
	8	56.5	±	1.3
	9	53.9	±	1.3
	10	50.6	±	1.3

Availability: Each vial contains one agarose insert of undigested genomic DNA of *Bacillus licheniformis* DSM 5749 for PFGE.

### IRMM-312: Genomic DNA (gDNA) of *Bacillus subtilis* DSM 5750 in agarose inserts for Pulsed Field Gel Electrophoresis (PFGE)

<i>Sfi</i> digested DNA fragments in the size interval 50 kb – 90 kb		Fragment length (kb)		
Band no	1	89.2	±	0.9
	2	81.4	±	0.8
	3	77.7	±	0.6
	4	62.5	±	1.8
	5	59.5	±	2.1
	6	44.0	±	2.4
	7	29.2	±	2.0
	8	23.6	±	1.3
	9	18.6	±	1.3

Availability: Each vial contains one agarose insert of undigested genomic DNA of *Bacillus subtilis* DSM 5750 for PFGE.

**IRMM-313: Genomic DNA (gDNA) of *Campylobacter coli* (CNET068) and *Campylobacter jejuni* (CNET112) in agarose inserts for Pulsed Field Gel Electrophoresis (PFGE)**

SmaI digested DNA fragments		DNA fragment sizes (kb)		
Fragment no	2	458.8	±	2.0
	3	351.7	±	2.4
	4	303.0	±	2.3
	5	263.2	±	1.9
	6	188.2	±	1.2
	7	173.2	±	1.3
	8	131.1	±	1.5
	9	114.4	±	1.2
	10	95.5	±	1.4
	11	81.2	±	1.7
	12	54.9	±	2.2
	13	40.7	±	1.6
	14	(25.4	±	1.3)
	15	(17.6	±	0.3)
	16	(10.9	±	0.4)

Values in brackets are not certified.

Availability: Each vial contains one agarose plug for PFGE with undigested genomic DNA of *Campylobacter coli* CNET068 and *Campylobacter jejuni* CNET112 embedded.

**IRMM-351: Escherichia coli 0157 in material spheres**

	Number of colony forming unit (cfu)		
cfu per material sphere on nutrient agar	4	±	2
cfu per material sphere on enterohemolysin agar	4	±	2

Availability: Each vial contains one material sphere of Escherichia coli 0157 (NCTC 12900).

**IRMM-352: Salmonella enteritidis in material spheres**

	Number of colony forming unit (cfu)		
cfu per material sphere on nutrient agar	5	±	2
cfu per material sphere on xylose lysine deoxycholate agar	4	±	2

Availability: Each vial contains one material sphere of Salmonella enteritidis (NCTC 12694).

**IRMM-354: Candida albicans in material spheres**

	Number of colony forming unit (cfu)		
cfu per material sphere on nutrient agar	917	±	168
cfu per material sphere on Oxytetracyclin-Glucose-Yeast Extract agar (OGYE)	912	±	173

Availability: Each vial contains one material sphere of Candida albicans (NCIPF 3179).

## IRMM-355: Enterococcus faecalis in material spheres

	Number of colony forming unit (cfu)		
cfu per material sphere on horse blood agar	890	±	135
cfu per material sphere on Slanetz and Bartley agar	823	±	126

Availability: Each vial contains one material sphere of *Enterococcus faecalis* (CIP 106877).

## IRMM-447: Genomic DNA (gDNA) of *Listeria monocytogenes* (strain 4B, NCTC 11994) with certified identity

	Indicative value (µg)		
Mass of genomic DNA <i>Listeria monocytogenes</i> per vial	(1.1	±	0.7)

Values in brackets are not certified.

Availability: Vial containing approximately 1.1 µg genomic DNA in lyophilised form and closed under argon atmosphere.

## IRMM-448: Genomic DNA (gDNA) of *Campylobacter Jejuni* (NCTC 11351) with certified identity

Property			
Identity	Confirmed by ceuE gene		
Mass of genomic DNA per vial (ng)	(71	±	39)

Values in brackets are not certified.

Availability: Vial containing approximately 71 ng genomic DNA in lyophilised form and closed under argon atmosphere.

## IRMM-449: Genomic DNA (gDNA) of *Escherichia coli* O157 (strain EDL 933) with certified identity

	Indicative value (µg)		
Mass of genomic DNA <i>Escherichia coli</i> per vial	(1.3	±	0.7)

Values in brackets are not certified.

Availability: Vial containing approximately 1.3 µg genomic DNA in lyophilised form and closed under argon atmosphere.

## 2.2.6 CERTIFIED FOR VETERINARY DRUGS

	Description	Substance	Hormones in lyophilised bovine urine Mass concentration in reconstituted sample (µg/L)		
<b>BCR-386<sup>(1)</sup></b> <b>BCR-387<sup>(1)</sup></b> <b>BCR-388<sup>(1)</sup></b> <b>BCR-390 (RM)<sup>(1)</sup></b> <b>BCR-391<sup>(1)</sup></b>	Bovine urine	Diethylstilboestrol (DES)	< 0.1		
	Bovine urine	Dienoestrol (DE)	< 0.1		
	Bovine urine	Hexoestrol (HEX)	< 0.1		
	Bovine urine	Dienoestrol (DE)	(34)		
	Bovine urine	Hexoestrol (HEX)	13.3	±	3.1
			Content	Relevant below the certified value	Relevant above the certified value
<b>BCR-502<sup>(2)</sup></b>	Bovine urine	Clenbuterol	< 0.1		
		Salbutamol	< 0.2		
<b>BCR-503<sup>(2)</sup></b>	Bovine urine	Clenbuterol	2.5	0.4	0.4
		Salbutamol	2.3	0.6	0.9
<b>BCR-504<sup>(2)</sup></b>	Bovine urine	Clenbuterol	6.0	0.5	0.7
		Salbutamol	5.6	1.1	1.9

Value in brackets is not certified.

Availability: <sup>(1)</sup> Units of lyophilised urine equivalent to about 2.0 mL in vials sealed under nitrogen.

<sup>(2)</sup> Units of lyophilised urine equivalent to about 5.0 mL in vials sealed under nitrogen.

	Description	Substance		Hormones in lyophilised bovine urine Mass concentration in reconstituted sample ( $\mu\text{g}/\text{kg}$ )		
<b>ERM-BB386<sup>(1)</sup></b>	Bovine urine	Diethylstilboestrol (DES)		< 0.6		
<b>ERM-BB389<sup>(2)</sup></b>	Bovine urine	Dienoestrol (DE)		< 0.6		
		Hexoestrol (HEX)		< 0.4		
		Diethylstilboestrol (DES)	1.1	$\pm$	0.5	
		Dienoestrol (DE)	5.5	$\pm$	1.4	
		Hexoestrol (HEX)	6.1	$\pm$	0.9	

Availability: <sup>(1)</sup> Vial containing approximately 0.36 g lyophilised bovine urine corresponding to 5.20 g of fresh bovine urine.

<sup>(2)</sup> Vial containing approximately 0.31 g lyophilised bovine urine corresponding to 5.18 g of fresh bovine urine.

	Description	Substance		Mass fraction in reconstituted sample ( $\mu\text{g}/\text{kg}$ ) <sup>*</sup>		
<b>BCR-648</b>	Bovine liver	Clenbuterol		< 0.5		
<b>BCR-649</b>	Bovine liver	Clenbuterol	1.2	$\pm$	0.3	
<b>BCR-474</b>	Bovine liver	17 $\alpha$ -trenbolone		< 0.5		
<b>BCR-475</b>	Bovine liver	17 $\alpha$ -trenbolone	7.6	$\pm$	2.2	
<b>BCR-411</b>	Bovine muscle	Diethylstilboestrol		> 0.5		
<b>BCR-412</b>	Bovine muscle	Diethylstilboestrol		< 0.1		
<b>BCR-673</b>	Bovine eye	Clenbuterol		< 0.5		
<b>BCR-674</b>	Bovine eye	Clenbuterol	9.4	$\pm$	1.1	

Availability: BCR-648 and -649 are provided in units of 10 g lyophilised bovine liver in vials sealed under argon. BCR-474 and -475 are sold as set and provided in brown glass vials in units of 2.8 g lyophilised liver corresponding to 10 g fresh liver. BCR-411 and -412 are provided brown glass vials as lyophilised bovine muscle in units equivalent to about 5 g of fresh bovine tissue. BCR-673 and BCR-674 are provided in brown glass vials containing about 0.1 g of material.

	<b>BCR-444</b> Porcine muscle (blank) ( $\mu\text{g}/\text{kg}$ )	<b>ERM-BB130</b> Pork muscle ( $\mu\text{g}/\text{kg}$ )	<b>BCR-445</b> Porcine muscle (incurred) ( $\mu\text{g}/\text{kg}$ )
Chloramphenicol	< 0.2	0.230 $\pm$ 0.021	8.9 $\pm$ 0.9

Availability: BCR-444 and BCR-445 are provided in brown glass vials, ERM-BB130 in an amber glass bottle, each containing about 7 g of lyophilised pork muscle tissue.

	Description	Substance		Mass fraction in reconstituted sample (mg/kg)		
<b>BCR-695</b>	Pig liver	Chlortetracycline		< 0.004		
<b>BCR-696</b>	Pig liver	Chlortetracycline	0.58	$\pm$	0.11	
<b>BCR-697</b>	Pig muscle	Chlortetracycline		< 0.006		
<b>BCR-706</b>	Pig kidney	Chlortetracycline		< 0.005		
<b>BCR-707</b>	Pig kidney	Chlortetracycline	1.30	$\pm$	0.20	

Availability: These CRMs are provided in sealed glass vials containing lyophilised tissue equivalent to 5 g of fresh tissue.

	Description	Substance		( $\mu\text{g}/\text{kg}$ )		
<b>BCR-492</b>	Milk powder	Oxytetracycline	307	$\pm$	14	
<b>BCR-493</b>	Milk powder	Oxytetracycline		< 10		
<b>ERM-BB492</b>	Milk powder	Sum of oxytetracycline and 4-epi-oxytetracycline	101	$\pm$	11	
<b>ERM-BB493</b>	Milk powder	Sum of oxytetracycline and 4-epi-oxytetracycline		< 5		

Availability: BCR-492 and BCR-493: Units of lyophilised milk equivalent to 10 mL sealed under dry nitrogen; sold as set of 1 unit BCR-492 and BCR-493 each.

ERM-BB492 and ERM-BB493: Units of approximately 5.5 g of spray-dried partially skimmed milk; sold as set of 1 unit ERM-BB492 and ERM-BB493 each.

Substance	BCR-725 Salmon tissue ( $\mu\text{g/kg}$ )		
Flumequine Oxolinic acid	1170	$\pm$	210
	600	$\pm$	100

Availability: BCR-725 is provided in amber glass vials containing 2.2 g of lyophilised salmon tissue material.

	ERM-BB124 Pork muscle		
Nitroimidazoles in the reconstituted material	Mass fraction ( $\mu\text{g/kg}$ )		
Ronidazole (RNZ)	2.09	$\pm$	0.25
Metronidazole (MNZ)	1.93	$\pm$	0.15
2-hydroxymethyl-1-methyl-5-nitroimidazole (HMMNI)	0.69	$\pm$	0.09
Hydroxymetronidazole (MNZOH)	6.2	$\pm$	0.9
Hydroxipronidazole (IPZOH)	1.67	$\pm$	0.12
Dimetridazole (DMZ)	< 0.25		

Availability: ERM-BB124 is provided in amber glass bottles containing 10 g of lyophilised pork muscle tissue.

## 2.2.7 CERTIFIED FOR IDENTITY

### Ewes'/Goats' Curd (BCR-599)

BCR-599 consists of a set of two freeze dried curd materials made from a mixture of ewes' and goats' milk, intended to detect adulteration by cows' milk in cheeses made from ewes' milk, goats' milk and mixtures thereof, according to the reference method described in Commission Regulation (EC) No. 1081/96. The 0 % material is not adulterated, the 1 % material is adulterated milk 1 % of cows' milk.

Availability: BCR-599 is available as a set of two brown glass vials containing each about 15 g of lyophilised curd powder under Argon atmosphere.

### Peanut Test Material Kit (IRMM-481)

The IRMM peanut test material is **not** a reference material, because it could not be tested for homogeneity and stability. The intention is to make a standard peanut matrix available to the research community who may wish to conduct wider studies on similar matrices pre-treated in different ways. At present only the nominal particle size and the pre-treatment applied to the peanuts before milling can be guaranteed as well as the tolerances of weighing of  $1.000 \pm 0.015$  g of each variety in the peanut mixture provided in the kit.

	Peanut variety, origin	Peanut treatment	Nominal net weight of peanut powder	Colour code on cap
IRMM-481a	Runners, Argentina	blanched, strong air-roasting	2 g	blue
IRMM-481b	Common Natal, South-Africa	raw, mild air-roasting	2 g	green
IRMM-481c	Virginia, USA,	blanched, strong oil roasting	2 g	gold
IRMM-481d	Virginia, China	blanched, mild oil-roasting	2 g	red
IRMM-481e	Jumbo Runners, USA,	blanched only	2 g	brown
IRMM-481f	Mixture of 481a to 481e	all above	5 g (1 g of each variety 481a - 481e)	silver

Availability: IRMM-481 consists of a kit with six different vials containing non-salted peanut powders with a nominal particle size from 0.5 to 1.0 mm. Five of the vials are filled with approximately 2 g of each variety. The sixth vial contains a mixture of all five varieties provided in IRMM-481a to 481e. All vials have been filled with argon prior to capping to provide a protective atmosphere.

## Calibration kit for ruminant detection by PCR

The calibration kit for ruminant detection by PCR is **not** a certified reference material because the copy number concentration values have only been determined in a single laboratory and the material have not yet been tested for long term stability. Consequently, the DNA copy number concentrations are provided as indicative values and not as certified values.

The materials are intended for the determination of a cut-off value to discriminate positive samples (containing the ruminant target sequence) from negative samples by quantitative PCR. As any reference material (RM), the materials can also be used for control charts or precision studies.

	<b>Copy number concentration of the plasmid</b>
	<b>Indicative value [cp/µL]</b>
IRMM-AD482a	128
IRMM-AD482b	32
IRMM-AD482c	8

Availability: IRMM-AD482 consists of a kit of three different vials containing plasmid solutions bearing a ruminant DNA target with indicative copy number concentrations of 128 cp/µL, 32 cp/µL and 8 cp/µL. The vials contain at least 1 mL of plasmid solution with Tris-EDTA buffer and maize genomic DNA at a concentration of 12 ng/µL.

## 2.2.8 OTHERS

Substance	ERM-BD273 Toasted bread
	<u>Mass fraction</u> (ng/g)
Acrylamide	425 ± 29

Availability: ERM-BD273 is available in a brown glass vial containing about 30 g of toasted bread powder.

## 3 MATERIALS RELATED TO CLINICAL CHEMISTRY

### 3.1 PURE STANDARDS AND SYNTHETIC MATERIALS

	Description	Purity (%)
BCR-546	Formaldehyde 2,4-dinitrophenylhydrazone	> 99.3
BCR-547	Acetaldehyde 2,4-dinitrophenylhydrazone	98.3 ± 0.5
BCR-548	Acrolein 2,4-dinitrophenylhydrazone	> 97.9
BCR-549	Acetone 2,4-dinitrophenylhydrazone	> 99.6
BCR-550	Glutaraldehyde 2,4-dinitrophenylhydrazone	> 98.1

Availability: Approximately 10 mg of crystals in glass vials.

Compounds	<b>BCR-551</b> Acetonitrile solution Mass concentration (µg/mL)	<b>BCR-552</b> Acetonitrile solution (blank) (µg/mL)
Formaldehyde 2,4-Dinitrophenylhydrazone	2.94 ± 0.05	< 0.08
Acetaldehyde 2,4-dinitrophenylhydrazone	4.89 ± 0.07	< 0.05
Acrolein 2,4-dinitrophenylhydrazone	0.483 ± 0.011	< 0.04
Acetone 2,4-dinitrophenylhydrazone	4.96 ± 0.07	< 0.05

Availability: Set BCR-551-2 consists of 4 samples of BCR-551 and 1 sample of BCR-552.

	<b>BCR-553</b> Glass fibre filters Spiked mass per filter (expressed as µg formaldehyde)	<b>BCR-554</b> Glass fibre filters Mass per filter (blank) (expressed as µg formaldehyde)
Formaldehyde 2,4-dinitrophenylhydrazone on glass fibre filters	4.96 ± 0.06	< 0.1

Availability: Set BCR-553-4 consists of 2 samples of BCR-553 and 1 sample of BCR-554.

Substance	<b>BCR-555</b> Chlorinated hydrocarbons on Tenax (ng)		
Dichloromethane	320	±	40
1,1,1-Trichloroethane	370	±	40
Trichloroethylene	390	±	40
Perchloroethylene	327	±	17
Toluene	57	±	7

Availability: Stainless steel tube of 9.0 cm length and 0.25 inches outer diameter containing a single section of 250 mg TENAX GR, charged with 4 chlorinated hydrocarbons and toluene at the levels shown above.

	Description	Latex spheres Parameters of the calibration line		
<b>BCR-165</b>	Nominal 2 µm latex (0.02 % solids)	2.223	±	0.013
<b>BCR-166</b>	Nominal 4.8 µm latex (0.2 % solids)	4.821	±	0.019
<b>BCR-167</b>	Nominal 9.6 µm latex (1.4 % solids)	9.475	±	0.018

Availability: Vials containing 2 mL of an aqueous suspension of latex spheres.

	Thyroxine (T <sub>4</sub> )	3,3',5-triiodothyronine (T <sub>3</sub> )		
<b>IRMM-468</b>	98.6 (1.50)	±	0.7 0.12)	(0.51 97.1 ± 0.17) 0.7
<b>IRMM-469</b>				

Values in brackets are not certified.

Availability: The material consists of an off-white crystalline powder in an amber glass vial sealed under N<sub>2</sub> atmosphere. Each vial contains about 100 mg of the powder.

## 3.2 MATRIX MATERIALS

### 3.2.1 CERTIFIED FOR THE HORMONE CONTENT

#### Cortisol reference panel of fresh frozen human sera ERM-DA451/IFCC

Serum No.	Certified value nmol/L	Uncertainty nmol/L	Serum No.	Certified value nmol/L	Uncertainty nmol/L
1	361	14	18	146	6
2	432	17	19	166	7
3	288	11	20	83	4
4	152	6	21	89	4
5	329	13	22	180	7
6	278	11	23	387	15
7	515	20	24	384	15
8	163	7	25	315	12
9	287	11	26	215	9
10	230	9	27	497	19
11	334	13	28	299	12
12	261	10	29	265	11
13	430	17	30	114	5
14	626	24	31	764	29
15	246	10	32	623	24
16	211	8	33	264	10
17	366	14	34	390	15

Availability: As panel of 34 x 1 mL serum in screw capped cryo-vials.

		Cortisol in human serum (concentration in the reconstituted material <sup>1)</sup> )						
		( $\mu\text{g/L}$ )		( $\text{nmol/L}$ )				
ERM-DA192		98.8 277	$\pm$ $\pm$	2.0 5	$\pm$ $\pm$	273 763	$\pm$ $\pm$	6 14
ERM-DA193								

Availability: In units of lyophilised material of a 1.25 mL portion of serum kept under nitrogen in sealed glass ampoules.

<sup>1)</sup> The sample is to be reconstituted with (1.25  $\pm$  0.01) mL of distilled water.

		Progesterone in human serum (concentration in the reconstituted material <sup>1)</sup> )						
		( $\mu\text{g/L}$ )		( $\text{nmol/L}$ )				
BCR-348R		8.5	$\pm$	0.4	$\pm$	26.9	$\pm$	1.2
ERM-DA347		3.19	$\pm$	0.07	$\pm$	10.13	$\pm$	0.21

Availability: In units of lyophilised material of a 1 mL portion of serum kept under nitrogen in sealed glass ampoules.

<sup>1)</sup> The sample is to be reconstituted with (1.0  $\pm$  0.01) mL of distilled water.

		17 $\beta$ -Estradiol in human serum (concentration in the reconstituted material) Amount-of-substance concentration ( $\text{nmol/L}$ )			
BCR-576 <sup>1)</sup>			0.114	$\pm$	0.005
BCR-577 <sup>2)</sup>			0.689	$\pm$	0.032
BCR-578 <sup>2)</sup>			1.34	$\pm$	0.07

Availability: BCR-576, -577, -578 are lyophilised material of a 5 mL (BCR-576) or 1 mL (BCR-577 and BCR-578) portion of serum kept under nitrogen in sealed glass ampoules.

<sup>1)</sup> The sample is to be reconstituted with (5.00  $\pm$  0.05) mL of distilled water.

<sup>2)</sup> The sample is to be reconstituted with (1.00  $\pm$  0.01) mL of distilled water.

### 3.2.2 CERTIFIED FOR THE TOTAL ELEMENT CONTENT AND OTHER PROPERTIES

	Description	Substance	Metal concentrations in the reconstituted material <sup>1)</sup> ( $\mu\text{g/L}$ )		
ERM-CE195	Lyophilised bovine blood	Pb Cd <sup>2)</sup>	416 5.06	$\pm$ $\pm$	9 0.15
ERM-CE196	Lyophilised bovine blood	Pb Cd <sup>2)</sup>	772 12.33	$\pm$ $\pm$	11 0.20
BCR-634	Lyophilised human blood	Pb Cd	46 1.4	$\pm$ $\pm$	5 0.4
BCR-635	Lyophilised human blood	Pb Cd	210 6.6	$\pm$ $\pm$	24 0.6
BCR-636	Lyophilised human blood	Pb Cd	$0.52 \cdot 10^3$ 11.6	$\pm$ $\pm$	$0.05 \cdot 10^3$ 0.6

Availability: In units of lyophilised material equivalent to about 5.75 mL of bovine blood with additives kept under nitrogen in rubber stoppered vials.

BCR-634, BCR-635 and BCR-636 are available in lyophilised form in brown glass vials, containing approximately 0.6 g dry matter equivalent to 3.0 mL of fresh whole blood.

<sup>1)</sup> The sample is to be reconstituted with (5.00  $\pm$  0.01) mL water.

<sup>2)</sup> Recertified by IRMM.

	Description	Substance	Element concentration in the reconstituted material <sup>1)</sup> (mmol/L)		
BCR-304	Lyophilised human serum	Ca Li Mg	2.201 0.985 1.85	± ± ±	0.019 0.029 0.03

Availability: In units of lyophilised material equivalent to about 5.3 mL of human serum kept under vacuum in rubber stoppered vials.

<sup>1)</sup> The sample is to be reconstituted with (5.00 ± 0.01) mL bi-distilled water.

	Description	Substance	Metal concentrations (µg/L)		
BCR-637	Human serum	Al Se Zn	12.5 81 1110	± ± ±	3.0 7 220
BCR-638	Human serum	Al Se Zn	55 104 1430	± ± ±	7 7 210
BCR-639	Human serum	Al Se Zn	194 133 2360	± ± ±	14 12 140

Availability: supplied in frozen form in white plastic vials containing approximately 4.5 mL serum.

Substance	ERM-DB001 Human hair (mg/kg)		
As	0.044	±	0.006
Cd	0.125	±	0.007
Cu	33	±	4
Hg	0.365	±	0.028
Pb	2.14	±	0.20
Se	3.24	±	0.24
Zn	209	±	12

Availability: supplied in amber glass bottle, provided in aluminium sachet, and contains a minimum amount of 3.5 g of a human hair homogeneous powder

### 3.2.3 CERTIFIED FOR PROTEIN CONTENT

	Description	Mass concentration in the reconstituted material <sup>1)</sup> (g/L)		
BCR-393	Lyophilised Apo A I from human serum	1.06	±	0.05

Availability: In units of lyophilised material equivalent to about 1.5 mL of Apolipoprotein solution in sealed glass ampoules under nitrogen.

<sup>1)</sup> The sample must be reconstituted with 1.0 mL of phosphate buffer.

	Description	Mass concentration in the reconstituted material <sup>1)</sup> (g/L)		
BCR-457	Human Thyroglobulin (Tg)	0.324	±	0.018

Availability: In units of lyophilised material in sealed glass ampoules under nitrogen.

<sup>1)</sup> The sample must be reconstituted with 1.0 mL of distilled water.

	Description	Protein mass per ampoule <sup>1)</sup> (µg)		
BCR-486	Purified alphafoetoprotein (AFP)	100	±	9

Availability: BCR-486 is provided in sealed glass ampoules. Each sample is in lyophilised form and it contains purified AFP without additives. The protein mass per ampoule is equivalent to (100 ± 9) µg when the material is reconstituted with 1.0 mL phosphate buffer according to the specified procedure.

<sup>1)</sup> Carbohydrate mass of the molecule is not included.

	Description	Protein mass/ampoule
<b>BCR-613</b>	Prostate specific antigen in the reconstituted material	71 ± 7 µg

Availability: Lyophilised PSA in sealed glass ampoules kept under argon gas.

	Description	HbA <sub>1c</sub> /Hb <sub>T</sub> in reconstituted material (%)
<b>BCR-405 (RM)</b>	Glycated haemoglobin (HbA <sub>1c</sub> ) in human haemolysate	(6.29 ± 0.18)

Value in brackets is not certified.

Availability: Sealed glass ampoules of lyophilised material equivalent to about 0.5 mL of a solution of haemolysate of human erythrocytes kept under carbonmonoxide.

- <sup>1)</sup> Sample to be reconstituted with 1 mL of deionised water and diluted with appropriate haemolyzing reagent, taking into account that the total haemoglobin (Fe<sub>4</sub>) concentration is about 0.23 mmol/L (15 g/L).

ERM-DA470k Human Serum Proteins			
Description	Mass concentration <sup>1)</sup> (g/L)	Description	Mass concentration <sup>1)</sup> (g/L)
$\alpha_2$ macroglobulin (A2M)	1.43 ± 0.06	Haptoglobin (HPT)	0.889 ± 0.021
$\alpha_1$ acid glycoprotein (AAG)	0.617 ± 0.013	Immunoglobulin A (IgA)	1.80 ± 0.05
$\alpha_1$ antitrypsin (AAT)	1.12 ± 0.03	Immunoglobulin G (IgG)	9.17 ± 0.18
Albumin (ALB)	37.2 ± 1.2	Immunoglobulin M (IgM)	0.723 ± 0.027
Complement 3c (C3c)	1.00 ± 0.04	Transferrin (TRF)	2.36 ± 0.08
Complement 4 (C4)	0.162 ± 0.007	Transthyretin (TTR)	0.220 ± 0.018

Availability: Glass bottle containing lyophilised materials equivalent to about 1 mL of serum with additives kept under nitrogen.

- 1) Sample to be reconstituted with (1.00 ± 0.01) g water.

	Description	Mass concentration (mg/L)
<b>ERM-DA471/IFCC</b>	Cystatin C	5.48 ± 0.15

Availability: Glass vial containing lyophilised human serum spiked with cystatin C.

	Description	Mass concentration (mg/L)
<b>ERM-DA474/IFCC</b>	C-reactive protein (CRP)	41.2 ± 2.5

Availability: Glass ampoule containing at least 1 mL processed human serum spiked with CRP.

	Description	Absorbance at 540 nm and 10.00 mm pathlength	Mass concentration (mg/L)	Substance concentration (µmol/L)
<b>BCR-522</b>	Haemiglobincyanide (HiCN) in bovine blood lysate	0.5457 ± 0.0009	800.3 ± 1.3	49.61 ± 0.08

Availability: Bovine blood lysate in sealed brown glass ampoules (10 mL) equivalent to about 800.3 mg/L of haemiglobincyanide.

	Description	Amount-of-substance concentration of creatinine (µmol/L)
<b>BCR-573</b>	Creatinine in human serum	68.7 ± 1.4
<b>BCR-574</b>	Creatinine in human serum	105.0 ± 1.3
<b>BCR-575</b>	Creatinine in human serum	404.1 ± 7.1

Availability: BCR-573, -574, -575 are the lyophilised form of approximately 1 mL portion of serum, with no additives. The mass of the lyophilised material contained in the ampoule is about 0.09 g.

## **BCR-573i (RM) Set of creatinine interfering substances**

Availability: Consists of three vials with lyophilised solutions

- 0.025 mg calcium dobesilate / 1.2 mg cefoxitin;
- 0.044 mg sodium pyruvate;
- 0.108 mg bilirubin ditaurate.

	Description	Amount-of-substance fraction (mmol/mol)	
<b>IRMM/IFCC-467</b>	Haemoglobin isolated from whole blood	HbA0/(HbA1c + HbA0)	> 976

Availability: Provided in vials containing approximately 39 mg a deep frozen buffered solution.

### **3.2.4 CERTIFIED FOR CATALYTIC ACTIVITY**

	Description	Catalytic concentration in reconstituted material Certified value	
		U/L	µkat/L
<b>BCR-410</b>	Prostatic acid phosphatase highly purified, from human prostate <sup>2)</sup>	28.0 ± 0.7	0.466 ± 0.012
<b>BCR-647</b>	Human adenosine deaminase (ADA1), from human erythrocytes <sup>2)</sup>		2.55 ± 0.09
<b>BCR-693</b>	Human pancreatic lipase from pancreatic juice <sup>4)</sup>		28.9 ± 1.2
<b>BCR-694</b>	Human pancreatic lipase (recombinant) <sup>4)</sup>		17.4 ± 1.0
<b>ERM-AD452/IFCC</b>	γ-Glutamyltransferase partially purified, from pig kidney <sup>3)</sup>	114.1 ± 2.4	1.90 ± 0.04
<b>ERM-AD454/IFCC</b>	Alanine aminotransferase partially purified, from pig heart <sup>3)</sup>	186 ± 4	3.09 ± 0.07
<b>ERM-AD455/IFCC</b>	Creatine kinase CK-MB from human heart <sup>3)</sup>	101 ± 4	1.68 ± 0.07
<b>IRMM/IFCC-456</b>	Human pancreatic α-Amylase <sup>3)</sup>		9.1 ± 0.3
<b>ERM-AD457/IFCC</b>	Aspartate Transaminase (AST)	104.6 ± 2.7	1.74 ± 0.05

Availability: Sealed glass ampoules of lyophilised material equivalent to about 1 mL of a solution of enzyme stabilized by incorporation in serum albumin matrix of human (BCR-410) or bovine (ERM-AD452/IFCC, ERM-AD454/IFCC and ERM-AD457/IFCC) origin kept under dry nitrogen. BCR-647 has been stabilised by incorporation in a matrix of 50 mmol/L Tris/HCl buffer 9pH=7.4) and human serum albumin (30 g/L). ERM-AD455/IFCC and IRMM/IFCC-456 are provided in sealed ampoules or vials filled with dry nitrogen. Samples are in lyophilised form and equivalent to about 1 mL of a solution of purified enzyme.

BCR-693 and BCR-694 are provided in ampoules of lyophilised material equivalent to about 1 mL of stabilised enzyme.

<sup>1)</sup> According to IFCC recommended method at 30 °C.

<sup>2)</sup> According to method specified in report.

<sup>3)</sup> According to IFCC recommended method at 37 °C.

<sup>4)</sup> According to method described in certification report at 37 °C.

### 3.2.5 CERTIFIED FOR DNA SEQUENCE

#### CRMs for monitoring leukaemia (ERM-AD623)

SET OF PLASMID SOLUTIONS		
	Number of specific DNA fragments per plasmid	
	Certified value	Uncertainty
<i>BCR-ABL</i> b3a2 transcript <i>BCR</i> transcript <i>GUSB</i> transcript	1	negligible
	1	negligible
	1	negligible
Copy number concentration of the plasmid		
	Certified value [cp/µL]	Uncertainty [cp/µL]
ERM-AD623a	$1.08 \times 10^6$	$0.13 \times 10^6$
ERM-AD623b	$1.08 \times 10^5$	$0.11 \times 10^5$
ERM-AD623c	$1.03 \times 10^4$	$0.10 \times 10^4$
ERM-AD623d	$1.02 \times 10^3$	$0.09 \times 10^3$
ERM-AD623e	$1.04 \times 10^2$	$0.10 \times 10^2$
ERM-AD623f	10.0	1.5

Availability: ERM-AD623 is a set of six plasmid solutions (a-f). Each of six vials contains approximately 600 µL of plasmid solution.

### 3.2.6 OTHERS

	Description	Parameters of the calibration line
ERM-AD148	Lyophilised thromboplastin Bovine (OBT/79)	Slope      1.011    ±    0.015 intercept   - 0.321    ±    0.025
ERM-AD149	Lyophilised rabbit thromboplastin	Slope      1.257    ±    0.013 Intercept   - 0.242    ±    0.019

Availability: ERM-AD148 in units of lyophilised bovine brain thromboplastin equivalent to about 2.2 g bovine brain tissue extract kept under vacuum in sealed glass ampoules.

ERM-AD149 in sealed glass ampoules containing the lyophilised form of a 0.5 mL aliquot of the extract of rabbit brain tissue, without calcium ion added.

	BCR-665 Asbestos fibres in lung tissue (Number of fibres of more than 1 µm in length in million per g dry tissue)	BCR-666 Asbestos fibres in lung tissue (Number of fibres of more than 1 µm in length in million per g dry tissue)
Amosite + crocidolite Anthophyllite	49    ±    16 1.8    ±    0.9	2.3    ±    0.9 5.1    ±    1.5

Availability: Sealed vials with 100 mg of lung tissue.

	IRMM-435 Pharmaceutical glass containers Alkali leaching and release
Volume of titration solution 0.01 mol/L HCl per 50 mL of leachate Sodium release per volume of leachate Release of Na <sub>2</sub> O per volume of leachate	0.38    ±    0.04    mL 1.41    ±    0.14    mg/L 1.91    ±    0.19    mg/L

Availability: Each unit of IRMM-435 consists of 20 vials of 18.9 mL brimful capacity, made of a semi-durable type of glass, which screw caps.

## 4 MATERIALS CERTIFIED FOR PHYSICAL PROPERTIES

### 4.1 CERTIFIED FOR THERMAL PROPERTIES

#### Resin bonded glass fibre board IRMM-440

The certified thermal conductivity between – 10 °C and + 50 °C is given by

$$\lambda [\text{W}/(\text{m}\cdot\text{K})] = 0.029\,394\,9 + 0.000\,106\,0 \times T [^{\circ}\text{C}] + 2.047 \times 10^{-7} \times T^2 [({}^{\circ}\text{C})^2]$$

This equation is valid for a sample of the reference material within the density range [64 kg/m<sup>3</sup> - 78 kg/m<sup>3</sup>].

The uncertainty of the certified thermal conductivity is ± 0.000 28 W/(m.K) at the 95 % confidence level over the range [- 10 °C / + 50 °C].

Availability: Boards which can be cut to: (300 x 300 x 35) mm, (500 x 500 x 35) mm, (600 x 600 x 35) mm, (1000 x 1000 x 35) mm.

#### Glass-ceramic BCR-724

The certified thermal diffusivity between 298 K and 1025 K is given by

$$\alpha [\text{m}^2/\text{s} \cdot 10^{-6}] = 4.406 - 1.351 \cdot 10^{-2} \cdot T + 2.133 \cdot 10^{-5} \cdot T^2 - 1.541 \cdot 10^{-8} \cdot T^3 + 4.147 \cdot 10^{-12} \cdot T^4$$

The uncertainty of the certified thermal diffusivity is ± 6.1 [ % ] at the 95 % confidence level over the range from 298 K to 1025 K.

The certified thermal conductivity between 298 K and 1025 K is given by

$$\lambda [\text{W}/(\text{m} \cdot \text{K})] = 2.332 + 515.1 / T$$

The uncertainty of the certified thermal conductivity is ± 6.5 [ % ] at the 95 % confidence level over the range from 298 K to 1025 K.

Availability: Glass-ceramic cylinders in different shapes (BCR-724A: diameter = 13.0 mm, height > 18 mm; BCR-724B: diameter = 13.9 mm, height > 21 mm; BCR-724C: diameter = 25.9 mm, height > 22 mm; BCR-724D: diameter = 26.9 mm, height > 22 mm; BCR-724E: diameter = 50.7 mm, height > 25 mm).

	Description	Cold filter plugging point CFPP Temperature (°C)			Cloud point CP (°C)		
BCR-395	Gasoil	– 5.8      ±      0.4			(- 5 °C)		
ERM-FC395k	Gasoil	– 7.9      ±      1.6			– 7.2      ±      3.0		

Values in brackets are not certified.

Availability: BCR-395 is available in sealed jar which conforms to the dimension given in EN 116, ready for use in the apparatus without need to transfer the material.

ERM-FC395k consists of a set two amber glass ampoules, each containing 27 mL of gas oil.

Substance	ERM-EF411 Hard coal			ERM-EF412 Brown coal			ERM-EF413 Furnace coke		
Gross calorific value (GCV) (MJ/kg)	29.0	±	0.4	26.02	±	0.22	29.5	±	0.4
Net calorific value (NCV) (MJ/kg)	28.0	±	0.4	24.98	±	0.25	29.4	±	0.5
Volatile matter (g/100 g)	38.1	±	1.0	50.1	±	0.7			
Ash (g/100 g)	8.3	±	0.7	4.11	±	0.23			
C (g/100 g)	71.4	±	1.0	66.2	±	0.7	87.8	±	1.9
Ca (g/kg)				9.8	±	0.4	2.92	±	0.22
Cd (mg/kg)				(0.012	±	0.004)			
Cl (mg/kg)	99	±	19				(350	±	130)
Co (mg/kg)	(3.5	±	0.8)						
Cu (mg/kg)				(0.68	±	0.22)			
H (g/100 g)	4.80	±	0.14	4.88	±	0.15			
Hg (mg/kg)	(0.079	±	0.015)	0.071	±	0.011			
K (mg/kg)				229	±	18			
Mg (g/kg)				(3.73	±	0.16)			
Mn (mg/kg)				48.6	±	1.9			
N (g/100 g)	1.43	±	0.10	0.74	±	0.06	1.10	±	0.07
Na (g/kg)				2.20	±	0.12	0.64	±	0.07
Pb (mg/kg)				(0.25	±	0.05)			
S (g/100 g)	0.598	±	0.017	0.360	±	0.023	0.58	±	0.12
Sb (mg/kg)	(1.5	±	0.4)	(0.024	±	0.004)			
Se (mg/kg)	5.1	±	1.0	0.96	±	0.14	1.33	±	0.26
Tl (mg/kg)	(0.24	±	0.07)						
V (mg/kg)	(22	±	7)	0.57	±	0.04			
Zn (mg/kg)	(13	±	4)	(0.99	±	0.18)	16.0	±	2.5

Values in brackets are not certified.

Availability: ERM-EF411, ERM-EF412 and ERM-EF413 are available in units of about 50 g in aluminium-laminated sachets.

## 4.2 CERTIFIED FOR MECHANICAL PROPERTIES

### Shear testing of powders BCR-116

The flow of powders or granulated materials under the force of gravity affects the design and operation of silos used for their bulk storage. The European Federation of Chemical Engineering (EFCE) therefore developed a test method, based on the Jenike Shear Cell, to determine the shear strength of powders under different compaction and loading conditions. The complexity of this method is such that errors due to poor technique can easily arise. A reference material has therefore been produced with which laboratories can verify both their equipment and experimental technique.

Essentially the EFCE method consists of using a known load to compact a powder sample into a cylindrical Jenike Cell composed of two metal rings one upon the other. Having reached critical compaction of the powder, the steady state force necessary to displace the upper ring horizontally with respect to the lower one is determined with the compaction load still applied. Having established steady state shear the normal load on the powder is then reduced and the horizontal force necessary to continue to shear the powder is determined. It is this variation of the shear force as a function of the reduced normal load for a given compaction load which characterizes the powder.

The reference material consists of 3 kg of limestone powder packed in a polyethylene jar. It is accompanied by a certificate giving shear stress as a function of normal applied stress for four different powder compaction stresses.

## Creep BCR-425

Creep is the progressive deformation of a material under load. Metallic materials are usually tested at elevated temperatures for periods of 1 000 to 100 000 hours by surrounding them with a suitable constant temperature furnace. The need to operate under such conditions gives rise to problems of alignment, strain measurement, temperature measurement etc. which can result in considerable differences in results between laboratories.

BCR-425 was therefore developed to allow laboratories to validate their creep testing rigs and procedures as a whole within a reasonable time (500 to 600 hours) using a material whose properties are sensitive to test conditions.

The test piece is delivered in the form of a bar of 14 mm diameter and 500 mm length which must be machined by the laboratory to the required shape and size.

At a test temperature of 600 °C, using an applied stress of 160 MPa the certified properties are as follows:

Creep rate at 400 hours of	(72	±	5)	$10^{-6} \text{ h}^{-1}$
Time to a creep strain of 2 %	(278	±	16)	h
Time to a creep strain of 4 %	(557	±	30)	h

## Nimonic 75 for ambient temperature tensile properties BCR-661

Property	Certified value			
	(300	±	8)	MPa
0.2 % proof stress $R_{p0.2}$	(318	±	7)	MPa
0.5 % proof stress $R_{p0.5}$	(750	±	14)	MPa
Ultimate tensile strength $R_m$	(40.9	±	0.9)	%
Elongation to fracture A	(60	±	4)	%
Reduction in area Z				

Availability: BCR-661 can be supplied in 3 bars of 150 mm long and diameter 14 m or as 1 bar of 500 m long, sufficient for the manufacture of three test-pieces.

## Scratch testing BCR-692

Failure event	Critical load	
	Certified value (N)	Uncertainty (N)
Forward chevron cracks at the borders of the scratch track. <i>(Lc<sub>1</sub> shall be taken at the closest end of the event to the scratch track start).</i>	(Lc <sub>1</sub> ) 13.6	1.8
Forward chevron cracks at the borders of the scratch track, with local interfacial spallation or with gross interfacial spallation. <i>(Lc<sub>2</sub> shall be taken at the failure event that occurs first and at the closest end of the event to the scratch track start).</i>	(Lc <sub>2</sub> ) 17.0	2.1
Gross interfacial shell-shaped spallation. <i>(Lc<sub>3</sub> shall be taken at the first point where the substrate can be seen at the centre of the track in a crescent that goes completely through the track).</i>	(Lc <sub>3</sub> ) 28	2.9

Availability: The reference samples are (30x30x5) mm steel coupons coated with a diamond-like carbon coating (DLC) applied by plasma-assisted chemical vapour deposition. The coupons are distributed in a reusable plastic box containing desiccant.

## Impact toughness Charpy specimens

Impact toughness is the ability of a material to resist fracture under the effect of shock loading. It is determined by means of a conventional test defined in ISO and CEN Standards.

The test is sensitive to many sources of errors and therefore already 40 years ago, ASTM specification required testing machines to be verified periodically by reference specimens. The recent CEN standard (EN 10045-2: 1993) specifies that impact toughness machines should be certified by using the BCR certified reference Charpy specimens or other specimens traceable to the latter.

The reference specimens described here are supplied by groups of five. Their use must be done in accordance with the instructions given in the certification report, in particular with regard to degreasing. The fracture energy is certified for the CEN striker (knife of 2 mm radius).

Measurements generally need to be performed at 20°C, with the exception of ERM-FA013 where one batch has certified values at 0°C and 20°C to avoid jamming. We will deliver by default a batch certified by 20°C unless the order explicitly mentions the value at 0°C.

For each energy level there are several batches of slightly different values. The following table gives the nominal values, the certificate gives the actual values.

ERM-FA013 – Low Energy 20°C	20-30 J
ERM-FA013 – Low Energy 0°C	20-30 J
ERM-FA015	80 J
ERM-FA016	120 J
ERM-FA415	150J

## 4.3 CERTIFIED FOR MORPHOLOGICAL PROPERTIES

Particle size distributions					
	Form of Quartz	Certified Property	Size Range ( $\mu\text{m}$ )	Unit Size (g)	
BCR-066	Powder	Stokes' diameter	0.35 - 3.50	10	
BCR-067	Powder	Stokes' diameter	2.40 - 32.00	10	
BCR-068	Sand	Volume diameter	160.0 - 630.0	100	
BCR-069	Powder	Stokes' diameter	14.0 - 90.0	10	
BCR-070	Powder	Stokes' diameter	1.20 - 20.00	10	

Particle size distributions					
	Form of Quartz	Certified Property	Size Range ( $\mu\text{m}$ )	Unit Size (g)	
BCR-130	Powder	Volume diameter	50 - 220	50	
BCR-131	Powder	Volume diameter	480 - 1800	200	
BCR-132	Gravel	Volume diameter	1400 - 5000	700	

	Description	Specific Surface Area ( $\text{m}^2 \cdot \text{g}^{-1}$ )	Unit Size (g)
BCR-169	Alpha alumina	0.104 $\pm$ 0.012	60
BCR-170	Alpha alumina	1.05 $\pm$ 0.05	60
BCR-171	Alumina	2.95 $\pm$ 0.13	50
BCR-172	Quartz	2.56 $\pm$ 0.10	10
BCR-173	Titanium dioxide	8.23 $\pm$ 0.21	46
BCR-175	Tungsten	0.18 $\pm$ 0.04	200

## Mullite ( $3\text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2$ ) BCR-301 (RM)

High crystallinity.

Vitreous phase 0.03 g/g. No other phase detected.

Impurities in g/kg:

$\text{Fe}_2\text{O}_3$	< 2	$\text{Na}_2\text{O}$	< 1
$\text{CaO}$	< 1.2	$\text{K}_2\text{O}$	< 0.5
$\text{MgO}$	< 0.5	$\text{TiO}_2$	< 0.5

Mullite BCR-301 (RM)

Reflection	Lattice spacing (nm)	Relative intensity
[110]	0.538 2	0.50
[210]	0.339 0	1
[220]	0.269 5	0.40
[121]	0.220 6	0.59
[331]	0.152 4	0.36

## Microcrystalline cellulose BCR-302 (Water content at 10 water activities)

The water content of the material when in equilibrium with the atmosphere above each saturated salt solution specified in the table was determined at 25 °C by the method recommended by COST 90. BCR-302 is specifically intended to check the correct application of the COST procedure for determination of water sorption isotherms of foods.

Nominal water activity $a_w$ at 25 °C	Certified equilibrium water content mass fraction g/kg)	Specified saturated salt aqueous solution
0.1105	21.3 ± 1.1	1 Lithium Chloride
0.2245	32.4 ± 1.3	2 Potassium Acetate
0.3300	41.5 ± 0.9	3 Magnesium Chloride
0.4276	51.6 ± 0.9	4 Potassium Carbonate
0.5286	59.7 ± 1.4	5 Magnesium Nitrate
0.5770	64.7 ± 1.5	6 Sodium Bromide
0.7083	82.5 ± 1.7	7 Strontium Chloride
0.7528	88.9 ± 2.4	8 Sodium Chloride
0.8426	110 ± 4	9 Potassium Chloride
0.9019	133 ± 5	10 Barium Chloride

Availability: 20 g in sealed sachets.

	Description	Micropore volume (cm <sup>3</sup> /g <sup>1</sup> )	Median micropore width (nm)
BCR-704	Faujasite type zeolite	0.205 ± 0.006	0.668 ± 0.019
BCR-705	Linde type A zeolite	0.181 ± 0.006	0.592 ± 0.020

Availability: Glass bottle containing 10 g of pellets.

## Thickness of silicon dioxide on silicon BCR-564

With the trend towards the use of gate oxides with thicknesses around 10 nm the silicon oxide/silicon interface characteristics gain more and more importance. Reference materials of silicon dioxide layers of nominal thickness of 10, 50 and 120 nm on silicon wafers have been prepared to be used as calibration standards for ellipsometry. They provide the means for a quantitative analysis of oxides thicker than 10 nm. Each standard is individually certified.

Availability: Set of three standards (16 × 16 mm): 1 standard each of 10, 50 and 120 nm SiO<sub>2</sub> on silicon wafer.

## Reference material for depth profiling by ion beam sputtering BCR-261T

In order to achieve the accuracy required when measuring compositional depth profiles using ion beam sputtering in association with Auger Electron Spectroscopy, a reference material of accurately known thickness on a stable substrate is required. BCR-261T is a tantalum pentoxide on tantalum foil reference material existing in two nominal thicknesses of 30 and 100 nm.

	Nominal thickness (nm)	Certified values			
		$10^{21}$ oxygen atoms/m <sup>2</sup>		oxide thickness ratio	
BCR-261T	(30) (100)	1.72 5.40	$\pm$ $\pm$	0.07 0.12	0.321 0.013

Values in brackets are not certified.

Availability: Four rectangular foils of 5 x 10 mm of each oxide thickness.

Colloidal Silica in water ERM-FD100	Equivalent spherical diameter	
	Certified value (nm)	Uncertainty (nm)
Intensity-weighted harmonic mean diameter (DLS)	19.0	0.6
Intensity-based modal Stokes diameter (CLS)	20.1	1.3
Number-based modal diameter (TEM/SEM)	19.4	1.3
Intensity-weighted mean diameter (SAXS)	21.8	0.7
Equivalent spherical diametr, volume-weighted mean (SAXS)	(20.4)	(1.6)
Zeta Potential	(- 43.0 mV)	(22 mV)

Values in brackets are not certified.

Availability: ERM-FD100 is available in 10 mL pre-scored amber glass ampoules containing approximately 9 mL of suspension.

Colloidal Silica in aqueous solution ERM-FD304	Equivalent spherical diameter	
	Certified value (nm)	Uncertainty (nm)
Scattering intensity-weighted harmonic mean diameter (DLS)	42.1	0.6
Extinction intensity-based modal Stokes (CLS)	33.0	3.0
Number-based modal diameter (TEM/SEM)	(27.8)	(1.5)

Values in brackets are not certified.

Availability: ERM-FD304 is available in 10 mL pre-scored amber glass ampoules containing approximately 9 mL of suspension.

## 4.4 CERTIFIED FOR OPTICAL PROPERTIES

### Tomato paste colour reference tile BCR-400

BCR-400 consists of a ceramic tile the colour of which is defined by Hunter L, a and b values. If other data are needed these may be calculated readily for the same geometry or illuminant / standard observer (45°/0° or 0°/45° viewing geometry, illuminant C/2° standard observer combination, aperture of ≥ 50 mm nominal diameter. The nominal Hunter values of the tiles are: L = 26, a = 33 and b = 14.5. Each tile is individually certified, the overall uncertainties on L, a and b being ± 0.3, ± 0.4 and ± 0.3 respectively.

BCR-400 is intended for calibration purpose only and does not represent a standard tomato paste colour.

## 5 MATERIALS RELATED TO INDUSTRIAL APPLICATIONS

### 5.1 CERTIFIED FOR COMPOSITION

Certified Parameter	IRMM-441 n-Heptane (g/kg)	IRMM-442 Isooctane (g/kg)
Isooctane, purity by difference		999.85 ± 0.05
n-Heptane, purity by difference	999.85 ± 0.05	
Impurities		
Total organics (other than isoctane)		0.11 ± 0.04
Total organics (other than n-Heptane)	0.12 ± 0.05	
Isooctane	0.07 ± 0.02	
n-Heptane		0.02 ± 0.02
Water	0.03 ± 0.02	0.04 ± 0.02

Availability: IRMM-441 and -442 are supplied in ampoules of 100 mL.

	Description	Substance	Certified values (g/kg)
BCR-010	Tin ore	Sn	765.9 ± 1.2

Availability: This CRM is contained in brown glass bottles. The approximate quantity per unit is 225 g.

	Description	Substance	Certified values (g/kg)
BCR-033	Super-phosphate	P <sub>2</sub> O <sub>5</sub> SO <sub>4</sub> CaO SiO <sub>2</sub> F Al <sub>2</sub> O <sub>3</sub> Fe <sub>2</sub> O <sub>3</sub> MgO	193.4 ± 1.2 428.0 ± 4.1 314.8 ± 3.1 29.2 ± 1.2 16.5 ± 0.5 11.0 ± 0.8 4.0 ± 0.2 2.1 ± 0.2
BCR-113	Potassium Chloride	K Cl Na Ca Mg water soluble K	502.5 ± 1.1 478.0 ± 0.9 15.3 ± 0.2 1.03 ± 0.04 0.24 ± 0.01 501.3 ± 0.7
BCR-114	Potassium Sulphate	K SO <sub>4</sub> Cl Na Ca Mg water soluble K	418.0 ± 0.9 533 ± 2 18.5 ± 0.1 11.0 ± 0.1 9.4 ± 0.2 0.74 ± 0.01 417.6 ± 0.8
BCR-178	Calcium Ammonium Nitrate	NH <sub>4</sub> – N NO <sub>3</sub> - N total - N Ca	130.44 ± 0.32 130.15 ± 0.57 260.19 ± 0.54 88.82 ± 0.27
BCR-179	Urea	total – N Uric – n Biuret	465.4 ± 0.8 460.9 ± 0.9 10.37 ± 0.11

Availability: Units of about 100 g in the form of fine powder.

	Description	Substance	Certified values			
<b>BCR-032</b>	Moroccan Phosphate rock	CaO	518	±	4	g/kg
		P <sub>2</sub> O <sub>5</sub>	329.8	±	1.7	g/kg
		CO <sub>2</sub>	51.0	±	0.8	g/kg
		F	40.4	±	0.6	g/kg
		SiO <sub>2</sub>	20.9	±	1.2	g/kg
		SO <sub>3</sub>	18.4	±	0.8	g/kg
		Al <sub>2</sub> O <sub>3</sub>	5.5	±	0.6	g/kg
		MgO	4.0	±	0.1	g/kg
		Fe <sub>2</sub> O <sub>3</sub>	2.3	±	0.1	g/kg
		As	(9.5	±	0.5	mg/kg)
		B	(22.6	±	2.2	mg/kg)
		Cd	(20.8	±	0.7	mg/kg)
		Cr	(257	±	16	mg/kg)
		Co	(0.59	±	0.06	mg/kg)
		Cu	(33.7	±	1.4	mg/kg)
		Hg	(0.055	±	0.011	mg/kg)
		Mn	(18.8	±	1.3	mg/kg)
		Ni	(34.6	±	1.9	mg/kg)
		Ti	(171	±	10	mg/kg)
		V	(153	±	7	mg/kg)
		Zn	(253	±	6	mg/kg)

Values in brackets are not certified.

Availability: Units of about 100 g in the form of fine powder.

Substance	<b>BCR-126A</b> Lead crystal glass (cg/g)		
SiO <sub>2</sub>	57.80	±	0.11
PbO	23.98	±	0.06
K <sub>2</sub> O	9.99	±	0.07
Al <sub>2</sub> O <sub>3</sub>	0.126	±	0.013
Fe <sub>2</sub> O <sub>3</sub>	0.005 5	±	0.001 2
Sb <sub>2</sub> O <sub>3</sub>	0.291	±	0.012
BaO	1.053	±	0.030
CaO	1.033	±	0.030
MgO	0.512	±	0.013
ZnO	1.01	±	0.04
Na <sub>2</sub> O	3.57	±	0.07
Li <sub>2</sub> O	0.494	±	0.016
Density at 20 °C	2.990 5	±	0.001 6 g/cm <sup>3</sup>
Refractive index n <sub>D</sub> <sup>20 °C</sup> at 589.3 nm	1.559 67	±	0.000 22

Availability: In the form of square plates (100 × 100 mm) and 10 mm thickness.

## 5.2 CERTIFIED FOR TRACE ELEMENT CONTENT

	Material	Certified value	Form	Unit
IRMM-521	Ni	< 0.1 mg Co kg <sup>-1</sup>	B: 0.5 mm wire R: 0.1 mm foil	100 cm <sup>2</sup> (1.8 g) 75 cm <sup>2</sup> (6.7 g)
IRMM-522	Cu	< 0.05 mg Co kg <sup>-1</sup> 0.95 ± 0.04 mg Ag kg <sup>-1</sup>	A: 0.1 mm foil B: 1.0 mm foil C: 0.5 mm wire D: 1.0 mm wire	100 cm <sup>2</sup> (8.9 g) 20 cm <sup>2</sup> (17.8 g) 1 m (1.8 g) 1 m (7.0 g)
IRMM-523	Al	< 0.1 mg Na kg <sup>-1</sup>	A: 0.1 mm foil B: 1.0 mm foil C: 1.0 mm wire	100 cm <sup>2</sup> (2.7 g) 20 cm <sup>2</sup> (5.4 g) 1 m (2.1 g)
IRMM-524	Fe	< 0.05 mg Co kg <sup>-1</sup> < 0.1 mg Mn kg <sup>-1</sup>	A: 0.1 mm foil B: 0.5 mm wire	100 cm <sup>2</sup> (7.9 g) 1 m (1.6 g)
IRMM-525	Nb	19.6 ± 1.8 mg Ta kg <sup>-1</sup>	A: 0.02 mm foil B: 0.1 mm foil C: 0.5 mm wire	20 cm <sup>2</sup> (0.3 g) 20 cm <sup>2</sup> (1.7 g) 1 m (1.7 g)
IRMM-526	Nb	0.30 ± 0.09 mg Ta kg <sup>-1</sup>	A: 0.02 mm foil B: 0.1 mm foil C: 0.5 mm wire	20 cm <sup>2</sup> (0.3 g) 20 cm <sup>2</sup> (1.7 g) 1 m (1.7 g)
IRMM-529	Rh	< 5 g Pt kg <sup>-1</sup> 26.0 ± 0.6 g Ir kg <sup>-1</sup>	0.05 mm foil	20 cm <sup>2</sup> (1.2 g)
IRMM-531	Ti	< 0.1 mg Sc kg <sup>-1</sup>	A: 0.1 mm foil B: 0.5 mm foil C: 0.5 mm wire	100 cm <sup>2</sup> (4.5 g) 20 cm <sup>2</sup> (4.5 g) 1 m (1 g)
IRMM-527R	Al – 0.1% Co	1.001 ± 0.024 g Co kg <sup>-1</sup>	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm <sup>2</sup> (2.7 g) 1 m (0.5 g) 1 m (2.1 g)
IRMM-528R	Al – 1.0% Co	10.02 ± 0.23 g Co kg <sup>-1</sup>	A: 0.1 mm foil C: 1.0 mm wire	100 cm <sup>2</sup> (2.7 g) 1 m (2.1 g)
IRMM-530R	Al – 0.1% Au	1.003 ± 0.012 g Au kg <sup>-1</sup>	A: 0.1 mm foil C: 1.0 mm wire	100 cm <sup>2</sup> (2.7 g) 1 m (2.1 g)
IRMM-532	Al – 0.01% Co	0.100 0 ± 0.002 5 g Co kg <sup>-1</sup>	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm <sup>2</sup> (2.7 g) 1 m (0.5 g) 1 m (2.1 g)
IRMM-533	Al – 0.1% Ag	0.996 ± 0.017 g Ag kg <sup>-1</sup>	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm <sup>2</sup> (2.7 g) 1 m (0.5 g) 1 m (2.1 g)
IRMM-534	Al – 2.0% Sc	19.95 ± 0.20 g Sc kg <sup>-1</sup>	A: 0.1 mm foil B: 0.5 mm wire C: 1.0 mm wire	100 cm <sup>2</sup> (2.7 g) 1 m (0.5 g) 1 m (2.1 g)

	Description	Substance	Certified value (mg/kg)	Form, dimensions <sup>1)</sup> and availability
BCR-017A	Copper	P	6.85 ± 0.29	
BCR-017B		S	10.4 ± 0.6	
BCR-022A	Copper (electrolytic tough pitch)	O	138 ± 7	
BCR-022B	Copper (electrolytic tough pitch)	O	138 ± 7	Ø 9 mm, h 50 mm
BCR-054R	Copper (low oxygen)	O	0.47 ± 0.07	Ø 7 mm, h 50 mm
BCR-058	Copper (continuous cast)	O	390 ± 12	Ø 7 mm, h 50 mm
BCR-099	Nickel	O	8.4 ± 1.3	25 cubes of 2 g
		N	1.1 ± 0.3	
BCR-024B	Titanium	N	117 ± 13	B: 25 cubes of 0.4 g
BCR-024C		O	608 ± 23	C: 25 cubes of 0.2 g
BCR-059A	Titanium alloy Ti6Al4V	O	1750 ± 70	A: Ø 26 mm, h 9 mm
BCR-059B		N	172 ± 27	B: 25 cubes of 0.2 g
BCR-318	Titanium	H	12.2 ± 0.6	Ø 7 mm, h 1 mm (bottle with approx. 100 discs)
BCR-275	Zirconium alloy Zircaloy-4	O	1670 ± 50	Ø 13 mm, h 1 mm
		N	39.0 ± 1.7	(bottle with 10 discs)
		C	113 ± 4	
BCR-276	Zirconium alloy Zircaloy-4	O	1540 ± 80	Ø 4.5 mm, h 2 mm
		N	41 ± 9	(bottle with approx. 100 discs)
		C	108 ± 101	
BCR-102	Tungsten carbide powder	O	185 ± 4	Bottles containing 2 - 3 g powder, sealed under argon in an aluminium container

1) Ø = diameter, h = height

Substance	BCR-286		BCR-287		BCR-288	
	Electrolytically refined lead (mg/kg)		Thermally refined lead (mg/kg)		Lead with added impurities (mg/kg)	
Ag			15.20	± 0.21		
As			67.3	± 1.1	55.7	± 1.6
Bi	21.5	± 0.5	0.356	± 0.024	215.8	± 2.4
Cd			0.98	± 0.05	33.3	± 0.9
Cu			0.040	± 0.015	19.3	± 0.4
Sb	0.099	± 0.021			< 0.2	
Se					32.8	± 1.3
Te					2.26	± 0.08
Tl	2.47	± 0.07	0.73	± 0.04	8.2	± 0.4
Zn	< 0.1		< 0.1			

Availability: CRMs are available as follows: BCR-286A and -287A: blocks of 60 × 60 × 12 mm, BCR-286B, -287B and -288B: chips in bottles containing about 160 g.

Substance	BCR-321 Unalloyed zinc (mg/kg)	ERM-EB322 Unalloyed zinc (mg/kg)	ERM-EB323 Unalloyed zinc (mg/kg)	ERM-EB324 Unalloyed zinc (mg/kg)	ERM-EB325 Unalloyed zinc (mg/kg)	BCR-326 Unalloyed zinc (mg/kg)	BCR-327 Unalloyed zinc (mg/kg)
Al	< 0.7						
Cd	(0.23 ± 0.03)	15.08 ± 0.30	6.51 ± 0.21	48.6 ± 1.1	94.7 ± 2.5	203.0 ± 2.0	301.4 ± 2.3
Cu	(0.97 ± 0.05)	5.89 ± 0.15	18.9 ± 0.4	9.87 ± 0.18	47.5 ± 2.0	104.8 ± 2.7	(0.56 ± 0.11)
Fe	(2.22 ± 0.14)	19.1 ± 0.8	11.3 ± 0.7	58.5 ± 1.6	56.1 ± 3.3	264.8 ± 2.1	144.0 ± 1.3
In	< 0.2						
Pb	4.85 ± 0.20	15.0 ± 0.5	48.6 ± 0.9	26.1 ± 0.5	142 ± 9	307.0 ± 1.6	409.4 ± 2.3
Sn	< 0.5	5.6 ± 0.6	18.7 ± 0.7	9.8 ± 0.5	46.1 ± 2.0		
Tl	0.78 ± 0.10	5.28 ± 0.30	10.8 ± 0.5	19.9 ± 0.5	36.8 ± 1.2		

Values in brackets are not certified.

Availability: Discs of 80 mm diameter and 20 mm thickness (BCR-321, -326, -327) and 60 mm diameter, 30 mm thickness, respectively (ERM-EB322, EB323, EB324 and EB325).

Substance	BCR-351 ZnAl4 (mg/kg)	BCR-352 ZnAl4 (mg/kg)	BCR-353 ZnAl4 (mg/kg)	BCR-354 ZnAl4 (mg/kg)	BCR-355 ZnAl4 (mg/kg)
Al	[43.55 ± 0.11] × 10 <sup>3</sup>	[41.50 ± 0.10] × 10 <sup>3</sup>	[39.5 ± 0.4] × 10 <sup>3</sup>	[37.27 ± 0.16] × 10 <sup>3</sup>	[34.43 ± 0.13] × 10 <sup>3</sup>
Cd	(0.21 ± 0.03)	2.88 ± 0.12	10.44 ± 0.16	29.7 ± 0.4	58.1 ± 0.4
Cu	12.13 ± 0.15	31.26 ± 0.29	100.0 ± 0.8	312.3 ± 2.5	1035 ± 6
In	< 0.2	3.02 ± 0.28	2.55 ± 0.23	9.8 ± 0.9	24.6 ± 1.4
Mg	131.0 ± 0.9	283.0 ± 1.8	452.5 ± 2.4	602 ± 5	786 ± 6
Ni	(1.9 ± 0.6)	6.74 ± 0.16		83.1 ± 2.9	268 ± 8
Pb	4.50 ± 0.20	(6.4 ± 1.6)	24.4 ± 1.3	30.8 ± 1.2	56.9 ± 1.9
Sn	< 1	3.0 ± 0.7	5.6 ± 0.6	14.1 ± 1.1	29.1 ± 2.0
Tl	0.74 ± 0.06	3.2 ± 0.4	3.95 ± 0.22	11.01 ± 0.20	23.25 ± 0.28

Values in brackets ( ) are not certified.

Availability: Discs of 80 mm diameter and 20 mm thickness.

	BCR-356 ZnAl4Cu1 (mg/kg)	BCR-357 ZnAl4Cu1 (mg/kg)	BCR-359 ZnAl4Cu1 (mg/kg)	BCR-360 ZnAl4Cu1 (mg/kg)	BCR-361 ZnAl4Cu1 (mg/kg)
Al	[44.34 ± 0.11] × 10 <sup>3</sup>	[42.27 ± 0.11] × 10 <sup>3</sup>	[37.11 ± 0.11] × 10 <sup>3</sup>	[34.27 ± 0.12] × 10 <sup>3</sup>	[40.68 ± 0.19] × 10 <sup>3</sup>
Cd	0.73 ± 0.09	2.83 ± 0.10	29.8 ± 0.4	59.5 ± 0.6	(0.80 ± 0.17)
Cu	[3.944 ± 0.022] × 10 <sup>3</sup>	[5.849 ± 0.021] × 10 <sup>3</sup>	[9.89 ± 0.04] × 10 <sup>3</sup>	[12.34 ± 0.05] × 10 <sup>3</sup>	[7.98 ± 0.04] × 10 <sup>3</sup>
Fe	31.5 ± 0.6	25.7 ± 1.2	119.7 ± 1.1		10.34 ± 0.26
In	< 0.2	3.30 ± 0.14	15.5 ± 0.6	29.8 ± 0.6	(< 0.2)
Mg	132.3 ± 1.8	273 ± 4	557 ± 5	705 ± 5	
Ni	3.43 ± 0.19	9.82 ± 0.25	92.6 ± 0.6	267 ± 8	
Pb	9.87 ± 0.23	13.8 ± 0.6	36.2 ± 0.8	73.9 ± 1.4	5.31 ± 0.20
Sn	(0.32 ± 0.16)	3.51 ± 0.14	16.93 ± 0.22	33.0 ± 0.8	46.3 ± 0.9
Tl	0.79 ± 0.05	2.76 ± 0.05	13.34 ± 0.24	25.9 ± 0.7	37.4 ± 0.5

Values in brackets ( ) are not certified.

Availability: Discs of 80 mm diameter and 20 mm thickness.

Substance	BCR-089 TiAl6V4 (g/kg)		BCR-090 Titanium with added impurities (g/kg)		
Al	59.7	± 0.4			
B			0.0282	±	0.0014
Co			0.501	±	0.014
Cr			0.533	±	0.011
Cu			0.513	±	0.009
Fe			0.563	±	0.016
Mn			0.314	±	0.010
Mo			0.488	±	0.011
Nb			(0.492	±	0.026)
Ni			0.667	±	0.007
Sn			(0.71	±	0.05)
V	39.76	± 0.29			
W			(0.50	±	0.04)
Zr			(0.436	±	0.013)

Values in brackets are not certified.

Availability: BCR-089: Cylinder of 40 mm Ø and 20 mm height. BCR-090A: Cylinder of 40 mm Ø and 20 mm height. BCR-090B: Cubes of about 0.2 g in bottles containing approximately 25 g.

Substance	BCR-098 Zircaloy-4				
Cr	906	µg/g	±	9	µg/g
Fe	2143	mg/g	±	0.020	mg/g
Hf	77.6	µg/g	±	3.0	µg/g
Sn	14.60	mg/g	±	0.09	mg/g

Availability: Bottles containing about 10 g of chips.

Substance	BCR-074 Electrolytic copper (OFHC) (mg/kg)		
Ag	12.8	±	0.7
As	0.78	±	0.14
Bi	(0.10	±	0.03)
Cd		< 0.02	
Co		< 0.05	
Cr		< 0.1	
Fe	1.14	±	0.06
Mn	1.27	±	0.05
Ni	1.04	±	0.11
Pb	0.97	±	0.05
Sb	0.576	±	0.030
Se	0.37	±	0.04
Sn		< 0.07	
Te	(0.21	±	0.08)
Zn	0.46	±	0.07

Values in brackets are not certified.

Availability: CRMs are available as follows:

BCR-074A: Cylinder of 40 mm Ø, 30 mm height

	Substance	Quaternary bronze (g/kg)	Brass (g/kg)	Arsenic-Copper (g/kg)	Lead-bronze (g/kg)	Tin-bronze (g/kg)
<b>BCR-691</b>	As	1.94 ± 0.10	0.99 ± 0.10	46.0 ± 2.7	2.85 ± 0.22	1.94 ± 0.20
	Pb	79 ± 7	3.9 ± 0.4	1.75 ± 0.14	92 ± 17	2.04 ± 0.18
	Sn	71.6 ± 2.1	20.6 ± 0.7	2.02 ± 0.29	101 ± 8	70 ± 6
	Zn	60.2 ± 2.2	148 ± 5	0.55 ± 0.05	1.48 ± 0.24	1.57 ± 0.25

Availability: Set of five discs (one of each composition) of 35 mm Ø and 2 mm thickness, packed in a box.

	Description	Certified S content (g/kg)		
BCR-331	Steam Coal	4.99	±	0.10
BCR-332	High Volatile Industrial Coal	9.61	±	0.17
BCR-333	Coking Steam Coal	13.44	±	0.26
BCR-334	Anthracite	16.09	±	0.19
BCR-335	Flame Coal	50.8	±	0.6
BCR-336	High Volatile Steam Coal	32.90	±	0.26

Availability: These CRMs are available in units of about 20 g in ampoules with argon atmosphere.

Substance	BCR-460		
	Total Fluorine in coal powder (mg/kg)		
Cl	(59	±	18)
F	225	±	6

Values in brackets are not certified.

Availability: BCR-460 in glass bottles containing about 40 g.

Substance	BCR-461		
	Fluorine in clay (mg/kg)		
F	568	±	60

Availability: The samples are provided in units of 30 g in glass bottles.

	Description	S content (g/kg)		
ERM-EF672	Gasoil	0.203	±	0.006
ERM-EF671	Gasoil	0.452	±	0.009
ERM-EF104	Gasoil	1.019	±	0.019
BCR-105	Gasoil	3.63	±	0.10
BCR-106	Gasoil	5.02	±	0.08
BCR-107	Gasoil	10.40	±	0.15
ERM-EF211	Petrol	0.048 8	±	0.001 7

Availability: The materials are available in dark glass ampoules sealed under nitrogen. ERM-EF104, -671 and -672 contain 8 mL, BCR-105, -106 and -107 contain 25 g. ERM-EF211 is available in clear borosilicate glass ampoules and contains 19 mL.

	Description	Solvent Yellow 124 (SY124) content (mg/kg)		
ERM-EF317	Gasoil	0.141	±	0.018
ERM-EF318	Gasoil	7.0	±	0.4

Availability: The materials are available in dark glass ampoules sealed under nitrogen, containing 20 mL.

Certified Parameter	IRMM-441 n-Heptane (%)	IRMM-442 Isooctane (%)
n-Heptane, purity by difference	99.985	99.985
Isooctane, purity by difference	± 0.005	± 0.005
<u>Impurities</u>		
Total organics (other than isoctane)		0.011 ± 0.004
Total organics (other than n-Heptane)	0.012 ± 0.005	
Isooctane	0.007 ± 0.002	
n-Heptane		0.002 ± 0.002
Water	0.003 ± 0.002	0.004 ± 0.002
Lead	< 0.5 µg/L	< 1 µg/L

Availability: IRMM-441 and -442 are supplied in ampoules of 100 mL.

	Description	Substance	Certified values (g/kg)		
<b>BCR-109</b>	Zinc ore (blende)	Pb	7.38	±	0.03
		Fe	145.1	±	0.6
		Cu	9.46	±	0.08
		Cd	4.61	±	0.09
		Mg	0.20	±	0.01
		F	0.081	±	0.004
		Hg	0.0096	±	0.00012
<b>BCR-110</b>	Zinc ore (blende)	Pb	97.8	±	0.4
		Fe	5.46	±	0.10
		Cu	16.28	±	0.12
		Cd	10.51	±	0.07
		Mg	1.36	±	0.04
		F	0.055	±	0.003
		Hg	0.1484	±	0.0025

Availability: These RMs are contained in brown glass bottles. The approximate quantity per unit is 200 g for BCR-109 and 75 g for BCR-110.

	Description	Substance	Certified values (g/kg)		
<b>BCR-032</b>	Moroccan Phosphate rock	CaO	517.6	±	3.2
		P <sub>2</sub> O <sub>5</sub>	329.8	±	1.7
		CO <sub>2</sub>	51.0	±	0.8
		F	40.4	±	0.6
		SiO <sub>2</sub>	20.9	±	1.2
		SO <sub>3</sub>	18.4	±	0.8
		Al <sub>2</sub> O <sub>3</sub>	5.5	±	0.6
		MgO	4.0	±	0.1
		Fe <sub>2</sub> O <sub>3</sub>	2.3	±	0.1
		As	9.5 × 10 <sup>-3</sup>	±	0.5 × 10 <sup>-3</sup>
		B	22.6 × 10 <sup>-3</sup>	±	2.2 × 10 <sup>-3</sup>
		Cd	20.8 × 10 <sup>-3</sup>	±	0.7 × 10 <sup>-3</sup>
		Cr	257 × 10 <sup>-3</sup>	±	16 × 10 <sup>-3</sup>
		Co	0.59 × 10 <sup>-3</sup>	±	0.06 × 10 <sup>-3</sup>
		Cu	33.7 × 10 <sup>-3</sup>	±	1.4 × 10 <sup>-3</sup>
		Hg	55 × 10 <sup>-6</sup>	±	11 × 10 <sup>-6</sup>
		Mn	18.8 × 10 <sup>-3</sup>	±	1.3 × 10 <sup>-3</sup>
		Ni	34.6 × 10 <sup>-3</sup>	±	1.9 × 10 <sup>-3</sup>
		Ti	171 × 10 <sup>-3</sup>	±	10 × 10 <sup>-3</sup>
		V	153 × 10 <sup>-3</sup>	±	7 × 10 <sup>-3</sup>
		Zn	253 × 10 <sup>-3</sup>	±	6 × 10 <sup>-3</sup>

Availability: Units of about 100 g in the form of fine powder.

Substance	<b>BCR-664</b> Glass (mg/kg)		
As	5.9	±	0.4
Ba	29.1	±	0.7
Cd	5.7	±	0.4
Cl	68	±	8
Co	2.77	±	0.21
Cr	2.65	±	0.13
Pb	53.1	±	2.6
Sb	24.3	±	1.0
Se	8.6	±	0.5

Availability: Glass plate of (50 x 50 x 7) mm.

	Material	Uranium mass fraction (mg/kg)		
<b>IRMM-540R</b>	Uranium-doped oxide glass	15.0	±	0.9
<b>IRMM-541</b>	Uranium-doped oxide glass	49.4	±	2.7

Availability: Glass disc of 15 mm diameter and 2 mm thickness, polished on both sides.

Substance	ERM-EC590 Polyethylene (LDPE) g/kg			ERM-EC591 Polypropylene (PP) g/kg		
Br	2.13	±	0.09	2.08	±	0.07
2,4,4'-TriBDE (BDE-28)				0.0025	±	0.0004
2,2',4,4'-TetraBDE (BDE-47)	0.23	±	0.04	0.245	±	0.023
2,2',3,4,4'-PentaBDE (BDE-99)	0.302	±	0.030	0.32	±	0.04
2,2',4,4',6-PentaBDE (BDE-100)	0.063	±	0.005	0.066	±	0.007
2,2',4,4',5,5'-HexaBDE (BDE-153)	0.047	±	0.006	0.044	±	0.006
2,2',4,4',5,6'-HexaBDE (BDE-154)	0.0257	±	0.0026	0.026	±	0.004
2,2',3,4,4',5,6'-HeptaBDE (BDE-183)	0.132	±	0.012	0.087	±	0.008
2,2',3,3',4,4',6-OctaBDE + 2,2',3,4,4',5,6,6'-OctaBDE (BDE-197+204)	0.076	±	0.010	0.052	±	0.009
DecaBDE (BDE-209)	0.65	±	0.10	0.78	±	0.09
DecaBB (BB-209)	0.63	±	0.10	0.74	±	0.08
Sb	(0.756	±	0.025)	(0.713	±	0.022)

Values in brackets are not certified.

Availability: Brown glass bottle with 20 g granulate.

Substance	ERM-EC680k Polyethylene (low level) mg/kg			ERM-EC681k Polyethylene (high level)		
As	4.1	±	0.5	29.1	±	1.8 mg/kg
Br	96	±	4	0.77	±	0.04 g/kg
Cd	19.6	±	1.4	137	±	4 mg/kg
Cl	102.2	±	3.0	0.80	±	0.05 g/kg
Cr	20.2	±	1.1	100	±	5 mg/kg
Hg	4.64	±	0.20	23.7	±	0.8 mg/kg
Pb	13.6	±	0.5	98	±	6 mg/kg
S	76	±	4	0.63	±	0.04 g/kg
Sb	10.1	±	1.6	99	±	6 mg/kg
Sn	(15.3	±	2.8)	(86	±	6 mg/kg)
Zn	(137	±	20)	(1.25	±	0.07 g/kg)

Values in brackets are not certified.

Availability: Brown glass bottle with 100 g granulate.

### Cd in polyethylene (VDA 001-004)

A set of four certified reference materials for Cd in polyethylene (40.9 mg/kg, 75.9 mg/kg, 197.9 mg/kg and 407 mg/kg) has been certified by IRMM on behalf of VDA (Verband der Automobilindustrie e.V., Frankfurt). Information can be obtained from IRMM, Geel (B).

## 5.3 OTHERS

### Antimony implanted in silicon ERM-EG001 (BAM-L001)

Certified quantity	Certified value ( $10^{16} \cdot \text{cm}^{-2}$ )
Areal density of Sb atoms	4.81 ± 0.06
Isotope amount ratio $n(^{121}\text{Sb}) / n(^{123}\text{Sb})$	1.435 ± 0.006

Availability: The sample is a 10 mm x 10 mm silicon chip with a thermally grown surface oxide layer and Sb ions implanted with an energy of 400 keV. The certified value for the areal density of Sb atoms is valid for fractions of the chip surface down to  $0.15 \text{ mm}^2$  in size.

### Cementite Grains in Carburised Pure Iron (IRMM-471)

	Certified value (g/kg)
Carbon mass fraction in cementite grains	66.9 ± 2.7

Availability: 4-5 mm long rod with 5 mm diameter.

## 6 MATERIALS RELATED TO ISOTOPIC MEASUREMENTS

### 6.1 CERTIFIED FOR ISOTOPE ABUNDANCE RATIO (AMOUNT RATIO)

BCR-123 Ethanol						
Parameter	Ethanol H		Ethanol M		Ethanol L	
(D/H) <sub>I</sub>	$109.65 \times 10^{-6}$	$\pm 0.20 \times 10^{-6}$	$101.69 \times 10^{-6}$	$\pm 0.17 \times 10^{-6}$	$90.30 \times 10^{-6}$	$\pm 0.18 \times 10^{-6}$
(D/H) <sub>II</sub>	$119.76 \times 10^{-6}$	$\pm 0.25 \times 10^{-6}$	$130.94 \times 10^{-6}$	$\pm 0.21 \times 10^{-6}$	$122.20 \times 10^{-6}$	$\pm 0.4 \times 10^{-6}$
R	2.184	$\pm 0.005$	2.575	$\pm 0.006$	2.708	$\pm 0.009$

Availability: Units of 3 sealed NMR tubes containing respectively H-, M-, and L-ethanols, to which the tetramethylurea internal standard and the C<sub>6</sub>F<sub>6</sub> lock substance are added. 10 mm (BCR-123A) or 15 mm (BCR-123B) O.D. NRM tubes can be supplied.

Parameter	Unit	BCR-656 (96% ethanol)	BCR-657 (Sugar)	BCR-658 (Synthetic wine)	BCR-659 (Synthetic wine)	BCR-660 (Ethanol in water)
(D/H) <sub>I</sub> by <sup>2</sup> H-NMR	ppm	102.84 $\pm$ 0.20				102.90 $\pm$ 0.16
(D/H) <sub>II</sub> by <sup>2</sup> H-NMR	ppm	132.07 $\pm$ 0.30				131.95 $\pm$ 0.23
R by <sup>2</sup> H-NMR		2.570 $\pm$ 0.005				2.567 $\pm$ 0.005
$\delta^{13}\text{C}_{\text{VPDB}}$ by IRMS	$\text{\textperthousand}$	-26.91 $\pm$ 0.07	-10.76 $\pm$ 0.04			-26.72 $\pm$ 0.09
$\delta^{18}\text{O}_{\text{VSMOW}}$ of water from wine by IRMS	$\text{\textperthousand}$			-7.19 $\pm$ 0.04	-7.18 $\pm$ 0.02	
(D/H) <sub>w</sub> of water (IRMS)	ppm					148.68 $\pm$ 0.14
Alcoholic grade t <sub>D</sub>	w/w %	94.61 $\pm$ 0.05				11.96 $\pm$ 0.06 <sup>1)</sup>

1) in v/v %

Availability: BCR-656: Units of 25 mL of 96 % vol. neutral ethanol from wine in glass bottle;  
BCR-657: Units of approx. 1 g of dry glucose in a sealed amber vial;  
BCR-658: Units of 25 mL of synthetic wine solution in glass bottle;  
BCR-659: Units of 25 mL of synthetic wine solution in glass bottle;  
BCR-660: Units of 450 mL of aqueous ethanol solution in glass bottle.

Code	Description	Isotope amount content	Amount ratios					Unit size
IRMM-  007/1  007/2  007/3  007/4  007/5  007/6	0.5 M HNO <sub>3</sub>	$n(^{68}\text{Zn})/n(^{67}\text{Zn})$	$n(^{66}\text{Zn})/n(^{64}\text{Zn})$	$n(^{67}\text{Zn})/n(^{64}\text{Zn})$	$n(^{68}\text{Zn})/n(^{64}\text{Zn})$	$n(^{70}\text{Zn})/n(^{64}\text{Zn})$		5 mL
		$148.261(49) \cdot 10^{-9} \text{ mol} (^{64}\text{Zn})\cdot\text{g}^{-1}$	1.070 00 (47)	0.004 679 7(58)	0.021 337 4 (98)	0.022 830 9 (89)	0.000 067 57 (32)	
		$142.842(47) \cdot 10^{-9} \text{ mol} (^{64}\text{Zn})\cdot\text{g}^{-1}$	1.033 83 (45)	0.005 275 8 (57)	0.043 039 (20)	0.044 495 (18)	0.000 083 00 (31)	
		$125.44(25) \cdot 10^{-9} \text{ mol} (^{64}\text{Zn})\cdot\text{g}^{-1}$	1.012 45 (45)	0.007 057 4 (57)	0.107 896 (50)	0.109 239 (45)	0.000 129 11 (32)	
		$107.096(40) \cdot 10^{-9} \text{ mol} (^{64}\text{Zn})\cdot\text{g}^{-1}$	1.005 44 (45)	0.009 954 0 (60)	0.213 339 (99)	0.214 499 (88)	0.000 204 07 (44)	
		$79.518(32) \cdot 10^{-9} \text{ mol} (^{64}\text{Zn})\cdot\text{g}^{-1}$	1.001 62 (45)	0.016 608 8 (80)	0.455 60 (21)	0.456 34 (19)	0.000 376 29 (90)	
		$45.821(19) \cdot 10^{-9} \text{ mol} (^{64}\text{Zn})\cdot\text{g}^{-1}$	0.999 61 (45)	0.035 323 (17)	1.136 83 (52)	1.136 39 (46)	0.000 860 6 (23)	

Code	Description	Amount ratios	Unit size
		$n(^{41}\text{Ca})/n(^{40}\text{Ca})$	
ERM-AE701/1	0.6 M HNO <sub>3</sub> solution	1.011 4(68) · 10 <sup>-6</sup>	25 mL
ERM-AE701/2		1.023 5(69) · 10 <sup>-7</sup>	
ERM-AE701/3		1.018 1(69) · 10 <sup>-8</sup>	
ERM-AE701/4		1.047 9(71) · 10 <sup>-9</sup>	
ERM-AE701/5		1.052 0(71) · 10 <sup>-10</sup>	
ERM-AE701/6		1.091 3(74) · 10 <sup>-11</sup>	
ERM-AE701/7		1.054 9(72) · 10 <sup>-12</sup>	
ERM-AE701/8		1.052 4(71) · 10 <sup>-13</sup>	

Code	Description	Isotope amount fraction (-100)						Amount ratios			Unit size		
IRMM-009	0.2 M HNO <sub>3</sub> solution	<sup>24</sup> Mg	<sup>25</sup> Mg	<sup>26</sup> Mg				$n(^{25}\text{Mg})/n(^{24}\text{Mg})$	$n(^{26}\text{Mg})/n(^{24}\text{Mg})$		4 mL		
		78.992(25)	10.003(9)	11.005(19)				0.126 63(13)	0.139 32(26)				
IRMM-010	Pt metal	<sup>190</sup> Pt	<sup>192</sup> Pt	<sup>194</sup> Pt	<sup>195</sup> Pt	<sup>196</sup> Pt	<sup>198</sup> Pt	$n(^{190}\text{Pt})/n(^{195}\text{Pt})$	$n(^{192}\text{Pt})/n(^{195}\text{Pt})$	$n(^{194}\text{Pt})/n(^{195}\text{Pt})$	30 mg (wire)		
		0.011 7(11)	0.782(17)	32.86(27)	33.78(16)	25.21(23)	7.356(82)	0.000 347(34)	0.023 15(48)	0.973(11)			
								$n(^{196}\text{Pt})/n(^{195}\text{Pt})$	$n(^{198}\text{Pt})/n(^{195}\text{Pt})$				
IRMM-011	H <sub>3</sub> BO <sub>3</sub> solid	<sup>10</sup> B	<sup>11</sup> B					$n(^{10}\text{B})/n(^{11}\text{B})$			1 g		
		19.824 (20)	80.176(20)					0.247 26(32)					
IRMM-012	1 M HCl solution	<sup>50</sup> Cr	<sup>52</sup> Cr	<sup>53</sup> Cr	<sup>54</sup> Cr			$n(^{50}\text{Cr})/n(^{52}\text{Cr})$	$n(^{53}\text{Cr})/n(^{52}\text{Cr})$	$n(^{54}\text{Cr})/n(^{52}\text{Cr})$	5 mL		
		4.345(9)	83.789(2)	9.501(11)	2.365(5)			0.051 86(10)	0.113 39(15)	0.028 22(06)			
IRMM-016	Li <sub>2</sub> CO <sub>3</sub> solid	<sup>6</sup> Li	<sup>7</sup> Li					$n(^{6}\text{Li})/n(^{7}\text{Li})$			1 g		
		7.588 9(75)	92.411 1(75)					0.082 121(87)					
IRMM-017	Si solid	<sup>28</sup> Si	<sup>29</sup> Si	<sup>30</sup> Si				$n(^{29}\text{Si})/n(^{28}\text{Si})$	$n(^{30}\text{Si})/n(^{28}\text{Si})$		50 mg		
		92.228 77(86)	4.682 59(58)	3.088 64(70)				0.050 771 5(76)	0.033 488 9(82)				
IRMM-018a	SiO <sub>2</sub> solid	<sup>28</sup> Si	<sup>29</sup> Si	<sup>30</sup> Si				$n(^{29}\text{Si})/n(^{28}\text{Si})$	$n(^{30}\text{Si})/n(^{28}\text{Si})$		5 g		
		92.220 36(49)	4.687 30(36)	3.092 34(37)				0.050 827 2(40)	0.033 532 0(42)				

## 6.2 CERTIFIED FOR ISOTOPE AMOUNT CONTENT

Code	Description	Isotope amount content	Isotope enrichment	Unit size
IRMM-610	H <sub>3</sub> BO <sub>3</sub> aqueous solution	3.683 11 (88) µmol <sup>10</sup> B·g <sup>-1</sup>	$n(^{10}\text{B})/n(^{11}\text{B}) = 18.80$ (2)	5 mL
IRMM-611	H <sub>3</sub> BO <sub>3</sub> aqueous solution	4.025 (40) µmol <sup>11</sup> B·g <sup>-1</sup>	$n(^{10}\text{B})/n(^{11}\text{B}) = 0.247$ 26 (32)	5 mL
IRMM-615	0.5 M HCl solution	3.850 (14) µmol <sup>6</sup> Li·g <sup>-1</sup>	$n(^6\text{Li})/n(^7\text{Li}) = 21.897$ (44)	5 mL
IRMM-618	0.5 M HNO <sub>3</sub> solution	112.13 (17) µmol <sup>87</sup> Rb·kg <sup>-1</sup>	$n(^{85}\text{Rb})/n(^{87}\text{Rb}) = 0.20$ 498 (24)	5 mL
IRMM-619	0.5 M HNO <sub>3</sub> solution	85.00 (24) µmol <sup>85</sup> Rb·kg <sup>-1</sup>	$n(^{85}\text{Rb})/n(^{87}\text{Rb}) = 2.593$ 0 (20)	5 mL
IRMM-620	4.5 M HCl solution	173.35 (16) µmol <sup>57</sup> Fe·kg <sup>-1</sup>	$n(^{54}\text{Fe})/n(^{57}\text{Fe}) = < 0.0001$ $n(^{56}\text{Fe})/n(^{57}\text{Fe}) = 0.025$ 39 (31) $n(^{58}\text{Fe})/n(^{57}\text{Fe}) = 0.025$ 16 (18)	5 mL
IRMM-621	1 M HNO <sub>3</sub> solution	97.35 (15) µmol <sup>111</sup> Cd·kg <sup>-1</sup>	$n(^{106}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{108}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{110}\text{Cd})/n(^{111}\text{Cd}) = 0.004$ 44 (42) $n(^{112}\text{Cd})/n(^{111}\text{Cd}) = 0.021$ 74 (10) $n(^{113}\text{Cd})/n(^{111}\text{Cd}) = 0.005$ 818 (56) $n(^{114}\text{Cd})/n(^{111}\text{Cd}) = 0.010$ 875 (88) $n(^{116}\text{Cd})/n(^{111}\text{Cd}) = 0.001$ 629 (44)	4 mL
IRMM-622	1 M HNO <sub>3</sub> solution	9.739 (18) µmol <sup>111</sup> Cd·kg <sup>-1</sup>	$n(^{106}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{108}\text{Cd})/n(^{111}\text{Cd}) = < 0.000$ 05 $n(^{110}\text{Cd})/n(^{111}\text{Cd}) = 0.004$ 44 (42) $n(^{112}\text{Cd})/n(^{111}\text{Cd}) = 0.021$ 74 (10) $n(^{113}\text{Cd})/n(^{111}\text{Cd}) = 0.005$ 818 (56) $n(^{114}\text{Cd})/n(^{111}\text{Cd}) = 0.010$ 875 (88) $n(^{116}\text{Cd})/n(^{111}\text{Cd}) = 0.001$ 629 (44)	4 mL
IRMM-624	1 M HCl solution	174.84 (42) µmol <sup>50</sup> Cr·kg <sup>-1</sup>	$n(^{52}\text{Cr})/n(^{50}\text{Cr}) = 0.066$ 41 (50) $n(^{53}\text{Cr})/n(^{50}\text{Cr}) = 0.000$ 323 (64) $n(^{54}\text{Cr})/n(^{50}\text{Cr}) = 0.000$ 11 (11)	5 mL
IRMM-625	1 M HCl solution	144.233 (90) µmol <sup>52</sup> Cr·kg <sup>-1</sup>	$n(^{50}\text{Cr})/n(^{52}\text{Cr}) = 0.051$ 85 (20) $n(^{53}\text{Cr})/n(^{52}\text{Cr}) = 0.113$ 33 (38) $n(^{54}\text{Cr})/n(^{52}\text{Cr}) = 0.028$ 35 (34)	5 mL
IRMM-632	1 M HNO <sub>3</sub> solution	0.096 84 (41) µmol <sup>65</sup> Cu·g <sup>-1</sup>	$n(^{63}\text{Cu})/n(^{65}\text{Cu}) = 0.002$ 892 1 (92)	5 mL
ERM-AE633	1 M HNO <sub>3</sub> solution	5.998 (36) µmol <sup>63</sup> Cu·g <sup>-1</sup>	$n(^{65}\text{Cu})/n(^{63}\text{Cu}) = 0.445$ 63 (42)	4 - 5 mL
IRMM-634	1.8 M HCl solution	163.61 (38) µmol <sup>56</sup> Fe·kg <sup>-1</sup>	$n(^{54}\text{Fe})/n(^{56}\text{Fe}) = 0.063$ 70 (27) $n(^{57}\text{Fe})/n(^{56}\text{Fe}) = 0.023$ 096 (72) $n(^{58}\text{Fe})/n(^{56}\text{Fe}) = 0.003$ 071 (29)	5 mL
IRMM-635	1 M HNO <sub>3</sub> solution	58.131(37) µmol <sup>84</sup> Sr·kg <sup>-1</sup>	$n(^{86}\text{Sr})/n(^{84}\text{Sr}) = 0.000$ 589 (10) $n(^{87}\text{Sr})/n(^{84}\text{Sr}) = 0.000$ 098 (10) $n(^{88}\text{Sr})/n(^{84}\text{Sr}) = 0.000$ 386 (10)	5 mL
ERM-AE637	0.2 M HNO <sub>3</sub> solution	0.791 37(30) µmol <sup>24</sup> Mg·g <sup>-1</sup>	$n(^{26}\text{Mg})/n(^{24}\text{Mg}) = 0.139$ 68 (32) $n(^{25}\text{Mg})/n(^{24}\text{Mg}) = 0.126$ 86 (18)	5 mL
ERM-AE638	0.1 M HNO <sub>3</sub> solution	0.857 4 (34) µmol <sup>26</sup> Mg·g <sup>-1</sup>	$n(^{24}\text{Mg})/n(^{26}\text{Mg}) = 0.003$ 104 (26) $n(^{25}\text{Mg})/n(^{26}\text{Mg}) = 0.001$ 084 (11)	5 mL
ERM-AE639	0.5 M HCl solution	11.891 (50) µmol <sup>202</sup> Hg·g <sup>-1</sup>	$n(^{196}\text{Hg})/n(^{202}\text{Hg}) = 0.004$ 972 (46) $n(^{198}\text{Hg})/n(^{202}\text{Hg}) = 0.330$ 6 (21) $n(^{199}\text{Hg})/n(^{202}\text{Hg}) = 0.561$ 9 (28) $n(^{200}\text{Hg})/n(^{202}\text{Hg}) = 0.770$ 5 (28) $n(^{201}\text{Hg})/n(^{202}\text{Hg}) = 0.441$ 26 (88) $n(^{204}\text{Hg})/n(^{202}\text{Hg}) = 0.230$ 27 (75)	5 mL

<b>ERM-AE640</b>	0.5 M HCl solution	14.71 (11) nmol $^{202}\text{Hg}\cdot\text{g}^{-1}$	$n(^{196}\text{Hg})/n(^{202}\text{Hg}) = 0.000\ 018\ 09\ (38)$ $n(^{198}\text{Hg})/n(^{202}\text{Hg}) = 0.000\ 623\ (11)$ $n(^{199}\text{Hg})/n(^{202}\text{Hg}) = 0.001\ 603\ (16)$ $n(^{200}\text{Hg})/n(^{202}\text{Hg}) = 0.005\ 499\ (34)$ $n(^{201}\text{Hg})/n(^{202}\text{Hg}) = 0.013\ 351\ (52)$ $n(^{204}\text{Hg})/n(^{202}\text{Hg}) = 0.002\ 595\ (21)$	5 mL
<b>ERM-AE641</b>	Cl in water	18.959 (15) $\mu\text{mol } ^{35}\text{Cl}\cdot\text{g}^{-1}$	$n(^{37}\text{Cl})/n(^{35}\text{Cl}) = 0.319\ 77\ (83)$	4 - 5 mL
<b>ERM-AE642</b>	Cl in water	4.375 (26) $\mu\text{mol } ^{37}\text{Cl}\cdot\text{g}^{-1}$	$n(^{35}\text{Cl})/n(^{37}\text{Cl}) = 0.019\ 14\ (48)$	4 - 5 mL
<b>IRMM-643</b>	2.8 M HNO <sub>3</sub> solution	334.33 (84) $\mu\text{mol } ^{32}\text{S}\cdot\text{kg}^{-1}$	$n(^{33}\text{S})/n(^{32}\text{S}) = 0.007\ 877\ 6\ (58)$ $n(^{34}\text{S})/n(^{32}\text{S}) = 0.044\ 149\ 3\ (78)$ $n(^{36}\text{S})/n(^{32}\text{S}) = 0.000\ 153\ 40\ (94)$	5 mL
<b>IRMM-644</b>	3.2 M HNO <sub>3</sub> solution	326.28 (80) $\mu\text{mol } ^{32}\text{S}\cdot\text{kg}^{-1}$	$n(^{33}\text{S})/n(^{32}\text{S}) = 0.007\ 969\ 8\ (70)$ $n(^{34}\text{S})/n(^{32}\text{S}) = 0.045\ 162\ 2\ (82)$ $n(^{36}\text{S})/n(^{32}\text{S}) = 0.000\ 170\ 0\ (58)$	5 mL
<b>IRMM-645</b>	2.8 M HNO <sub>3</sub> solution	371.96 (57) $\mu\text{mol } ^{32}\text{S}\cdot\text{kg}^{-1}$	$n(^{33}\text{S})/n(^{32}\text{S}) = 0.007\ 747\ 6\ (38)$ $n(^{34}\text{S})/n(^{32}\text{S}) = 0.042\ 747\ 3\ (62)$ $n(^{36}\text{S})/n(^{32}\text{S}) = 0.000\ 145\ 1\ (42)$	5 mL
<b>IRMM-646</b>	2.8 M HNO <sub>3</sub> solution	4586 (27) $\mu\text{mol } ^{34}\text{S}\cdot\text{kg}^{-1}$	$n(^{32}\text{S})/n(^{34}\text{S}) = 0.038\ 314\ 9\ (31)$ $n(^{33}\text{S})/n(^{34}\text{S}) = 0.000\ 470\ 88\ (15)$ $n(^{36}\text{S})/n(^{34}\text{S}) = 0.000\ 018\ 1\ (37)$	5 mL
<b>ERM-AE647</b>	1 M HNO <sub>3</sub> solution	134.974 (73) $\mu\text{mol } ^{63}\text{Cu}\cdot\text{g}^{-1}$	$n(^{65}\text{Cu})/n(^{63}\text{Cu}) = 0.445\ 60\ (74)$	4 mL
<b>ERM-AE649</b>	1 M HNO <sub>3</sub> solution	0.836 88 (27) $\mu\text{mol } ^{205}\text{Tl}\cdot\text{g}^{-1}$	$n(^{203}\text{Tl})/n(^{205}\text{Tl}) = 0.418\ 91\ (18)$	4 - 5 mL
<b>IRMM-651</b>	0.5 M HNO <sub>3</sub> solution	0.077 506 (30) $\mu\text{mol } ^{64}\text{Zn}\cdot\text{g}^{-1}$	$n(^{66}\text{Zn})/n(^{64}\text{Zn}) = 0.557\ 17\ (30)$ $n(^{67}\text{Zn})/n(^{64}\text{Zn}) = 0.080\ 702\ (34)$ $n(^{68}\text{Zn})/n(^{64}\text{Zn}) = 0.366\ 27\ (12)$ $n(^{70}\text{Zn})/n(^{64}\text{Zn}) = 0.011\ 981\ (22)$	5 mL
<b>IRMM-652</b>	0.5 M HNO <sub>3</sub> solution	0.156 000 (50) $\mu\text{mol } ^{64}\text{Zn}\cdot\text{g}^{-1}$	$n(^{66}\text{Zn})/n(^{64}\text{Zn}) = 0.004\ 107\ 3\ (59)$ $n(^{67}\text{Zn})/n(^{64}\text{Zn}) = 0.000\ 499\ 87\ (96)$ $n(^{68}\text{Zn})/n(^{64}\text{Zn}) = 0.002\ 029\ 5\ (23)$ $n(^{70}\text{Zn})/n(^{64}\text{Zn}) = 0.000\ 052\ 76\ (34)$	5 mL
<b>IRMM-653</b>	0.5 M HNO <sub>3</sub> solution	0.138 014 (60) $\mu\text{mol } ^{67}\text{Zn}\cdot\text{g}^{-1}$	$n(^{64}\text{Zn})/n(^{67}\text{Zn}) = 0.013\ 191\ 5\ (81)$ $n(^{66}\text{Zn})/n(^{67}\text{Zn}) = 0.024\ 551\ 6\ (70)$ $n(^{68}\text{Zn})/n(^{67}\text{Zn}) = 0.051\ 086\ (36)$ $n(^{70}\text{Zn})/n(^{67}\text{Zn}) = 0.000\ 527\ 8\ (18)$	5 mL
<b>IRMM-654</b>	0.5 M HNO <sub>3</sub> solution	0.146 098 (48) $\mu\text{mol } ^{68}\text{Zn}\cdot\text{g}^{-1}$	$n(^{64}\text{Zn})/n(^{68}\text{Zn}) = 0.00\ 489\ 4\ (38)$ $n(^{66}\text{Zn})/n(^{68}\text{Zn}) = 0.003\ 186\ 8\ (46)$ $n(^{68}\text{Zn})/n(^{68}\text{Zn}) = 0.001\ 411\ 3\ (17)$ $n(^{70}\text{Zn})/n(^{68}\text{Zn}) = 0.000\ 194\ 98\ (78)$	5 mL
<b>IRMM-3702</b>	1 M HNO <sub>3</sub> solution	1.512 (30) $\mu\text{mol } ^{64}\text{Zn}\cdot\text{g}^{-1}$	$n(^{66}\text{Zn})/n(^{64}\text{Zn}) = 0.563\ 97\ (30)$ $n(^{67}\text{Zn})/n(^{64}\text{Zn}) = 0.082\ 166\ (35)$ $n(^{68}\text{Zn})/n(^{64}\text{Zn}) = 0.375\ 19\ (16)$ $n(^{70}\text{Zn})/n(^{64}\text{Zn}) = 0.012\ 418\ (23)$	3 mL

## **INDEX**

### **NUMERICAL LIST**

ERM / CRM	DESIGNATION	PAGE NO.
ERM-AC057	AFLATOXIN B1 IN ACETONITRILE	27
ERM-AC058	AFLATOXIN B2 IN ACETONITRILE	27
ERM-AC059	AFLATOXIN G1 IN ACETONITRILE	27
ERM-AC060	AFLATOXIN G2 IN ACETONITRILE	27
ERM-AC213	PAHs IN ACETONITRILE / TOLUENE	9
ERM-AC699	ZEARALENONE IN ACETONITRILE	26
ERM-AD148	THROMBOPLASTIN BOVINE (OBT/79) (prothrombin time)	58
ERM-AD149	THROMBOPLASTIN RABBIT (prothrombin time)	58
ERM-AD413	PLASMID DNA FRAGMENTS OF MON 810 MAIZE	29
ERM-AD415	PLASMID DNA FRAGMENTS OF NK603 MAIZE	30
ERM-AD425	PLASMID DNA FRAGMENTS OF 356043 SOYBEAN	33
ERM-AD427	PLASMID DNA FRAGMENTS OF 98140 MAIZE	34
ERM-AD452/IFCC	$\gamma$ -GLUTAMYLTRANSFERASE (catalytic concentration)	57
ERM-AD454/IFCC	ALANINE AMINOTRANSFERASE (catalytic concentration)	57
ERM-AD455/IFCC	CREATINE KINASE (CK-MB iso-enzyme) (catalytic concentration)	57
ERM-AD457/IFCC	ASPARTATE TRANSAMINASE (AST)	57
ERM-AD623	BCR-ABL pDNA CALIBRANT	58
ERM-AE633	COPPER (natural) spike, nitrate solution	75
ERM-AE637	MAGNESIUM (natural) spike, nitrate solution	75
ERM-AE638	MAGNESIUM-26 spike, nitrate solution	75
ERM-AE639	MERCURY (natural) spike, chloride solution	75
ERM-AE640	MERCURY-202 spike, chloride solution	76
ERM-AE641	CHLORIDE (natural) spike, chloride solution	76
ERM-AE642	CHLORIDE-37 spike, chloride solution	76
ERM-AE647	COPPER-63, nitrate solution	76
ERM-AE649	THALLIUM (natural) spike, nitrate solution	76
ERM-AE701	CALCIUM-41 isotopic, nitrate solution (set of 8 units)	74
ERM-BB124	PORK MUSCLE	51
ERM-BB130	PORK MUSCLE	50
ERM-BB184	BOVINE MUSCLE (trace elements)	40
ERM-BB186	PIG KIDNEY (trace elements)	40
ERM-BB350	FISH OIL	37
ERM-BB384	PORK MUSCLE	42, 44
ERM-BB386	BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol) (blank)	50
ERM-BB389	BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol)	50
ERM-BB422	FISH MUSCLE (trace elements)	40
ERM-BB430	PORK FAT (pesticides)	37
ERM-BB444	NATURAL PORK FAT (blank)	36
ERM-BB445	SPIKED PORK FAT (very low level)	36
ERM-BB446	SPIKED PORK FAT (low level)	36
ERM-BB492	MILK POWDER (oxytetracycline)	50
ERM-BB493	MILK POWDER (oxytetracycline) (blank)	50
ERM-BC190	RAPSEED (colza) (S, total glucosinolate, medium level)	38
ERM-BC211	RICE (As species)	21
ERM-BC366	RAPSEED (colza) (S, total glucosinolate, low level)	38
ERM-BC367	RAPSEED (colza) (S, total glucosinolate, high level)	38
ERM-BC381	RYE FLOUR	42, 44

ERM-BC382	WHEAT FLOUR	42, 44
ERM-BC514	HARICOTS BEANS (dietary fibre)	46
ERM-BC515	CARROT (dietary fibre)	46
ERM-BC516	APPLE (dietary fibre)	46
ERM-BC517	FULL FAT SOYA (dietary fibre)	46
ERM-BC716	MAIZE	40
ERM-BC717	MAIZE	40
ERM-BD150	SKIMMED MILK POWDER (trace elements)	40
ERM-BD151	SKIMMED MILK POWDER (trace elements)	40
ERM-BD273	TOASTED BREAD	52
ERM-BD282	WHOLE MILK POWDER (aflatoxin M1, zero level)	39
ERM-BD283	WHOLE MILK POWDER (aflatoxin M1, low level)	39
ERM-BD284	WHOLE MILK POWDER (aflatoxin M1, high level)	39
ERM-BD518	BRAN BREAKFAST CEREAL (dietary fibre)	46
ERM-BD600	WHOLE MILK POWDER (vitamins)	44
ERM-BE375	COMPOUND FEEDINGSTUFF (very low level)	39
ERM-BE376	COMPOUND FEEDINGSTUFF (high level)	39
ERM-BF410k	GENETICALLY MODIFIED ROUNDUP READY SOYA	28
ERM-BF411	GENETICALLY MODIFIED Bt-176 MAIZE	28
ERM-BF412	GENETICALLY MODIFIED Bt-11 MAIZE	28
ERM-BF413k	GENETICALLY MODIFIED MON 810 MAIZE	29
ERM-BF414	GENETICALLY MODIFIED GA21 MAIZE	29
ERM-BF415	GENETICALLY MODIFIED NK603 MAIZE	30
ERM-BF416	GENETICALLY MODIFIED MON 863 MAIZE	30
ERM-BF417	GENETICALLY MODIFIED MON 863 x MON 810 MAIZE	30
ERM-BF418	GENETICALLY MODIFIED 1507 MAIZE	31
ERM-BF419	GENETICALLY MODIFIED H7-1 SUGAR BEET	31
ERM-BF420	GENETICALLY MODIFIED 3272 MAIZE	31
ERM-BF421	GENETICALLY MODIFIED EH92-527-1 POTATO	31
ERM-BF422	GENETICALLY MODIFIED 281-24-236 X 3006-210-23 COTTON SEED	32
ERM-BF423	GENETICALLY MODIFIED MIR604 MAIZE	32
ERM-BF424	GENETICALLY MODIFIED 59122 MAIZE	32
ERM-BF425	GENETICALLY MODIFIED SOYA 356043	32
ERM-BF426	GENETICALLY MODIFIED SOYA 305423	33
ERM-BF427	GENETICALLY MODIFIED 98140 MAIZE	33
ERM-BF428	GENETICALLY MODIFIED GHB119 COTTON	34
ERM-BF429	GENETICALLY MODIFIED T304-40 COTTON	34
ERM-BF430	GENETICALLE MODIFIED AM04-1020 POTATO	34
ERM-BF431	GENETICALLE MODIFIED AV43-6-G7 POTATO	34
ERM-BF432	GENETICALLY MODIFIED SOYA DAS-68416-4	35
ERM-BF433	GENETICALLY MODIFIED DAS-40278-9 MAIZE	35
ERM-BF434	GENETICALLY MODIFIED 73496 RAPSEED	35
ERM-BF435	GENETICALLY MODIFIED PH05-026-0048 POTATO	35
ERM-BF436	GENETICALLY MODIFIED DAS-44406-6 SOYA	36
ERM-CA408	SIMULATED RAINWATER (major components)	17
ERM-CA615	GROUND WATER	18
ERM-CA616	GROUND WATER	18
ERM-CA713	WASTE WATER	18
ERM-CC141	LOAM SOIL	10
ERM-CC580	ESTUARINE SEDIMENT (Hg, methylmercury)	11, 20
ERM-CC690	CALCAREOUS SOIL	10
ERM-CD200	BLADDERWRACK ( <i>Fucus vesiculosus</i> ) (trace elements)	15
ERM-CD281	RYE GRASS	14
ERM-CE195	BOVINE BLOOD (Pb, Cd)	54

ERM-CE196	BOVINE BLOOD (Pb, Cd)	54
ERM-CE278k	MUSSEL TISSUE (elements)	16
ERM-CE464	TUNA FISH (total and methylmercury)	16, 21
ERM-CE477	MUSSEL TISSUE (butyltins)	21
ERM-CZ100	FINE DUST (PM <sub>10</sub> -like) (PAHs)	21
ERM-CZ120	FINE DUST (PM <sub>10</sub> -like) (elements)	13
ERM-DA192	HUMAN SERUM (cortisol unspiked)	54
ERM-DA193	HUMAN SERUM (cortisol spiked)	54
ERM-DA347	HUMAN SERUM (progesterone)	54
ERM-DA451/IFCC	CORTISOL REFERENCE SERUM PANEL	54
ERM-DA470k/IFCC	HUMAN SERUM (proteins)	56
ERM-DA471/IFCC	HUMAN SERUM (cystatin C)	56
ERM-DA474/IFCC	HUMAN SERUM (CRP)	56
ERM-DB001	HUMAN HAIR (trace elements)	55
ERM-EB322	UNALLOYED ZINC (trace elements)	68
ERM-EB323	UNALLOYED ZINC (trace elements)	68
ERM-EB324	UNALLOYED ZINC (trace elements)	68
ERM-EB325	UNALLOYED ZINC (trace elements)	68
ERM-EC590	POLYETHYLENE (LDPE)	72
ERM-EC591	POLYPROPYLENE (PP)	72
ERM-EC680k	POLYETHYLENE (low level)	72
ERM-EC681k	POLYETHYLENE (high level)	72
ERM-EF104	GAS OIL (0.1019 % S)	70
ERM-EF211	PETROL	70
ERM-EF317	GAS OIL (Solvent Yellow 124)	70
ERM-EF318	GAS OIL (Solvent Yellow 124)	70
ERM-EF411	HARD COAL	60
ERM-EF412	BROWN COAL	60
ERM-EF413	FURNACE COKE	60
ERM-EF671	GAS OIL (0.0452 % S)	70
ERM-EF672	GAS OIL (0.0203 % S)	70
ERM-EG001	ANTIMONY IMPLANTED IN SILICON	72
ERM-FA013	CHARPY SPECIMENS Low Energy 20°C (impact toughness)	62
ERM-FA013	CHARPY SPECIMENS Low Energy 0°C (impact toughness)	62
ERM-FA015	CHARPY SPECIMENS 80 J (impact toughness)	62
ERM-FA016	CHARPY SPECIMENS 120 J (impact toughness)	62
ERM-FA415	CHARPY SPECIMENS 150 J (impact toughness)	62
ERM-FC395k	GAS OIL (CFPP and CP)	59
ERM-FD100	COLLOIDAL SILICA	64
ERM-FD304	COLLOIDAL SILICA	64
BCR-010	TIN ORE CONCENTRATE (Sn)	65
BCR-017A, B	COPPER (S, P)	67
BCR-022A, B	ELECTROLYTIC TOUGH PITCH COPPER (O)	67
BCR-024B, C	TITANIUM (O, N)	67
BCR-032	MOROCCAN PHOSPHATE ROCK (trace elements)	66, 71
BCR-033	SUPERPHOSPHATE (various parameters)	65
BCR-038	FLY ASH FROM PULVERISED COAL (trace elements)	12
BCR-046	BENZO[b]CHRYSENE (purity)	1
BCR-047	BENZO[b]FLUORANTHENE (purity)	1
BCR-048R	BENZO[k]FLUORANTHENE (purity)	1
BCR-049	BENZO[jj]FLUORANTHENE (purity)	1
BCR-050	BENZO[e]PYRENE (purity)	1
BCR-052	BENZO[ghi]PERYLENE (purity)	1

BCR-054R	COPPER (O)	67
BCR-058	CONTINUOUS CAST COPPER (O)	67
BCR-059A, B	Ti 6AL 4V ALLOY (O)	67
BCR-060	AQUATIC PLANT ( <i>Lagarosiphon major</i> ) (trace elements)	13
BCR-063R	SKIM MILK POWDER (major and trace elements)	15
BCR-066	QUARTZ (particle size 0.35 – 3.50 microns)	62
BCR-067	QUARTZ (particle size 2.40 – 32.00 microns)	62
BCR-068	QUARTZ (particle size 160 – 630 microns)	62
BCR-069	QUARTZ (particle size 14 – 90 microns)	62
BCR-070	QUARTZ (1.20 – 20.00 microns)	62
BCR-074A	ELECTROLYTIC COPPER (trace elements)	69
BCR-077R	1-METHYLCHRYSENE (purity)	1
BCR-078R	2-METHYLCHRYSENE (purity)	1
BCR-079R	3-METHYLCHRYSENE (purity)	1
BCR-080R	4-METHYLCHRYSENE (purity)	1
BCR-081R	5-METHYLCHRYSENE (purity)	1
BCR-082R	6-METHYLCHRYSENE (purity)	1
BCR-089	TiAl6V4 (Al, V)	69
BCR-090A, B	TITANIUM (impurities)	69
BCR-091	ANTHANTHRENE (purity)	1
BCR-092	10-AZABENZO[a]PYRENE (purity)	1
BCR-093R	1-METHYLBENZ[a]ANTHRACENE (purity)	1
BCR-094	DIBENZ[a,c]ANTHRACENE (purity)	1
BCR-095	DIBENZ[a,j]ANTHRACENE (purity)	1
BCR-096	DIBENZO[a,l]PYRENE (80mylas)	1
BCR-097	BENZO[a]FLUORANTHENE (purity)	1
BCR-098	ZIRCALOY-4 (trace element impurities)	69
BCR-099	NICKEL (N, O)	67
BCR-100	BEECH LEAVES (trace elements)	13
BCR-102	TUNGSTEN CARBIDE POWDER (O)	67
BCR-105	GAS OIL (0.363 % S)	70
BCR-106	GAS OIL (0.502 % S)	70
BCR-107	GAS OIL (1.040 % S)	70
BCR-109	ZINC ORE CONCENTRATE (trace elements)	71
BCR-110	ZINC ORE CONCENTRATE (trace elements)	71
BCR-113	POTASSIUM CHLORIDE FERTILIZER (elemental composition)	65
BCR-114	POTASSIUM SULPHATE FERTILIZER (elemental composition)	65
BCR-115	ANIMAL FEED (Organochlorine pesticides)	38
BCR-116	LIMESTONE POWDERS (for shear testing)	60
BCR-121	WHOLEMEAL FLOUR (vitamins)	44
BCR-122	MARGARINE (vitamins)	44
BCR-123A, B	3 REFERENCE ETHANOLS (H, M, L) (for SNIF-NMR)	25, 73
BCR-126A	LEAD GLASS (composition/refractive index)	66
BCR-129	HAY POWDER (elements)	13
BCR-130	QUARTZ (particle size 50 – 220 microns)	62
BCR-131	QUARTZ (particle size 480 – 1800 microns)	62
BCR-132	QUARTZ (particle size 1400 – 5000 microns)	62
BCR-133	DIBENZO[a,e]PYRENE (80mylas)	1
BCR-134	BENZO[c]PHENANTHRENE (purity)	1
BCR-136R	BENZO[b]NAPHTHO[2,3-d]THIOPHENE (purity)	1
BCR-137R	BENZO[b]NAPHTHO[1,2-d]THIOPHENE (purity)	1
BCR-138	DIBENZO[a,h]ANTHRACENE (purity)	1
BCR-139	BENZO[ghi]FLUORANTHENE (purity)	1
BCR-140	BENZO[c]CHRYSENE (purity)	1

BCR-142R	LIGHT SANDY SOIL (trace elements)	10
BCR-143R	SEWAGE SLUDGE AMENDED SOIL (trace elements)	10
BCR-145R	SEWAGE SLUDGE (mixed origin) (trace elements)	11
BCR-146R	SEWAGE SLUDGE (industrial origin) (trace elements)	11
BCR-152	DIBENZ[a,I]ACRIDINE (81mylas)	1
BCR-153R	DIBENZ[a,h]ACRIDINE (purity)	1
BCR-154	DIBENZ[a,j]ACRIDINE (purity)	1
BCR-155	DIBENZ[a,c]ACRIDINE (81mylas)	1
BCR-156R	DIBENZ[c,h]ACRIDINE (purity)	1
BCR-157	BENZ[a]ACRIDINE (purity)	1
BCR-158	BENZ[c]ACRIDINE (purity)	1
BCR-159	DIBENZO[a,h]PYRENE (81mylas)	1
BCR-160R	FLUORANTHENE (purity)	1
BCR-162R	SOYA-MAIZE OIL BLEND (fatty acid profile)	42
BCR-163	BEEF-PORK FAT BLEND (fatty acid profile)	42
BCR-165	LATEX SPHERES (particle diameter 2 microns)	53
BCR-166	LATEX SPHERES (particle diameter 4.8 microns)	53
BCR-167	LATEX SPHERES (particle diameter 9.6 microns)	53
BCR-168	PICENE (purity)	1
BCR-169	ALPHA ALUMINA (0.10 m <sup>2</sup> /g) (nitrogen BET specific surface area)	62
BCR-170	ALPHA ALUMINA (1.05 m <sup>2</sup> /g) (nitrogen BET specific surface area)	62
BCR-171	ALUMINA (2.95 m <sup>2</sup> /g) (nitrogen BET specific surface area)	62
BCR-172	QUARTZ (2.50 m <sup>2</sup> /g) (nitrogen BET specific surface area)	62
BCR-173	TITANIA (8.23 m <sup>2</sup> /g) (nitrogen BET specific surface area)	62
BCR-175	TUNGSTEN (0.18 m <sup>2</sup> /g) (nitrogen BET specific surface area)	62
BCR-176R	FLY ASH (trace elements)	12
BCR-177R	PYRENE (purity)	1
BCR-178	CALCIUM AMMONIUM NITRATE FERTILIZER (composition)	65
BCR-179	UREA FERTILIZER (composition)	65
BCR-185R	BOVINE LIVER (trace elements)	40
BCR-187	NATURAL MILK POWDER (pesticides)	38
BCR-188	SPIKED MILK POWDER (pesticides)	38
BCR-191	BROWN BREAD (trace elements)	40
BCR-261T	TANTALUM PENTOXIDE ON TANTALUM FOIL	64
BCR-262R	DEFATTED PEANUT MEAL (aflatoxin B1, blank)	39
BCR-263R	DEFATTED PEANUT MEAL (aflatoxin B1, medium level)	39
BCR-264	DEFATTED PEANUT MEAL (aflatoxin B1, high level)	39
BCR-265	DIBENZO[a,e]FLUORANTHENE (81mylas)	1
BCR-266	7H-DIBENZO (c,g) CARBAZOLE (purity)	1
BCR-267	INDENO[1,2,3-cd]FLUORANTHENE (purity)	1
BCR-269	CHRYSENE (purity)	1
BCR-270	TRIPHENYLENE (purity)	1
BCR-271	BENZ[a]ANTHRACENE (purity)	1
BCR-272	CORONENE (purity)	1
BCR-273	SINGLE CELL PROTEIN (major elements)	41
BCR-274	SINGLE CELL PROTEIN (trace elements)	41
BCR-275	ZIRCALOY (C, N, O)	67
BCR-276	ZIRCALOY (C, N, O)	67
BCR-277R	ESTUARINE SEDIMENT (trace elements)	11
BCR-280R	LAKE SEDIMENT (trace elements)	11
BCR-286A, B	ELECTROLYTICALLY REFINED LEAD (trace elements)	68
BCR-287A, B	THERMALLY REFINED LEAD (trace elements)	68
BCR-288B	LEAD WITH ADDED IMPURITIES (trace elements)	68
BCR-289	2,4'-DICHLOROBIPHENYL (IUPAC No. 8) (purity)	2

BCR-290	2,3,3'-TRICHLOROBIPHENYL (IUPAC No. 20) (purity)	2
BCR-291	2,4,4'-TRICHLOROBIPHENYL (IUPAC No. 28) (purity)	2
BCR-293	2,2',5,5'-TETRACHLOROBIPHENYL (IUPAC No. 52) (purity)	2
BCR-296	2,2',3,4,4',5'-HEXACHLOROBIPHENYL (IUPAC No. 138) (purity)	2
BCR-297	2,2',4,4',5,5'-HEXACHLOROBIPHENYL (IUPAC No. 153) (purity)	2
BCR-298	2,2',3,4,4',5,5'-HEPTACHLOROBIPHENYL (IUPAC No. 180) (purity)	2
BCR-301 (RM)	MULLITE (lattice spacing, other parameters)	63
BCR-302	MICROCRYSTALLINE CELLULOSE (water content above saturated solutions)	63
BCR-304	HUMAN SERUM (Ca, Mg, Li)	55
BCR-305	1-NITROPYRENE (purity)	2
BCR-306	1-NITRONAPHTALENE (purity)	2
BCR-307	2-NITRONAPHTALENE (purity)	2
BCR-308	9-NITROANTHRACENE (purity)	2
BCR-309	6-NITROCHRYSENE (purity)	2
BCR-310	3-NITROFLUORANTHENE (purity)	2
BCR-311	6-NITROBENZO[a]PYRENE (purity)	2
BCR-312	2-NITRO-7-METHOXYNAPHTHO[2,1-b]FURAN (purity)	2
BCR-318	TITANIUM (H)	67
BCR-320R	CHANNEL SEDIMENT (trace elements)	11
BCR-321	UNALLOYED ZINC (trace elements)	68
BCR-326	UNALLOYED ZINC (disc) (trace elements)	68
BCR-327	UNALLOYED ZINC (disc) (trace elements)	68
BCR-331	LOW VOLATILE STEAM COAL (S)	70
BCR-332	HIGH VOLATILE INDUSTRIAL COAL (S)	70
BCR-333	COKING STEAM COAL (S)	70
BCR-334	ANTHRACITE (S)	70
BCR-335	FLAME COAL (S)	70
BCR-336	HIGH VOLATILE STEAM COAL (S)	70
BCR-337	DIBENZO[b,d]FURAN (82mylas)	2
BCR-338	4H-CYCLOPENTA[def]PHENANTHREN-4-ONE (purity)	2
BCR-339	BENZO[c,d]PYREN-6-ONE (purity)	2
BCR-340	BENZO[b]NAPHTHO (1,2-d) FURAN (purity)	2
BCR-341	BENZO[b]NAPHTHO (2,1-d) FURAN (purity)	2
BCR-342	BENZO[a]FLUORENONE (purity)	2
BCR-348R	HUMAN SERUM (high progesterone)	54
BCR-349	COD LIVER OIL (PCBs)	37
BCR-351	ZnAl4 (trace elements)	68
BCR-352	ZnAl4 (trace elements)	68
BCR-353	ZnAl4 (trace elements)	68
BCR-354	ZnAl4 (trace elements)	68
BCR-355	ZnAl4 (trace elements)	68
BCR-356	ZnAl4Cu1 (trace elements)	68
BCR-357	ZnAl4Cu1 (trace elements)	68
BCR-359	ZnAl4Cu1 (trace elements)	68
BCR-360	ZnAl4Cu1 (trace elements)	68
BCR-361	ZnAl4Cu1 (trace elements)	68
BCR-365	PCB STANDARD SOLUTION	2
BCR-375	COMPOUND FEED (aflatoxin B1 blank)	39
BCR-377	MAIZE FLOUR (deoxynivalenol blank)	39
BCR-380R	WHOLE MILK POWDER (major nutrients)	45
BCR-383	HARICOTS VERTS (major nutrients)	42, 44
BCR-385R	PEANUT BUTTER (aflatoxins low level)	39
BCR-386	BOVINE URINE (diethylstilboestrol blank)	49
BCR-387	BOVINE URINE (dienoestrol blank)	49

BCR-388	BOVINE URINE (hexoestrol blank)	49
BCR-390 (RM)	BOVINE URINE (dienoestrol positive)	49
BCR-391	BOVINE URINE (hexoestrol positive)	49
BCR-393	HUMAN APOLIPOPROTEIN A I (mass concentration)	55
BCR-395	GAS OIL (CFPP 6°C)	59
BCR-396	WHEAT FLOUR (deoxynivalenol blank)	38
BCR-400	TOMATO PASTE COLOUR REFERENCE TILE (colour values)	64
BCR-401R	PEANUT BUTTER (aflatoxins very low level)	39
BCR-402	WHITE CLOVER (trace elements)	13
BCR-405 (RM)	HUMAN HAEMOLYSATE (glycated haemoglobin (HbA <sub>1c</sub> )	56
BCR-410	HUMAN PROSTATIC ACID PHOSPHATASE (catalytic concentration)	57
BCR-411	BOVINE MUSCLE (diethylstilboestrol positive)	50
BCR-412	BOVINE MUSCLE (diethylstilboestrol blank)	50
BCR-414	PLANKTON (trace elements)	14
BCR-420	WASTE MINERAL OIL (low PCB level)	24
BCR-423 (RM)	AFLATOXIN M1 STANDARD SOLUTION	26
BCR-425	NIMONIC 75 FOR CREEP TESTING	61
BCR-431	BRUSSELS SPROUT (vitamins)	44
BCR-444	PORCINE MUSCLE (chloramphenicol blank)	50
BCR-445	PORCINE MUSCLE (chloramphenicol positive)	50
BCR-446	RAPESSEED (oil, moisture, volatiles)	45
BCR-447	RAPESSEED (oil, moisture, volatiles)	45
BCR-449	WASTE MINERAL OIL (high PCB level)	24
BCR-450	NATURAL MILK POWDER (PCBs )	36
BCR-457	HUMAN THYROGLOBULIN (Tg) (mass concentration)	55
BCR-458	COCONUT OIL (PAH doped)	36
BCR-459	COCONUT OIL (PAH blank)	36
BCR-460	COAL (F)	70
BCR-461	CLAY (F)	70
BCR-462	COASTAL SEDIMENT (butyltins)	20
BCR-463	TUNA FISH (total and methylmercury)	16, 21
BCR-465	RICE FLOUR (83mylase, low level)	44
BCR-466	RICE FLOUR (83mylase, medium level)	44
BCR-467	RICE FLOUR (83mylase, high level)	44
BCR-471	WHEAT (ochratoxin A, blank)	39
BCR-474-5	BOVINE LIVER (trenbolone blank and positive)	50
BCR-479	FRESH WATER (nitrate, low level)	17
BCR-480	FRESH WATER (nitrate, high level)	17
BCR-481	INDUSTRIAL SOIL (PCBs )	22
BCR-482	LICHEN (trace elements)	15
BCR-483	SEWAGE SLUDGE AMENDED SOIL (trace elements)	19
BCR-484	SEWAGE SLUDGE AMENDED (terra rossa) SOIL (trace elements)	19
BCR-485	MIXED VEGETABLES (vitamins)	44
BCR-486	PURIFIED HUMAN ALFAFOETOPROTEIN (protein mass)	55
BCR-487	PIG LIVER (vitamins)	44
BCR-490	FLY ASH (PCDDs and PCDFs)	23
BCR-492	MILK POWDER (oxytetracycline)	50
BCR-493	MILK POWDER (oxytetracycline) (blank)	50
BCR-502	BOVINE URINE (clenbuterol and salbutamol)	49
BCR-503	BOVINE URINE (clenbuterol and salbutamol)	49
BCR-504	BOVINE URINE (clenbuterol and salbutamol)	49
BCR-505	ESTUARINE WATER (trace elements)	17
BCR-519	ANHYDROUS BUTTER FAT (triglycerides)	43
BCR-522	BOVINE BLOOD LYSATE (haemiglobincyanide)	56

BCR-524	INDUSTRIAL SOIL (PAHs)	22
BCR-528	BACILLUS CEREUS (number of colony forming particles)	47
BCR-529	INDUSTRIAL SANDY SOIL (PCDDs, PCDFs)	22
BCR-535	FRESHWATER HARBOUR SEDIMENT (PAHs)	22
BCR-536	FRESHWATER HARBOUR SEDIMENT (PCBs)	22
BCR-537	PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film A)	46
BCR-538	PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film B)	46
BCR-539	PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film C)	46
BCR-543	MUSSEL (dc-saxitoxin)	40
BCR-545	WELDING DUST LOADED ON FILTER (Cr VI, Cr)	21
BCR-546	FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	52
BCR-547	ACETALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	52
BCR-548	ACROLEIN-2,4-DINITROPHENYLHYDRAZONE (purity)	52
BCR-549	ACETONE-2,4-DINITROPHENYLHYDRAZONE (purity)	52
BCR-550	GLUTARALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	52
BCR-551	2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE	52
BCR-552	2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE (blank)	52
BCR-553	FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter	53
BCR-554	FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter (blank)	53
BCR-555	CHLORINATED HYDROCARBONS ON TENAX	53
BCR-563	COMMON WHEAT FLOUR (properties)	46
BCR-564	SiO <sub>2</sub> /Si (10,20,30 nm)	63
BCR-573	HUMAN SERUM (low creatinine)	56
BCR-573i	CREATININE (interfering substances)	57
BCR-574	HUMAN SERUM (medium creatinine)	56
BCR-575	HUMAN SERUM (high creatinine)	56
BCR-576	HUMAN SERUM (17 $\beta$ -ESTRADIOL, low level)	54
BCR-577	HUMAN SERUM (17 $\beta$ -ESTRADIOL, medium level)	54
BCR-578	HUMAN SERUM (17 $\beta$ -ESTRADIOL, high level)	54
BCR-579	COASTAL SEAWATER (Hg)	17
BCR-596	AQUATIC PLANT (Cr)	13
BCR-597	SEWAGE SLUDGE (Cr)	11
BCR-598	COD LIVER OIL (Organochlorine pesticides)	37
BCR-599	EWES'/GOATS' CURD (for adulteration with cows' milk) 0 and 1 % cows' milk	51
BCR-605	URBAN DUST (trimethyllead)	20
BCR-607	MILK POWDER (PCDDs, PCDFs)	38
BCR-609	GROUND WATER (trace elements, low level)	18
BCR-610	GROUND WATER (trace elements, high level)	18
BCR-611	GROUND WATER (Br, high level)	17
BCR-612	GROUND WATER (Br, low level)	17
BCR-613	PROSTATE SPECIFIC ANTIGEN (protein mass)	56
BCR-614	POLYCHLORODIBENZO-P-DIOXINS (PCDD) AND POLYCHLORODIBENZOFURANS (PCDFs)	3
BCR-615	FLY ASH (LOW LEVEL) (PCDDs and PCDFs)	23
BCR-617	ARTIFICIAL GROUND WATER (trace elements, low carbonate content)	18
BCR-627	TUNA FISH TISSUE (As species)	21
BCR-632	BUTTER FAT	43
BCR-633	ANHYDROUS BUTTER FAT (tracers)	43
BCR-634	HUMAN BLOOD (Pb, Cd)	54
BCR-635	HUMAN BLOOD (Pb, Cd)	54
BCR-636	HUMAN BLOOD (Pb, Cd)	54
BCR-637	HUMAN SERUM (Al, Se, Zn)	55
BCR-638	HUMAN SERUM (Al, Se, Zn)	55
BCR-639	HUMAN SERUM (Al, Se, Zn)	55
BCR-644	ARTIFICIAL FOODSTUFF (major nutrients)	44

BCR-645	ARTIFICIAL FOODSTUFF (major nutrients)	44
BCR-646	FRESHWATER SEDIMENT (butyltin and phenyltin compounds)	20
BCR-647	HUMAN ADENOSINE DEAMINASE (ADA 1)	57
BCR-648	BOVINE LIVER (CLENBUTEROL BLANK)	50
BCR-649	BOVINE LIVER (CLENBUTEROL POSITIVE)	50
BCR-651	BEER (EtOH, low level)	44
BCR-652	BEER (EtOH, very low level)	44
BCR-653	WINE (EtOH, low level)	44
BCR-656	ISOTOPE RATIOS IN ABSOLUTE ALCOHOL	25, 73
BCR-657	ISOTOPE RATIOS IN GLUCOSE	26, 73
BCR-658	ISOTOPE RATIOS IN SYNTHETIC WINE	26, 73
BCR-659	ISOTOPE RATIOS IN SYNTHETIC WINE	26, 73
BCR-660	ISOTOPE RATIOS IN ALCOHOLIC SOLUTION	26, 73
BCR-661	NIMONIC 75 FOR TENSILE PROPERTIES	61
BCR-663	SAXITOXIN IN ACETIC ACID	26
BCR-664	GLASS (trace elements)	71
BCR-665	LUNG TISSUE (asbestos fibres)	58
BCR-666	LUNG TISSUE (asbestos fibres)	58
BCR-667	ESTUARINE SEDIMENT	10
BCR-668	MUSSEL TISSUE	16
BCR-670	LEMNA MINOR (aquatic plant)	15
BCR-673	BOVINE EYE (CLENBUTEROL BLANK)	50
BCR-674	BOVINE EYE (CLENBUTEROL POSITIVE)	50
BCR-677	SEWAGE SLUDGE (PCDDs and PCDFs)	23
BCR-679	TRACE ELEMENTS IN WHITE CABBAGE	41
BCR-682	MUSSEL TISSUE	24
BCR-683	BEECH WOOD	24
BCR-684	RIVER SEDIMENT (extractable phosphorous)	19
BCR-685	SKIM MILK POWDER	45
BCR-691	COPPER ALLOYS	69
BCR-692	SCRATCH TESTING	61
BCR-693	HUMAN PANCREATIC LIPASE (from pancreatic juice)	57
BCR-694	HUMAN PANCREATIC LIPASE (recombinant)	57
BCR-695	PIG LIVER (CTC free)	50
BCR-696	PIG LIVER (CTC incurred)	50
BCR-697	PIG MUSCLE (CTC free)	50
BCR-700	ORGANIC-RICH SOIL (extractable elements)	19
BCR-701	LAKE SEDIMENT (trace elements)	20
BCR-704	FAUJASITE TYPE ZEOLITE (micropore volume and width)	63
BCR-705	LINDE TYPE A ZEOLITE (micropore volume and width)	63
BCR-706	PIG KIDNEY (CTC free)	50
BCR-707	PIG KIDNEY (CTC incurred)	50
BCR-708	DAIRY FEED (nutritional properties)	45
BCR-709	PIG FEED (nutritional properties)	45
BCR-718	HERRING (PCBs)	24
BCR-719	CHUB (PCBs)	24
BCR-723	ROAD DUST (trace elements)	12
BCR-724A-E	GLASS-CERAMIC	59
BCR-725	SALMON TISSUE	51
IRMM/IFCC-456	ALPHA-AMYLASE (catalytic concentration)	57
IRMM/IFCC-467	HAEMOGLOBIN HbA0	57
IRMM-007/1-6	Zinc-64, nitrate solution	73
IRMM-009	Mg, isotopic, nitrate solution	74
IRMM-010	PLATINUM, isotopic, metal	74

IRMM-011	BORIC ACID, isotopic, solid	74
IRMM-012	CHROMIUM, isotopic, chloride solution	74
IRMM-016	LITHIUM CARBONATE, isotopic, solid	74
IRMM-017	SILICON, isotopic, Si single crystal	74
IRMM-018a	SILICON DIOXIDE, isotopic, solid	74
IRMM-311	Genomic DNA of <i>Bacillus Licheniformis</i> DSM 5749	47
IRMM-312	Genomic DNA of <i>Bacillus Subtilis</i> DSM 5750	47
IRMM-315	4-DEOXYNIVALENOL in acetonitrile	27
IRMM-316	NIVALENOL in acetonitrile	27
IRMM-351	ESCHERICHIA COLI 0157 (NCTC 12900)	48
IRMM-352	SALMONELLA ENTERITIDIS (NCTC 12694)	48
IRMM-354	CANDIDA ALBICANS (NCPP 3179)	48
IRMM-355	ENTEROCOCCUS FAECALIS (CIP 106877)	49
IRMM-3702	ZINC-64 spike, nitrate solution	76
IRMM-435	PHARMACEUTICAL GLASS	58
IRMM-440	RESIN-BONDED FIBRE BOARD (thermal conductivity)	59
IRMM-441	n-HEPTANE (purity)	65, 70
IRMM-442	ISOOCTANE (purity)	65, 70
IRMM-443-1	EUROSOIL (adsorption coefficients and pH)	25
IRMM-443-2	EUROSOIL (adsorption coefficients and pH)	25
IRMM-443-3	EUROSOIL (adsorption coefficients and pH)	25
IRMM-443-4	EUROSOIL (adsorption coefficients and pH)	25
IRMM-443-5	EUROSOIL (adsorption coefficients and pH)	25
IRMM-443-7	EUROSOIL (adsorption coefficients and pH)	25
IRMM-447	Genomic DNA of <i>Listeria Monocytogenes</i>	49
IRMM-448	Genomic DNA of <i>Campylobacter Jejuni</i>	49
IRMM-449	Genomic DNA of <i>Escherichia coli</i>	49
IRMM-468	THYROXINE (T4)	53
IRMM-469	3,3',5 TRIIODOTHYRONINE (T3)	53
IRMM-471	CEMENTITE GRAINS IN CARBURISED PURE IRON	72
IRMM-481	Peanut Test Material Kit	51
IRMM-521	Ni	67
IRMM-522	Cu	67
IRMM-523	Al	67
IRMM-524	Fe	67
IRMM-525	Nb	67
IRMM-526	Nb	67
IRMM-527R	Al-0.1 % Co	67
IRMM-528R	Al-1.0 % Co	67
IRMM-529	Rh	67
IRMM-530R	Al-0.1 % Au	67
IRMM-531	Ti	67
IRMM-532	Al-0.01 % Co	67
IRMM-533	Al-0.1 % Ag	67
IRMM-534	Al-2.0 % Sc	67
IRMM-540R	OXIDE GLASS (15 ppm U)	71
IRMM-541	OXIDE GLASS (50 ppm U)	71
IRMM-610	BORON-10 spike, aqueous solution	75
IRMM-611	BORON (natural) spike, aqueous solution	75
IRMM-615	LITHIUM-6 spike, chloride solution	75
IRMM-618	RUBIDIUM-87 spike, nitrate solution	75
IRMM-619	RUBIDIUM (natural) spike, nitrate solution	75
IRMM-620	IRON-57 spike, chloride solution	75
IRMM-621	CADMIUM-111 spike, nitrate solution	75

IRMM-622	CADMIUM-111 spike, nitrate solution	75
IRMM-624	CHROMIUM-50 spike, chloride solution	75
IRMM-625	CHROMIUM (natural) spike, chloride solution	75
IRMM-632	COPPER-65 spike, nitrate solution	75
IRMM-634	IRON (natural) spike, chloride solution	75
IRMM-635	STRONTIUM-84 spike, nitrate solution	75
IRMM-643	SULPHUR-32 spike, nitrate solution	76
IRMM-644	SULPHUR-32 spike, nitrate solution	76
IRMM-645	SULPHUR-32 spike, nitrate solution	76
IRMM-646	SULPHUR-34 spike, nitrate solution	76
IRMM-651	ZINC-64 spike, nitrate solution	76
IRMM-652	ZINC-64 spike, nitrate solution	76
IRMM-653	ZINC-67 spike, nitrate solution	76
IRMM-654	ZINC-68 spike, nitrate solution	76
IRMM-801	COCOA BUTTER	43
IRMM-804	RICE FLOUR	41
STA-003m	TETRAMETHYLUREA	26
VDA 001-004	POLYETHYLENE (40, 75, 200, 400 mg/kg Cd)	72

## ALPHABETICAL LIST

DESIGNATION	ERM / CRM	PAGE NO.
$\gamma$ -GLUTAMYLTRANSFERASE (catalytic concentration)	ERM-AD452/IFCC	57
10-AZABENZO[a]PYRENE (purity)	BCR-092	1
1-METHYLBENZ[a]ANTHRACENE (purity)	BCR-093R	1
1-METHYLCHRYSENE (purity)	BCR-077R	1
1-NITRONAPHTALENE (purity)	BCR-306	2
1-NITROPYRENE (purity)	BCR-305	2
2,2',3,4,4',5,5'-HEPTACHLOROBIPHENYL (IUPAC No. 180) (purity)	BCR-298	2
2,2',3,4,4',5'-HEXACHLOROBIPHENYL (IUPAC No. 138) (purity)	BCR-296	2
2,2',4,4',5,5'-HEXACHLOROBIPHENYL (IUPAC No. 153) (purity)	BCR-297	2
2,2',5,5'-TETRACHLOROBIPHENYL (IUPAC No. 52) (purity)	BCR-293	2
2,3,3'-TRICHLOROBIPHENYL (IUPAC No. 20) (purity)	BCR-290	2
2,4,4'-TRICHLOROBIPHENYL (IUPAC No. 28) (purity)	BCR-291	2
2,4'-DICHLOROBIPHENYL (IUPAC No. 8) (purity)	BCR-289	2
2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE	BCR-551	52
2,4-DINITROPHENYLHYDRAZONES in ACETONITRILE (blank)	BCR-552	52
2-METHYLCHRYSENE (purity)	BCR-078R	1
2-NITRO-7-METHOXYNAPHTHO[2,1-b]FURAN (purity)	BCR-312	2
2-NITRONAPHTALENE (purity)	BCR-307	2
3 REFERENCE ETHANOLS (H, M, L) (for SNIF-NMR)	BCR-123A, B	25, 73
3,3',5 TRIIODOTHYRONINE (T3)	IRMM-469	53
3-METHYLCHRYSENE (purity)	BCR-079R	1
3-NITROFLUORANTHENE (purity)	BCR-310	2
4-DEOXYNIVALENOL in acetonitrile	IRMM-315	27
4H-CYCLOPENTA[def]PHENANTHREN-4-ONE (purity)	BCR-338	2
4-METHYLCHRYSENE (purity)	BCR-080R	1
5-METHYLCHRYSENE (purity)	BCR-081R	1
6-METHYLCHRYSENE (purity)	BCR-082R	1
6-NITROBENZO[a]PYRENE (purity)	BCR-311	2
6-NITROCHRYSENE (purity)	BCR-309	2
7H-DIBENZO (c,g) CARBAZOLE (purity)	BCR-266	1
9-NITROANTHRACENE (purity)	BCR-308	2
ACETALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-547	52
ACETONE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-549	52
ACROLEIN-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-548	52
AFLATOXIN B1 IN ACETONITRILE	ERM-AC057	27
AFLATOXIN B2 IN ACETONITRILE	ERM-AC058	27
AFLATOXIN G1 IN ACETONITRILE	ERM-AC059	27
AFLATOXIN G2 IN ACETONITRILE	ERM-AC060	27
AFLATOXIN M1 STANDARD SOLUTION	BCR-423 (RM)	26
Al	IRMM-523	67
Al-0.01 % Co	IRMM-532	67
Al-0.1 % Ag	IRMM-533	67
Al-0.1 % Au	IRMM-530R	67
Al-0.1 % Co	IRMM-527R	67
Al-1.0 % Co	IRMM-528R	67
Al-2.0 % Sc	IRMM-534	67
ALANINE AMINOTRANSFERASE (catalytic concentration)	ERM-AD454/IFCC	57
ALPHA ALUMINA (0.10 m <sup>2</sup> /g) (nitrogen BET specific surface area)	BCR-169	62
ALPHA ALUMINA (1.05 m <sup>2</sup> /g) (nitrogen BET specific surface area)	BCR-170	62
ALPHA-AMYLASE (catalytic concentration)	IRMM/IFCC-456	57

ALUMINA (2.95 m <sup>2</sup> /g) (nitrogen BET specific surface area)	BCR-171	62
ANHYDROUS BUTTER FAT (tracers)	BCR-633	43
ANHYDROUS BUTTER FAT (triglycerides)	BCR-519	43
ANIMAL FEED (Organochlorine pesticides)	BCR-115	38
ANTHANTHRENE (purity)	BCR-091	1
ANTHRACITE (S)	BCR-334	70
ANTIMONY IMPLANTED IN SILICON	ERM-EG001	72
APPLE (dietary fibre)	ERM-BC516	46
AQUATIC PLANT (Cr)	BCR-596	13
AQUATIC PLANT (Lagarosiphon major) (trace elements)	BCR-060	13
ARTIFICIAL FOODSTUFF (major nutrients)	BCR-644	44
ARTIFICIAL FOODSTUFF (major nutrients)	BCR-645	44
ARTIFICIAL GROUND WATER (trace elements, low carbonate content)	BCR-617	18
ASPARTATE TRANSAMINASE (AST)	ERM-AD457/IFCC	57
BACILLUS CEREUS (number of colony forming particles)	BCR-528	47
BCR-ABL pDNA CALIBRANT	ERM-AD623	58
BEECH LEAVES (trace elements)	BCR-100	13
BEECH WOOD	BCR-683	24
BEEF-PORK FAT BLEND (fatty acid profile)	BCR-163	42
BEER (EtOH, low level)	BCR-651	44
BEER (EtOH, very low level)	BCR-652	44
BENZ[a]ACRIDINE (purity)	BCR-157	1
BENZ[a]ANTHRACENE (purity)	BCR-271	1
BENZ[c]ACRIDINE (purity)	BCR-158	1
BENZO[a]FLUORANTHENE (purity)	BCR-097	1
BENZO[a]FLUORENONE (purity)	BCR-342	2
BENZO[b]CHRYSENE (purity)	BCR-046	1
BENZO[b]FLUORANTHENE (purity)	BCR-047	1
BENZO[b]NAPHTHO (1,2-d) FURAN (purity)	BCR-340	2
BENZO[b]NAPHTHO (2,1-d) FURAN (purity)	BCR-341	2
BENZO[b]NAPHTHO[1,2-d]THIOPHENE (purity)	BCR-137R	1
BENZO[b]NAPHTHO[2,3-d]THIOPHENE (purity)	BCR-136R	1
BENZO[c,d]PYREN-6-ONE (purity)	BCR-339	2
BENZO[c]CHRYSENE (purity)	BCR-140	1
BENZO[c]PHENANTHRENE (purity)	BCR-134	1
BENZO[e]PYRENE (purity)	BCR-050	1
BENZO[ghi]FLUORANTHENE (purity)	BCR-139	1
BENZO[ghi]PERYLENE (purity)	BCR-052	1
BENZO[jj]FLUORANTHENE (purity)	BCR-049	1
BENZO[k]FLUORANTHENE (purity)	BCR-048R	1
BLADDERWRACK ( <i>Fucus vesiculosus</i> ) (trace elements)	ERM-CD200	15
BORIC ACID, isotopic, solid	IRMM-011	74
BORON (natural) spike, aqueous solution	IRMM-611	75
BORON-10 spike, aqueous solution	IRMM-610	75
BOVINE BLOOD (Pb, Cd)	ERM-CE195	54
BOVINE BLOOD (Pb, Cd)	ERM-CE196	54
BOVINE BLOOD LYSATE (haemiglobincyanide)	BCR-522	56
BOVINE EYE (CLENBUTEROL BLANK)	BCR-673	50
BOVINE EYE (CLENBUTEROL POSITIVE)	BCR-674	50
BOVINE LIVER (CLENBUTEROL BLANK)	BCR-648	50
BOVINE LIVER (CLENBUTEROL POSITIVE)	BCR-649	50
BOVINE LIVER (trace elements)	BCR-185R	40
BOVINE LIVER (trenbolone blank and positive)	BCR-474-5	50
BOVINE MUSCLE (diethylstilboestrol blank)	BCR-412	50

BOVINE MUSCLE (diethylstilboestrol positive)	BCR-411	50
BOVINE MUSCLE (trace elements)	ERM-BB184	40
BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol)	ERM-BB389	50
BOVINE URINE (diethylstilboestrol, dienoestrol and hexoestrol) (blank)	ERM-BB386	50
BOVINE URINE (clenbuterol and salbutamol)	BCR-502	49
BOVINE URINE (clenbuterol and salbutamol)	BCR-503	49
BOVINE URINE (clenbuterol and salbutamol)	BCR-504	49
BOVINE URINE (dienoestrol blank)	BCR-387	49
BOVINE URINE (dienoestrol positive)	BCR-390 (RM)	49
BOVINE URINE (diethylstilboestrol blank)	BCR-386	49
BOVINE URINE (hexoestrol blank)	BCR-388	49
BOVINE URINE (hexoestrol positive)	BCR-391	49
BRAN BREAKFAST CEREAL (dietary fibre)	ERM-BD518	46
BROWN BREAD (trace elements)	BCR-191	41
BROWN COAL	ERM-EF412	60
BRUSSELS SPROUT (vitamins)	BCR-431	44
BUTTER FAT	BCR-632	43
CADMIUM-111 spike, nitrate solution	IRMM-621	75
CADMIUM-111 spike, nitrate solution	IRMM-622	75
CALCAREOUS SOIL	ERM-CC690	10
CALCIUM AMMONIUM NITRATE FERTILIZER (composition)	BCR-178	65
CALCIUM-41 isotopic, nitrate solution (set of 8 units)	ERM-AE701	74
CANDIDA ALBICANS (NCPF 3179)	IRMM-354	48
CARROT (dietary fibre)	ERM-BC515	46
CEMENTITE GRAINS IN CARBURISED PURE IRON	IRMM-471	72
CHANNEL SEDIMENT (trace elements)	BCR-320R	11
CHARPY SPECIMENS 120 J (impact toughness)	ERM-FA016	62
CHARPY SPECIMENS 150 J (impact toughness)	ERM-FA415	62
CHARPY SPECIMENS 80 J (impact toughness)	ERM-FA015	62
CHARPY SPECIMENS Low Energy 0°C (impact toughness)	ERM-FA013	62
CHARPY SPECIMENS Low Energy 20°C (impact toughness)	ERM-FA013	62
CHLORIDE (natural) spike, chloride solution	ERM-AE641	76
CHLORIDE-37 spike, chloride solution	ERM-AE642	76
CHLORINATED HYDROCARBONS ON TENAX	BCR-555	53
CHROMIUM (natural) spike, chloride solution	IRMM-625	75
CHROMIUM, isotopic, chloride solution	IRMM-012	74
CHROMIUM-50 spike, chloride solution	IRMM-624	75
CHRYSENE (purity)	BCR-269	1
CHUB (PCBs)	BCR-719	24
CLAY (F)	BCR-461	70
COAL (F)	BCR-460	70
COASTAL SEAWATER (Hg)	BCR-579	17
COASTAL SEDIMENT (butyltins)	BCR-462	20
COCOA BUTTER	IRMM-801	43
COCONUT OIL (PAH blank)	BCR-459	36
COCONUT OIL (PAH doped)	BCR-458	36
COD LIVER OIL (Organochlorine pesticides)	BCR-598	37
COD LIVER OIL (PCBs)	BCR-349	37
COKING STEAM COAL (S)	BCR-333	70
COLLOIDAL SILICA	ERM-FD100	64
COLLOIDAL SILICA	ERM-FD304	64
COMMON WHEAT FLOUR (properties)	BCR-563	46
COMPOUND FEED (aflatoxin B1 blank)	BCR-375	39
COMPOUND FEEDINGSTUFF (high level)	ERM-BE376	39

COMPOUND FEEDINGSTUFF (very low level)	ERM-BE375	39
CONTINUOUS CAST COPPER (O)	BCR-058	67
COPPER (natural) spike, nitrate solution	ERM-AE633	75
COPPER (O)	BCR-054R	67
COPPER (S, P)	BCR-017A, B	67
COPPER ALLOYS	BCR-691	69
COPPER-63, nitrate solution	ERM-AE647	76
COPPER-65 spike, nitrate solution	IRMM-632	75
CORONENE (purity)	BCR-272	1
CORTISOL REFERENCE SERUM PANEL	ERM-DA451/IFCC	53
CREATINE KINASE (CK-MB iso-enzyme) (catalytic concentration)	ERM-AD455/IFCC	57
CREATININE (interfering substances)	BCR-573i	57
Cu	IRMM-522	67
DAIRY FEED (nutritional properties)	BCR-708	45
DEFATTED PEANUT MEAL (aflatoxin B1, blank)	BCR-262R	39
DEFATTED PEANUT MEAL (aflatoxin B1, high level)	BCR-264	39
DEFATTED PEANUT MEAL (aflatoxin B1, medium level)	BCR-263R	39
DIBENZ[a,c]ACRIDINE (91mylas)	BCR-155	1
DIBENZ[a,h]ACRIDINE (purity)	BCR-153R	1
DIBENZ[a,l]ACRIDINE (91mylas)	BCR-152	1
DIBENZ[a,j]ACRIDINE (purity)	BCR-154	1
DIBENZ[a,c]ANTHRACENE (purity)	BCR-094	1
DIBENZ[a,j]ANTHRACENE (purity)	BCR-095	1
DIBENZ[c,h]ACRIDINE (purity)	BCR-156R	1
DIBENZO[a,e]FLUORANTHENE (91mylas)	BCR-265	1
DIBENZO[a,h]PYRENE (91mylas)	BCR-159	1
DIBENZO[a,e]PYRENE (91mylas)	BCR-133	1
DIBENZO[a,h]ANTHRACENE (purity)	BCR-138	1
DIBENZO[a,l]PYRENE (91mylas)	BCR-096	1
DIBENZO[b,d]FURAN (91mylas)	BCR-337	2
ELECTROLYTIC COPPER (trace elements)	BCR-074A	69
ELECTROLYTIC TOUGH PITCH COPPER (O)	BCR-022A, B	67
ELECTROLYTICALLY REFINED LEAD (trace elements)	BCR-286A, B	68
ENTEROCOCCUS FAECALIS (CIP 106877)	IRMM-355	49
ESCHERICHIA COLI 0157 (NCTC 12900)	IRMM-351	48
ESTUARINE SEDIMENT	BCR-667	10
ESTUARINE SEDIMENT (Hg, methylmercury)	ERM-CC580	11, 20
ESTUARINE SEDIMENT (trace elements)	BCR-277R	11
ESTUARINE WATER (trace elements)	BCR-505	17
EUROSOIL (adsorption coefficients and Ph)	IRMM-443-1	25
EUROSOIL (adsorption coefficients and Ph)	IRMM-443-2	25
EUROSOIL (adsorption coefficients and Ph)	IRMM-443-3	25
EUROSOIL (adsorption coefficients and Ph)	IRMM-443-4	25
EUROSOIL (adsorption coefficients and Ph)	IRMM-443-5	25
EUROSOIL (adsorption coefficients and Ph)	IRMM-443-7	25
EWES'/GOATS' CURD (for adulteration with cows' milk) 0 and 1 % cows' milk	BCR-599	51
FAUJASITE TYPE ZEOLITE (micropore volume and width)	BCR-704	63
Fe	IRMM-524	67
FINE DUST (PM <sub>10</sub> -like) (elements)	ERM-CZ120	13
FINE DUST (PM <sub>10</sub> -like) (PAHs)	ERM-CZ100	21
FISH MUSCLE (trace elements)	ERM-BB422	40
FISH OIL	ERM-BB350	37
FLAME COAL (S)	BCR-335	70
FLUORANTHENE (purity)	BCR-160R	1

FLY ASH (LOW LEVEL) (PCDDs and PCDFs)	BCR-615	23
FLY ASH (PCDDs and PCDFs)	BCR-490	23
FLY ASH (trace elements)	BCR-176R	12
FLY ASH FROM PULVERISED COAL (trace elements)	BCR-038	12
FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-546	52
FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter	BCR-553	53
FORMALDEHYDE-2,4-DINITROPHENYLHYDRAZONE on filter (blank)	BCR-554	53
FRESH WATER (nitrate, high level)	BCR-480	17
FRESH WATER (nitrate, low level)	BCR-479	17
FRESHWATER HARBOUR SEDIMENT (PAHs)	BCR-535	22
FRESHWATER HARBOUR SEDIMENT (PCBs)	BCR-536	22
FRESHWATER SEDIMENT (butyltin and phenyltin compounds)	BCR-646	20
FULL FAT SOYA (dietary fibre)	ERM-BC517	46
FURNACE COKE	ERM-EF413	60
GAS OIL (0.0203 % S)	ERM-EF672	70
GAS OIL (0.0452 % S)	ERM-EF671	70
GAS OIL (0.1019 % S)	ERM-EF104	70
GAS OIL (0.363 % S)	BCR-105	70
GAS OIL (0.502 % S)	BCR-106	70
GAS OIL (1.040 % S)	BCR-107	70
GAS OIL (CFPP 6°C)	BCR-395	59
GAS OIL (CFPP and CP)	ERM-FC395k	59
GAS OIL (Solvent Yellow 124)	ERM-EF317	70
GAS OIL (Solvent Yellow 124)	ERM-EF318	70
GENETICALLE MODIFIED AM04-1020 POTATO	ERM-BF430	34
GENETICALLE MODIFIED AV43-6-G7 POTATO	ERM-BF431	34
GENETICALLE MODIFIED DAS-40278-9 MAIZE	ERM-BF433	35
GENETICALLE MODIFIED DAS-44406-6 SOYA	ERM-BF436	36
GENETICALLY MODIFIED 1507 MAIZE	ERM-BF418	31
GENETICALLY MODIFIED 281-24-236 X 3006-210-23 COTTON SEED	ERM-BF422	32
GENETICALLY MODIFIED 3272 MAIZE	ERM-BF420	31
GENETICALLY MODIFIED 59122 MAIZE	ERM-BF424	32
GENETICALLY MODIFIED 73496 RAPESEED	ERM-BF434	35
GENETICALLY MODIFIED 98140 MAIZE	ERM-BF427	33
GENETICALLY MODIFIED Bt-11 MAIZE	ERM-BF412	28
GENETICALLY MODIFIED Bt-176 MAIZE	ERM-BF411	28
GENETICALLY MODIFIED EH92-527-1 POTATO	ERM-BF421	31
GENETICALLY MODIFIED GA21 MAIZE	ERM-BF414	29
GENETICALLY MODIFIED GHB119 COTTON	ERM-BF428	34
GENETICALLY MODIFIED H7-1 SUGAR BEET	ERM-BF419	31
GENETICALLY MODIFIED MIR604 MAIZE	ERM-BF423	32
GENETICALLY MODIFIED MON 810 MAIZE	ERM-BF413k	29
GENETICALLY MODIFIED MON 863 MAIZE	ERM-BF416	30
GENETICALLY MODIFIED MON 863 x MON 810 MAIZE	ERM-BF417	30
GENETICALLY MODIFIED NK603 MAIZE	ERM-BF415	30
GENETICALLY MODIFIED PH05-026-0048 POTATO	ERM-BF435	35
GENETICALLY MODIFIED ROUNDUP READY SOYA	ERM-BF410k	28
GENETICALLY MODIFIED SOYA 305423	ERM-BF426	33
GENETICALLY MODIFIED SOYA 356043	ERM-BF425	32
GENETICALLY MODIFIED SOYA DAS-68416-4	ERM-BF432	35
GENETICALLY MODIFIED T304-40 COTTON	ERM-BF429	34
Genomic DNA of <i>Bacillus Licheniformis</i> DSM 5749	IRMM-311	47
Genomic DNA of <i>Bacillus Subtilis</i> DSM 5750	IRMM-312	47
Genomic DNA of <i>Campylobacter Jejuni</i>	IRMM-448	49

Genomic DNA of Escherichia coli	IRMM-449	49
Genomic DNA of Listeria Monocytogenes	IRMM-447	49
GLASS (trace elements)	BCR-664	71
GLASS-CERAMIC	BCR-724A-E	59
GLUTARALDEHYDE-2,4-DINITROPHENYLHYDRAZONE (purity)	BCR-550	52
GROUND WATER	ERM-CA615	18
GROUND WATER	ERM-CA616	18
GROUND WATER (Br, high level)	BCR-611	17
GROUND WATER (Br, low level)	BCR-612	17
GROUND WATER (trace elements, high level)	BCR-610	18
GROUND WATER (trace elements, low level)	BCR-609	18
HAEMOGLOBIN HbA0	IRMM/IFCC-467	57
HARD COAL	ERM-EF411	60
HARICOTS BEANS (dietary fibre)	ERM-BC514	46
HARICOTS VERTS (major nutrients)	BCR-383	42, 44
HAY POWDER (elements)	BCR-129	13
HERRING (PCBs)	BCR-718	24
HIGH VOLATILE INDUSTRIAL COAL (S)	BCR-332	70
HIGH VOLATILE STEAM COAL (S)	BCR-336	70
HUMAN ADENOSINE DEAMINASE (ADA 1)	BCR-647	57
HUMAN APOLIPOPROTEIN A I (mass concentration)	BCR-393	55
HUMAN BLOOD (Pb, Cd)	BCR-634	54
HUMAN BLOOD (Pb, Cd)	BCR-635	54
HUMAN BLOOD (Pb, Cd)	BCR-636	54
HUMAN HAEMOLYSATE (glycated haemoglobin (HbA1c))	BCR-405 (RM)	56
HUMAN HAIR (trace elements)	ERM-DB001	55
HUMAN PANCREATIC LIPASE (from pancreatic juice)	BCR-693	57
HUMAN PANCREATIC LIPASE (recombinant)	BCR-694	57
HUMAN PROSTATIC ACID PHOSPHATASE (catalytic concentration)	BCR-410	57
HUMAN SERUM (17 $\beta$ -ESTRADIOL, high level)	BCR-578	54
HUMAN SERUM (17 $\beta$ -ESTRADIOL, low level)	BCR-576	54
HUMAN SERUM (17 $\beta$ -ESTRADIOL, medium level)	BCR-577	54
HUMAN SERUM (Al, Se, Zn)	BCR-637	55
HUMAN SERUM (Al, Se, Zn)	BCR-638	55
HUMAN SERUM (Al, Se, Zn)	BCR-639	55
HUMAN SERUM (Ca, Mg, Li)	BCR-304	55
HUMAN SERUM (cortisol spiked)	ERM-DA193	54
HUMAN SERUM (cortisol unspiked)	ERM-DA192	54
HUMAN SERUM (CRP)	ERM-DA474/IFCC	56
HUMAN SERUM (cystatin C)	ERM-DA471/IFCC	56
HUMAN SERUM (high creatinine)	BCR-575	56
HUMAN SERUM (high progesterone)	BCR-348R	54
HUMAN SERUM (low creatinine)	BCR-573	56
HUMAN SERUM (medium creatinine)	BCR-574	56
HUMAN SERUM (progesterone)	ERM-DA347	54
HUMAN SERUM (proteins)	ERM-DA470k/IFCC	56
HUMAN THYROGLOBULIN (Tg) (mass concentration)	BCR-457	55
INDENO[1,2,3-cd]FLUORANTHENE (purity)	BCR-267	1
INDUSTRIAL SANDY SOIL (PCDDs, PCDFs)	BCR-529	22
INDUSTRIAL SOIL (PAHs)	BCR-524	22
INDUSTRIAL SOIL (PCBs)	BCR-481	22
IRON (natural) spike, chloride solution	IRMM-634	75
IRON-57 spike, chloride solution	IRMM-620	75
ISOOCTANE (purity)	IRMM-442	65, 70

ISOTOPE RATIOS IN ABSOLUTE ALCOHOL	BCR-656	25, 73
ISOTOPE RATIOS IN ALCOHOLIC SOLUTION	BCR-660	26, 73
ISOTOPE RATIOS IN GLUCOSE	BCR-657	26, 73
ISOTOPE RATIOS IN SYNTHETIC WINE	BCR-658	26, 73
ISOTOPE RATIOS IN SYNTHETIC WINE	BCR-659	26, 73
LAKE SEDIMENT (trace elements)	BCR-280R	11
LAKE SEDIMENT (trace elements)	BCR-701	20
LATEX SPHERES (particle diameter 2 microns)	BCR-165	53
LATEX SPHERES (particle diameter 4.8 microns)	BCR-166	53
LATEX SPHERES (particle diameter 9.6 microns)	BCR-167	53
LEAD GLASS (composition/refractive index)	BCR-126A	66
LEAD WITH ADDED IMPURITIES (trace elements)	BCR-288B	68
LEMNA MINOR (aquatic plant)	BCR-670	15
LICHEN (trace elements)	BCR-482	15
LIGHT SANDY SOIL (trace elements)	BCR-142R	10
LIMESTONE POWDERS (for shear testing)	BCR-116	60
LINDE TYPE A ZEOLITE (mircopore volume and width)	BCR-705	63
LITHIUM CARBONATE, isotopic, solid	IRMM-016	74
LITHIUM-6 spike, chloride solution	IRMM-615	75
LOAM SOIL	ERM-CC141	10
LOW VOLATILE STEAM COAL (S)	BCR-331	70
LUNG TISSUE (asbestos fibres)	BCR-665	58
LUNG TISSUE (asbestos fibres)	BCR-666	58
MAGNESIUM (natural) spike, nitrate solution	ERM-AE637	75
MAGNESIUM-26 spike, nitrate solution	ERM-AE638	75
MAIZE	ERM-BC716	40
MAIZE	ERM-BC717	40
MAIZE FLOUR (deoxynivalenol blank)	BCR-377	39
MARGARINE (vitamins)	BCR-122	44
MERCURY (natural) spike, chloride solution	ERM-AE639	75
MERCURY-202 spike, chloride solution	ERM-AE640	76
Mg, isotopic, nitrate solution	IRMM-009	74
MICROCRYSTALLINE CELLULOSE (water content above saturated solutions)	BCR-302	63
MILK POWDER (oxytetracycline)	ERM-BB492	50
MILK POWDER (oxytetracycline) (blank)	ERM-BB493	50
MILK POWDER (oxytetracycline)	BCR-492	50
MILK POWDER (oxytetracycline) (blank)	BCR-493	50
MILK POWDER (PCDDs, PCDFs)	BCR-607	38
MIXED VEGETABLES (vitamins)	BCR-485	44
MOROCCAN PHOSPHATE ROCK (trace elements)	BCR-032	66, 71
MULLITE (lattice spacing, other parameters)	BCR-301 (RM)	63
MUSSEL (dc-saxitoxin)	BCR-543	40
MUSSEL TISSUE	BCR-668	16
MUSSEL TISSUE	BCR-682	24
MUSSEL TISSUE (butyltins)	ERM-CE477	21
MUSSEL TISSUE (elements)	ERM-CE278k	16
NATURAL MILK POWDER (PCBs )	BCR-450	36
NATURAL MILK POWDER (pesticides)	BCR-187	38
NATURAL PORK FAT (blank)	ERM-BB444	36
Nb	IRMM-525	67
Nb	IRMM-526	67
n-HEPTANE (purity)	IRMM-441	65, 70
Ni	IRMM-521	67
NICKEL (N, O)	BCR-099	67

NIMONIC 75 FOR CREEP TESTING	BCR-425	61
NIMONIC 75 FOR TENSILE PROPERTIES	BCR-661	61
NIVALENOL in acetonitrile	IRMM-316	27
ORGANIC-RICH SOIL (extractable elements)	BCR-700	19
OXIDE GLASS (15 ppm U)	IRMM-540R	71
OXIDE GLASS (50 ppm U)	IRMM-541	71
PAHs IN ACETONITRILE / TOLUENE	ERM-AC213	9
PCB STANDARD SOLUTION	BCR-365	2
PEANUT BUTTER (aflatoxins low level)	BCR-385R	39
PEANUT BUTTER (aflatoxins very low level)	BCR-401R	39
Peanut Test Material Kit	IRMM-481	51
PETROL	ERM-EF211	70
PHARMACEUTICAL GLASS	IRMM-435	58
PICENE (purity)	BCR-168	1
PIG FEED (nutritional properties)	BCR-709	45
PIG KIDNEY (CTC free)	BCR-706	50
PIG KIDNEY (CTC incurred)	BCR-707	50
PIG KIDNEY (trace elements)	ERM-BB186	40
PIG LIVER (CTC free)	BCR-695	50
PIG LIVER (CTC incurred)	BCR-696	50
PIG LIVER (vitamins)	BCR-487	44
PIG MUSCLE (CTC free)	BCR-697	50
PLANKTON (trace elements)	BCR-414	14
PLASMID DNA FRAGMENTS OF 356043 SOYBEAN	ERM-AD425	33
PLASMID DNA FRAGMENTS OF 98140 MAIZE	ERM-AD427	34
PLASMID DNA FRAGMENTS OF MON 810 MAIZE	ERM-AD413	29
PLASMID DNA FRAGMENTS OF NK603 MAIZE	ERM-AD415	30
PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film A)	BCR-537	46
PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film B)	BCR-538	46
PLASTIC FILM (OVERALL MIGRATION IN OLIVE OIL (film C)	BCR-539	46
PLATINUM, isotopic, metal	IRMM-010	74
POLYCHLORODIBENZO-P-DIOXINS (PCDD) AND POLYCHLORODIBENZOFURANS (PCDFS)	BCR-614	3
POLYETHYLENE (40, 75, 200, 400 mg/kg Cd)	VDA 001-004	72
POLYETHYLENE (high level)	ERM-EC681k	72
POLYETHYLENE (LDPE)	ERM-EC590	72
POLYETHYLENE (low level)	ERM-EC680k	72
POLYPROPYLENE (PP)	ERM-EC591	72
PORCINE MUSCLE (chloramphenicol blank)	BCR-444	50
PORCINE MUSCLE (chloramphenicol positive)	BCR-445	50
PORK FAT (pesticides)	ERM-BB430	37
PORK MUSCLE	ERM-BB124	51
PORK MUSCLE	ERM-BB130	50
PORK MUSCLE	ERM-BB384	42, 44
POTASSIUM CHLORIDE FERTILIZER (elemental composition)	BCR-113	65
POTASSIUM SULPHATE FERTILIZER (elemental composition)	BCR-114	65
PROSTATE SPECIFIC ANTIGEN (protein mass)	BCR-613	56
PURIFIED HUMAN ALFAFOETOPROTEIN (protein mass)	BCR-486	55
PYRENE (purity)	BCR-177R	1
QUARTZ (1.20 – 20.00 microns)	BCR-070	62
QUARTZ (2.50 m <sup>2</sup> /g) (nitrogen BET specific surface area)	BCR-172	62
QUARTZ (particle size 0.35 – 3.50 microns)	BCR-066	62
QUARTZ (particle size 14 – 90 microns)	BCR-069	62
QUARTZ (particle size 1400 – 5000 microns)	BCR-132	62
QUARTZ (particle size 160 – 630 microns)	BCR-068	62

QUARTZ (particle size 2.40 – 32.00 microns)	BCR-067	62
QUARTZ (particle size 480 – 1800 microns)	BCR-131	62
QUARTZ (particle size 50 – 220 microns)	BCR-130	62
RAPESEED (colza) (S, total glucosinolate, high level)	ERM-BC367	38
RAPESEED (colza) (S, total glucosinolate, low level)	ERM-BC366	38
RAPESEED (colza) (S, total glucosinolate, medium level)	ERM-BC190	38
RAPESEED (oil, moisture, volatiles)	BCR-446	45
RAPESEED (oil, moisture, volatiles)	BCR-447	45
RESIN-BONDED FIBRE BOARD (thermal conductivity)	IRMM-440	59
Rh	IRMM-529	67
RICE (As species)	ERM-BC211	21
RICE FLOUR	IRMM-804	41
RICE FLOUR (96mylase, low level)	BCR-465	44
RICE FLOUR (96mylase, medium level)	BCR-466	44
RICE FLOUR (96mylase, high level)	BCR-467	44
RIVER SEDIMENT (extractable phosphorous)	BCR-684	19
ROAD DUST (trace elements)	BCR-723	12
RUBIDIUM (natural) spike, nitrate solution	IRMM-619	75
RUBIDIUM-87 spike, nitrate solution	IRMM-618	75
RYE FLOUR	ERM-BC381	42, 44
RYE GRASS	ERM-CD281	14
SALMON TISSUE	BCR-725	51
SALMONELLA ENTERITIDIS (NCTC 12694)	IRMM-352	48
SAXITOXIN IN ACETIC ACID	BCR-663	26
SCRATCH TESTING	BCR-692	61
SEWAGE SLUDGE (Cr)	BCR-597	11
SEWAGE SLUDGE (industrial origin) (trace elements)	BCR-146R	11
SEWAGE SLUDGE (mixed origin) (trace elements)	BCR-145R	11
SEWAGE SLUDGE (PCDDs and PCDFs)	BCR-677	23
SEWAGE SLUDGE AMENDED (terra rossa) SOIL (trace elements)	BCR-484	19
SEWAGE SLUDGE AMENDED SOIL (trace elements)	BCR-143R	10
SEWAGE SLUDGE AMENDED SOIL (trace elements)	BCR-483	19
SILICON DIOXIDE, isotopic, solid	IRMM-018a	74
SILICON, isotopic, Si single crystal	IRMM-017	74
SIMULATED RAINWATER (major components)	ERM-CA408	17
SINGLE CELL PROTEIN (major elements)	BCR-273	41
SINGLE CELL PROTEIN (trace elements)	BCR-274	41
SiO <sub>2</sub> /Si (10,20,30 nm)	BCR-564	63
SKIM MILK POWDER	BCR-685	45
SKIM MILK POWDER (major and trace elements)	BCR-063R	15
SKIMMED MILK POWDER (trace elements)	ERM-BD150	40
SKIMMED MILK POWDER (trace elements)	ERM-BD151	40
SOYA-MAIZE OIL BLEND (fatty acid profile)	BCR-162R	42
SPIKED MILK POWDER (pesticides)	BCR-188	38
SPIKED PORK FAT (low level)	ERM-BB446	36
SPIKED PORK FAT (very low level)	ERM-BB445	36
STRONTIUM-84 spike, nitrate solution	IRMM-635	75
SULPHUR-32 spike, nitrate solution	IRMM-643	76
SULPHUR-32 spike, nitrate solution	IRMM-644	76
SULPHUR-32 spike, nitrate solution	IRMM-645	76
SULPHUR-34 spike, nitrate solution	IRMM-646	76
SUPERPHOSPHATE (various parameters)	BCR-033	65
TANTALUM PENTOXIDE ON TANTALUM FOIL	BCR-261T	64
TETRAMETHYLUREA	STA-003m	26

THALLIUM (natural) spike, nitrate solution	ERM-AE649	76
THERMALLY REFINED LEAD (trace elements)	BCR-287A, B	68
THROMBOPLASTIN BOVINE (OBT/79) (prothrombin time)	ERM-AD148	58
THROMBOPLASTIN RABBIT (prothrombin time)	ERM-AD149	58
THYROXINE (T4)	IRMM-468	53
Ti	IRMM-531	67
Ti 6AL 4V ALLOY (O)	BCR-059A, B	67
TiAl6V4 (Al, V)	BCR-089	69
TIN ORE CONCENTRATE (Sn)	BCR-010	65
TITANIA (8.23 m <sup>2</sup> /g) (nitrogen BET specific surface area)	BCR-173	62
TITANIUM (H)	BCR-318	67
TITANIUM (impurities)	BCR-090A, B	69
TITANIUM (O, N)	BCR-024B, C	67
TOasted BREAD	ERM-BD273	52
TOMATO PASTE COLOUR REFERENCE TILE (colour values)	BCR-400	64
TRACE ELEMENTS IN WHITE CABBAGE	BCR-679	41
TRIPHENYLENE (purity)	BCR-270	1
TUNA FISH (total and methylmercury)	ERM-CE464	16, 21
TUNA FISH (total and methylmercury)	BCR-463	16, 21
TUNA FISH TISSUE (As species)	BCR-627	21
TUNGSTEN (0.18 m <sup>2</sup> /g) (nitrogen BET specific surface area)	BCR-175	62
TUNGSTEN CARBIDE POWDER (O)	BCR-102	67
UNALLOYED ZINC (disc) (trace elements)	BCR-326	68
UNALLOYED ZINC (disc) (trace elements)	BCR-327	68
UNALLOYED ZINC (trace elements)	ERM-EB322	68
UNALLOYED ZINC (trace elements)	ERM-EB323	68
UNALLOYED ZINC (trace elements)	ERM-EB324	68
UNALLOYED ZINC (trace elements)	ERM-EB325	68
UNALLOYED ZINC (trace elements)	BCR-321	68
URBAN DUST (trimethyllead)	BCR-605	20
UREA FERTILIZER (composition)	BCR-179	65
WASTE MINERAL OIL (high PCB level)	BCR-449	24
WASTE MINERAL OIL (low PCB level)	BCR-420	24
WASTE WATER	ERM-CA713	18
WELDING DUST LOADED ON FILTER (Cr VI, Cr)	BCR-545	21
WHEAT (ochratoxin A, blank)	BCR-471	39
WHEAT FLOUR	ERM-BC382	42, 44
WHEAT FLOUR (deoxynivalenol blank)	BCR-396	38
WHITE CLOVER (trace elements)	BCR-402	13
WHOLE MILK POWDER (aflatoxin M1, high level)	ERM-BD284	39
WHOLE MILK POWDER (aflatoxin M1, low level)	ERM-BD283	39
WHOLE MILK POWDER (aflatoxin M1, zero level)	ERM-BD282	39
WHOLE MILK POWDER (major nutrients)	BCR-380R	45
WHOLE MILK POWDER (vitamins)	ERM-BD600	44
WHOLEMEAL FLOUR (vitamins)	BCR-121	44
WINE (EtOH, low level)	BCR-653	44
ZEARALENONE IN ACETONITRILE	ERM-AC699	26
ZINC ORE CONCENTRATE (trace elements)	BCR-109	71
ZINC ORE CONCENTRATE (trace elements)	BCR-110	71
ZINC-64 spike, nitrate solution	IRMM-3702	76
ZINC-64 spike, nitrate solution	IRMM-651	76
ZINC-64 spike, nitrate solution	IRMM-652	76
Zinc-64, nitrate solution	IRMM-007/1-6	73
ZINC-67 spike, nitrate solution	IRMM-653	76

ZINC-68 spike, nitrate solution	IRMM-654	76
ZIRCALOY (C, N, O)	BCR-275	67
ZIRCALOY (C, N, O)	BCR-276	67
ZIRCALOY-4 (trace element impurities)	BCR-098	69
ZnAl4 (trace elements)	BCR-351	68
ZnAl4 (trace elements)	BCR-352	68
ZnAl4 (trace elements)	BCR-353	68
ZnAl4 (trace elements)	BCR-354	68
ZnAl4 (trace elements)	BCR-355	68
ZnAl4Cu1 (trace elements)	BCR-356	68
ZnAl4Cu1 (trace elements)	BCR-357	68
ZnAl4Cu1 (trace elements)	BCR-359	68
ZnAl4Cu1 (trace elements)	BCR-360	68
ZnAl4Cu1 (trace elements)	BCR-361	68

13/03/2014